Zeitschrift: Elemente der Mathematik

Herausgeber: Schweizerische Mathematische Gesellschaft

Band: 62 (2007)

Artikel: Smallest limited snakes

Autor: Szabó, Lázló / Ujváry-Menyhárt, Zoltán

DOI: https://doi.org/10.5169/seals-98914

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 01.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Elemente der Mathematik

Smallest limited snakes

László Szabó and Zoltán Ujváry-Menyhárt

László Szabó was born in Budapest in 1966. He studied mathematics and computer science at Eötvös Loránd University and received his Ph.D. in mathematics from the Hungarian Academy of Sciences in 1996. His research interests are discrete geometry and convexity.

Zoltán Ujváry-Menyhárt was born in Budapest in 1973. He studied mathematics at Eötvös Loránd University and computer science at Budapest University of Technology and Economics. His research interests are combinatorics and discrete geometry.

1 Introduction

A (topological) disk is a subset of the euclidean plane homeomorphic to the unit ball. If two disks have a common interior point then we say that the disks overlap. A sequence $\mathcal{C} = \langle C_1, \dots, C_n \rangle$ of mutually non overlapping congruent disks where $C_i \cap C_j \neq \emptyset$ if and only if $|i-j| \leq 1$ is called a snake. If the snake \mathcal{C} is not a proper subset of another snake of disks congruent to the members of \mathcal{C} then we say that the snake is limited.

We are concerned with the following question: What is the minimum number of mutually non overlapping congruent disks which can form a limited snake? Here we prove

Theorem. The minimum number of mutually non overlapping congruent disks which can form a limited snake is four.

Surprisingly, under the assumption of convexity the above problem seems to be much more complicated. Fig. 1 shows that six mutually non overlapping congruent copies of a

Auf einem Tisch legt man mit lauter gleichen Münzen eine "Münzschlange": an eine erste Münze anstossend legt man eine zweite, daran anstossend eine dritte usw. Bei diesem Legespiel kann eine Konfiguration entstehen, bei der man weder am Kopf noch am Schwanz der Schlange eine weitere Münze anschliessen kann, weil der Platz durch andere Münzen des Schlangenkörpers versperrt wird. Welches ist die kleinste Anzahl Münzen, bei der dies vorkommen kann? Die Autoren untersuchen und beantworten die entsprechende Frage, wenn man die runden Münzen durch eine beliebige einfach zusammenhängende beschränkte Menge der Euklidischen Ebene ersetzt. Das entsprechende Problem für konvexe beschränkte Mengen ist hingegen noch offen.

certain rectangle can form a limited snake. Do there exist convex disks whose less then six mutually non overlapping congruent copies could form limited snakes? We conjecture that the answer to this question is in the negative.





Fig. 1

Also, the problem of determining the minimum number of mutually non overlapping congruent copies of a given disk which can form a limited snake is very complicated. The only known result in this direction is that the minimum number of mutually non overlapping congruent balls which can form a limited snake is ten (see [1]).

For additional results on more restrictive variants of the snake problem, see [2, 3, 4, 5, 6, 7, 8].

2 Proof of the theorem

Fig. 2 shows that this minimum number is at most four.



To complete the proof we have to show that two or three mutually non overlapping congruent disks cannot form a limited snake. Let C be an arbitrary disk and let $\mathcal{C} = \langle C_1, \ldots, C_n \rangle$ be a limited snake consisting of disks congruent with C.

We start with the case n=2. First assume that $\operatorname{conv} C_1=\operatorname{conv} C_2$, i.e., the convex hulls of C_1 and C_2 coincide. If every boundary point of $\operatorname{conv} C_1$ belongs to C_1 , i.e., C_1 is convex, then C_1 and C_2 coincide, which is impossible. Thus there exists a boundary point of $\operatorname{conv} C_1$ which does not belong to C_1 . This point lies in the relative interior of a segment joining two extreme points, say A and B, of $\operatorname{conv} C_1$. Recall that a point of a disk is an extreme point of the disk if there exists no segment in the disk that contains the point in its relative interior. The points A and B are extreme points of C_1 and they can be joined with a path C_1 whose points different from C_1 and C_2 and they can be joined with a path C_2 whose points different from C_1 and C_2 and they can be joined with a path C_2 whose points different from C_2 and the segment C_3 and the segment C_4 and the segmen

Thus there exists a point P of C_1 which does not belong to conv C_2 . Then P can be strictly separated from C_2 by a line l. Let l' be the support line of C_1 which is parallel to l and does not separate C_1 and C_2 . Reflecting C_1 with respect to l' we obtain a third copy of C which forms with C_1 and C_2 a three element snake, a contradiction.

Now we turn to the case n=3. Let \overline{DE} be a diameter of C_1 and consider the stripe S_1 whose boundary lines, say l_1 and l_2 , go through D and E, respectively, and are perpendicular to \overline{DE} . If C_3 is not contained in S_1 then consider the support line l of C_3 which is parallel to l_1 and whose distance from S_1 is maximal. Without loss of generality we may assume that l_2 separates l and l_1 . Let F be a common point of C_3 and l. The disk C_2 cannot intersect both l and l_1 since the distance between the two lines is greater than the diameter of C_2 . Thus either reflecting C_1 with respect to l_1 or reflecting C_3 with respect to l we obtain a fourth copy of C which forms with C_1 , C_2 and C_3 a four element snake, a contradiction.

Thus C_3 lies in S_1 . Let \overline{GH} be a diameter of C_3 and consider the stripe S_3 whose boundary lines, say l_3 and l_4 , go through G and G, respectively, and are perpendicular to \overline{GH} . If C_1 is not contained in S_3 then repeating the previous argument we obtain a contradiction. Therefore C_1 lies in S_3 . If $S_1 = S_3$, i.e., $S_4 = S_4$ and $S_4 = S_4$ without loss of generality, then $S_4 = S_4$ are different points since $S_4 = S_4$ and $S_4 = S_4$ are different points since $S_4 = S_4$ and $S_4 = S_4$ and $S_4 = S_4$ and $S_4 = S_4$ and $S_4 = S_4$ are different stripes then their intersection is a parallelogram which contains both $S_4 = S_4$ are different stripes then their intersection is a parallelogram which contains both $S_4 = S_4$ are different stripes then their intersection is a parallelogram which contains both $S_4 = S_4$ and $S_4 = S_$

References

- [1] Bisztriczky, T.; Böröczky Jr, K.; Harborth, H.; Piepmeyer, L.: On the smallest limited snake of unit disks. *Geom. Dedicata* 40 (1991), 319–324.
- [2] Bisztriczky, T.; Harborth, H.: Smallest limited edge-to-edge snakes in Euclidean tessellations. *Congr. Numer.* 149 (2001), 155–159.

- [3] Böröczky Jr, K.; Soltan, V.: Smallest maximal snakes of translates of convex domains. *Geom. Dedicata* 54 (1995), 31–44.
- [4] Harborth, H.: Problem 45: Kleinste endliche Schlange. Math. Semesterber. 36 (1989), 269-270.
- [5] Harborth, H.; Szabó, L.; Ujváry-Menyhárt, Z.: Smallest limited vertex-to-vertex snakes of unit triangles. *Geom. Dedicata* 78 (1999), 171–181.
- [6] Heidelberg, R.; Stege, L.; Weiß, H.: Lösung zu Problem 45. Math. Semesterber. 38 (1991), 137–138.
- [7] Hering, F.: Beweis einer Vermutung von Heiko Harborth über Polyominos aus Quadraten. *Math. Semesterber.* 38 (1991), 223–237.
- [8] Szabó, L.; Ujváry-Menyhárt, Z.: Maximal facet-to-facet snakes of unit cubes. Beiträge Algebra Geom. 42 (2001), 203–217.

László Szabó and Zoltán Ujváry-Menyhárt Computer and Automation Institute Hungarian Academy of Sciences Lágymányosi utca 11 H–1111 Budapest, Hungary

e-mail: lszabo@csillag.ilab.sztaki.hu