

Zeitschrift:	Elemente der Mathematik
Herausgeber:	Schweizerische Mathematische Gesellschaft
Band:	62 (2007)
Artikel:	Note from the editorial board : note on rectangles with vertices on prescribed circles
Autor:	Blatter, Christian / Wanner, Gerhard
DOI:	https://doi.org/10.5169/seals-98919

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Note from the Editorial Board

Note on rectangles with vertices on prescribed circles

Christian Blatter and Gerhard Wanner

Inspired by [2], Ionascu and Stanica considered the following problem: The four subsequent vertices of a rectangle R in the plane are at distances r_1, r_2, r_3, r_4 from the origin. Given these data, what can be said about the area A of R ? In a recent paper [1] they proved

Theorem 1. (a) *For given $r_i \geq 0$, rectangles R of the described kind exist iff*

$$r_1^2 + r_3^2 = r_2^2 + r_4^2. \quad (1)$$

(b) *The areas of these rectangles lie between the bounds*

$$A_{\min} = |r_2 r_4 - r_1 r_3|, \quad A_{\max} = r_2 r_4 + r_1 r_3.$$

The authors call their problem “unusual because of its surprisingly simple answer in spite of our rather laborious solution” (which takes 10 pages). In this note we shall prove Theorem 1 in a simple way, making use of no more than the Pythagorean theorem and the formula for the derivative of a product.

After cyclic reordering, and neglecting special or degenerate cases, we may assume that $0 < r_1 < r_i$ ($2 \leq i \leq 4$). We may also assume that the sides of the rectangle are parallel to the axes and that the point $P_1 = (x, y)$ on the circle of radius r_1 is the lower lefthand vertex of R (Fig. 1). From P_1 we draw a horizontal to the right and obtain the vertex $P_2 = (\bar{x}, y)$ on the circle of radius r_2 ; in a similar way, going from P_1 vertically upwards, we obtain the vertex $P_4 = (x, \bar{y})$ on the circle of radius r_4 . So far we have

$$\begin{aligned} x^2 + y^2 &= r_1^2, \\ \bar{x}^2 + y^2 &= r_2^2, \quad \bar{x} > 0, \\ x^2 + \bar{y}^2 &= r_4^2, \quad \bar{y} > 0. \end{aligned} \quad (2)$$

The upper righthand vertex of R is $P_3 = (\bar{x}, \bar{y})$, and it lies on the circle of radius r_3 iff

$$r_3^2 = \bar{x}^2 + \bar{y}^2 = r_2^2 - y^2 + r_4^2 - x^2 = r_2^2 + r_4^2 - r_1^2,$$




Figure 1

i.e., iff (1) holds. Since the above construction is always possible, part (a) of the theorem follows.

Remark. The midpoint of R bisects both diagonals. It follows that (1) can be seen as an instance of the so-called Pappus-Fagnano-Legendre formula $a^2 + b^2 = (c^2 + z^2)/2$ where $z/2$ is the length of the median through C of a triangle with sides a, b, c .

For our extremal problem we have to consider admissible variations $dx, dy, d\bar{x}, d\bar{y}$ of the quantities x, y, \bar{x}, \bar{y} . To this end we differentiate the equations (2) and obtain

$$x dx + y dy = 0, \quad \bar{x} d\bar{x} + y dy = 0, \quad x dx + \bar{y} d\bar{y} = 0 \quad (3)$$

from which we easily infer

$$d\bar{x} = \frac{x}{\bar{x}} dx, \quad d\bar{y} = \frac{y}{\bar{y}} dy. \quad (4)$$

The area of R is $A = (\bar{x} - x)(\bar{y} - y)$; it is maximal or minimal iff

$$dA = (d\bar{x} - dx)(\bar{y} - y) + (\bar{x} - x)(d\bar{y} - dy) = 0.$$

Using (4) we get

$$dA = \frac{x - \bar{x}}{\bar{x}}(\bar{y} - y)dx + (\bar{x} - x)\frac{y - \bar{y}}{\bar{y}}dy = -\frac{A}{\bar{x}\bar{y}}(\bar{y}dx + \bar{x}dy).$$

So the condition $dA = 0$ reduces to $\bar{y}dx + \bar{x}dy = 0$. Combined with the first equation (3) this shows that in the stationary situation we necessarily have

$$\frac{y}{x} = \frac{\bar{x}}{\bar{y}}. \quad (5)$$

In order to determine the maximal and minimal areas of the rectangle we argue as follows: From (5) and (2) we get

$$\frac{\bar{x}^2}{\bar{y}^2} = \frac{y^2}{x^2} = \frac{r_2^2 - \bar{x}^2}{r_4^2 - \bar{y}^2}, \quad \text{whence} \quad \bar{x}^2(r_4^2 - \bar{y}^2) = \bar{y}^2(r_2^2 - \bar{x}^2),$$

and after cancelling terms we see that

$$\frac{\bar{y}}{\bar{x}} = \frac{r_4}{r_2}, \quad \frac{y}{x} = \frac{r_2}{r_4},$$

the latter using (5) again.

We now know the ratios of these coordinates as well as their Pythagorean sums, whence they must be

$$x = \pm \frac{r_4}{\sqrt{r_2^2 + r_4^2}} r_1, \quad y = \pm \frac{r_2}{\sqrt{r_2^2 + r_4^2}} r_1, \quad \bar{x} = \frac{r_2}{\sqrt{r_2^2 + r_4^2}} r_3, \quad \bar{y} = \frac{r_4}{\sqrt{r_2^2 + r_4^2}} r_3.$$

Inserting these values into the formula for A we obtain, using (1):

$$A = \frac{(r_2 r_3 \mp r_4 r_1)(r_4 r_3 \mp r_2 r_1)}{r_2^2 + r_4^2} = \frac{r_1^2 + r_3^2}{r_2^2 + r_4^2} r_2 r_4 \mp r_1 r_3 = r_2 r_4 \mp r_1 r_3.$$

This leads to

$$A_{\min} = r_2 r_4 - r_1 r_3, \quad A_{\max} = r_2 r_4 + r_1 r_3,$$

as stated. It is easily seen that for the maximum the origin lies in the interior of R whereas for the minimum it is in the exterior.

References

- [1] Ionascu, E.J.; Stanica, P.: Extremal values for the area of rectangles with vertices on concentrical circles. *Elem. Math.* 62 (2007), 30–39.
- [2] Zahlreich Problems Group: Problem 11057. *Amer. Math. Monthly* 111 (2004), 64. Solution: *ibid.* 113 (2006), 82.

Christian Blatter
Albertus-Walder-Weg 16
CH-8606 Greifensee, Switzerland
e-mail: christian.blatter@math.ethz.ch

Gerhard Wanner
Université/Mathématiques
Case postale 64
CH-1211 Génève 4, Switzerland
e-mail: Gerhard.Wanner@math.unige.ch