

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 63 (2008)

Artikel: On summing to arbitrary real numbers
Autor: Hanl, Jaroslav / Šustek, Jan / Nair, Radhakrishnan
DOI: <https://doi.org/10.5169/seals-99061>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On summing to arbitrary real numbers

Jaroslav Hančl, Jan Šustek, Radhakrishnan Nair,
Pavel Rucki and Dmitry Bodyagin

Jaroslav Hančl completed his studies in mathematics at Charles University in Prague in 1984. Presently he holds a position at the University of Ostrava. His main field of research is diophantine approximation.

Jan Šustek is a Ph.D. student at the Department of Mathematics of the University of Ostrava working in diophantine approximation.

Radhakrishnan Nair received his Ph.D. at the University of Warwick in 1986. Presently he holds a position at the University of Liverpool. His main field of research are questions on distributions in number theory.

Pavel Rucki completed his studies in mathematics at the University of Ostrava in 2006. Presently he is a postdoctoral fellow at the same university. His main field of research is diophantine approximation.

Dmitry Bodyagin is a postgraduate student at the Institute of Mathematics of the Belarusian Academy of Sciences. His main field of research is diophantine approximation.

Following Erdős [1] we say a sequence $\{a_n\}_{n=1}^{\infty}$ is *irrational* if the set $\{\sum_{n \geq 1} \frac{1}{a_n c_n} \mid c_n \in \mathbb{N}\}$, which we refer to henceforth as its *expressible set*, contains no rational numbers. In [1] it is shown that if $\lim_{n \rightarrow \infty} a_n^{1/2^n} = \infty$ and $a_n \in \mathbb{N}$ for all $n \in \mathbb{N}$ then $\sum_{n \geq 1} a_n^{-1}$ is an irrational number. From this Erdős deduces that the sequence $\{2^{2^n}\}_{n=1}^{\infty}$ is an irrational sequence. Thus its expressible set contains no rational numbers. In [2] it is shown that if $a_n \in \mathbb{R}^+$ for all $n \in \mathbb{N}$ and $\limsup_{n \rightarrow \infty} \frac{1}{n} \log_2 \log_2 a_n < 1$ then the expressible set of the sequence $\{a_n\}_{n=1}^{\infty}$ contains an interval. It seems to be the case that in general finding the expressible set for the sequence $\{a_n\}_{n=1}^{\infty}$ is not easy.

Ein interessantes zahlentheoretisches Problem ist die Frage nach der Rationalität des Werts einer konvergenten Reihe reeller Zahlen. An diese Fragestellung anknüpfend nennen wir mit P. Erdős eine Folge $\{a_n\}_{n=1}^{\infty}$ reeller Zahlen irrational, falls die Menge $E = \{\sum_{n=1}^{\infty} 1/(a_n c_n) \mid c_n \in \mathbb{N}\}$ keine rationale Zahl enthält. In der vorliegenden Arbeit beweisen die Autoren für den Fall, dass die Reihe $\sum_{n=1}^{\infty} 1/a_n$ bedingt konvergent ist, dass die Menge E jeweils die gesamte reelle Zahlengerade ausschöpft.

In this paper we give conditions on $\{a_n\}_{n=1}^{\infty}$ to ensure that its expressible set is equal to \mathbb{R} . We prove the following:

Theorem 1. *Let $\{a_n\}_{n=1}^{\infty}$ be a sequence of nonzero real numbers such that the series $\sum_{n=1}^{\infty} \frac{1}{a_n}$ is conditionally convergent. Then its expressible set is equal to \mathbb{R} .*

A series is conditionally convergent if it is convergent but the series of the absolute values of its terms is not. Theorem 1 is an immediate consequence of the following more general theorem.

Theorem 2. *Let $\{a_n\}_{n=1}^{\infty}$ be a sequence of nonzero real numbers such that the series $\sum_{n=1}^{\infty} \frac{1}{a_n}$ is conditionally convergent. Then for every pair α, β of real numbers with $\alpha \leq \beta$ there exists a sequence $\{c_n\}_{n=1}^{\infty}$ of positive integers such that*

$$\alpha = \liminf_{N \rightarrow \infty} \sum_{n=1}^N \frac{1}{a_n c_n} \quad \text{and} \quad \beta = \limsup_{N \rightarrow \infty} \sum_{n=1}^N \frac{1}{a_n c_n}. \quad (1)$$

For the proof of Theorem 2 we need the following two lemmas.

Lemma 1. *Let $\{a_n\}_{n=1}^{\infty}$ be a sequence of nonzero real numbers such that the series $\sum_{n=1}^{\infty} \frac{1}{a_n}$ is conditionally convergent. Then for every real number $A \geq 0$ and every integer $N \geq 0$ there exist a number $K \in \mathbb{N}$ and numbers $c_{N+1}, \dots, c_{N+K} \in \mathbb{N}$ such that*

$$\sum_{n=N+1}^{N+K} \frac{1}{a_n c_n} \in \left(A, A + \frac{1}{a_{N+K}} \right].$$

Proof. Define $\mathcal{P} = \{n \mid a_n > 0\}$ and $\mathcal{N} = \{n \mid a_n < 0\}$. The series $\sum_{n=1}^{\infty} \frac{1}{a_n}$ is conditionally convergent, hence

$$\sum_{\substack{n=N+1 \\ n \in \mathcal{P}}}^{\infty} \frac{1}{a_n} = \infty.$$

This implies that there exists a positive integer K such that

$$\sum_{\substack{n=N+1 \\ n \in \mathcal{P}}}^{N+K-1} \frac{1}{a_n} \leq A \quad \text{and} \quad \sum_{\substack{n=N+1 \\ n \in \mathcal{P}}}^{N+K} \frac{1}{a_n} > A.$$

The fact that

$$0 < \sum_{\substack{n=N+1 \\ n \in \mathcal{P}}}^{N+K} \frac{1}{a_n} - \sum_{\substack{n=N+1 \\ n \in \mathcal{P}}}^{N+K-1} \frac{1}{a_n} = \frac{1}{a_{N+K}}$$

immediately gives

$$s = \sum_{\substack{n=N+1 \\ n \in \mathcal{P}}}^{N+K} \frac{1}{a_n} \in \left(A, A + \frac{1}{a_{N+K}} \right].$$

Now consider two cases:

(1) Assume that $\mathcal{N} \cap \{N+1, \dots, N+K\} = \emptyset$. In this case put $c_n = 1$ for every $n = N+1, \dots, N+K$ and the result follows.

(2) Now suppose that

$$r = \sum_{\substack{n=N+1 \\ n \in \mathcal{N}}}^{N+K} \frac{1}{a_n} < 0.$$

Put $C = \lceil \frac{r}{A-s} \rceil + 1$. Then

$$0 > \sum_{\substack{n=N+1 \\ n \in \mathcal{N}}}^{N+K} \frac{1}{Ca_n} = \frac{1}{C} \sum_{\substack{n=N+1 \\ n \in \mathcal{N}}}^{N+K} \frac{1}{a_n} > \frac{A-s}{r} \cdot r = A-s.$$

Hence the result follows by taking $c_n = 1$ for $n \in \{N+1, \dots, N+K\} \cap \mathcal{P}$ and $c_n = C$ for $n \in \{N+1, \dots, N+K\} \cap \mathcal{N}$. \square

Lemma 2. Let $\{a_n\}_{n=1}^{\infty}$ be a sequence of nonzero real numbers such that the series $\sum_{n=1}^{\infty} \frac{1}{a_n}$ is conditionally convergent. Then for every real number $A \leq 0$ and every integer $N \geq 0$ there exist a number $K \in \mathbb{N}$ and numbers $c_{N+1}, \dots, c_{N+K} \in \mathbb{N}$ such that

$$\sum_{n=N+1}^{N+K} \frac{1}{a_n c_n} \in \left[A - \left| \frac{1}{a_{N+K}} \right|, A \right).$$

Proof. Using the transformation $a_n \mapsto -a_n$ and Lemma 1 we obtain Lemma 2. \square

Proof of Theorem 2. In the following we set

$$S_k = \sum_{n=1}^k \frac{1}{a_n c_n}.$$

If $\beta \geq 0$ then putting $A = \beta$ and $N = 0$ into Lemma 1 we obtain a number K and a sequence $\{c_n\}_{n=1}^K$ such that

$$S_K \in \left(\beta, \beta + \frac{1}{a_K} \right].$$

Then set $N_0 = 0$ and $N_1 = K$.

Similarly, if $\beta < 0$ then $\alpha < 0$, and putting $A = \alpha$ and $N = 0$ into Lemma 2 we get K and $\{c_n\}_{n=1}^K$ with

$$S_K \in \left[\alpha - \left| \frac{1}{a_K} \right|, \alpha \right).$$

Then set $N_0 = K$.

Now we will construct the sequence $\{c_n\}_{n=1}^{\infty}$ by induction. Consider two cases:

(1) Suppose that we have constructed the sequence $\{N_m\}_{m=0}^{2t+1}$, $t \in \mathbb{N}_0$, with

$$S_{N_{2t+1}} \in \left(\beta, \beta + \frac{1}{a_{N_{2t+1}}} \right].$$

Lemma 2 implies that there exist K and $\{c_n\}_{n=N_{2t+1}+1}^{N_{2t+1}+K}$ such that

$$\sum_{n=N_{2t+1}+1}^{N_{2t+1}+K} \frac{1}{a_n c_n} \in \left[\alpha - S_{N_{2t+1}} - \left| \frac{1}{a_{N_{2t+1}+K}} \right|, \alpha - S_{N_{2t+1}} \right).$$

Let $N_{2t+2} = N_{2t+1} + K$. Then we have

$$S_{N_{2t+2}} \in \left[\alpha - \left| \frac{1}{a_{N_{2t+2}}} \right|, \alpha \right).$$

(2) Suppose that we have constructed the sequence $\{N_m\}_{m=0}^{2t}$, $t \in \mathbb{N}_0$, with

$$S_{N_{2t}} \in \left[\alpha - \left| \frac{1}{a_{N_{2t}}} \right|, \alpha \right).$$

Lemma 1 implies that there exist K and $\{c_n\}_{n=N_{2t}+1}^{N_{2t}+K}$ such that

$$\sum_{n=N_{2t}+1}^{N_{2t}+K} \frac{1}{a_n c_n} \in \left(\beta - S_{N_{2t}}, \beta - S_{N_{2t}} + \frac{1}{a_{N_{2t}+K}} \right).$$

Let $N_{2t+1} = N_{2t} + K$. Then we have

$$S_{N_{2t+1}} \in \left(\beta, \beta + \frac{1}{a_{N_{2t+1}}} \right].$$

Using alternatively cases (1) and (2) we construct the whole sequence $\{c_n\}_{n=1}^{\infty}$. From the construction it follows that

- $\alpha - \left| \frac{1}{a_k} \right| \leq S_k \leq \beta + \left| \frac{1}{a_k} \right|$ for every $k \geq N_1$,
- $S_{N_{2t}} < \alpha$ for every $t \in \mathbb{N}$,
- $S_{N_{2t+1}} > \beta$ for every $t \in \mathbb{N}_0$.

The series $\sum_{n=1}^{\infty} \frac{1}{a_n}$ is conditionally convergent, hence $\frac{1}{a_n} \rightarrow 0$. This implies that

$$\lim_{t \rightarrow \infty} S_{N_{2t}} = \alpha \quad \text{and} \quad \lim_{t \rightarrow \infty} S_{N_{2t+1}} = \beta$$

and the result follows. □

References

- [1] Erdős, P.: Some problems and results on the irrationality of the sum of infinite series. *J. Math. Sci.* 10 (1975), 1–7.
- [2] Hančl, J.: Expression of Real Numbers With the Help of Infinite Series. *Acta Arith.* LIX.2, (1991), 97–104.

Jaroslav Hančl, Jan Šustek
Department of Mathematics and Institute for
Research and Applications of Fuzzy Modeling
University of Ostrava
30. dubna 22
701 03 Ostrava 1, Czech Republic
e-mail: hancl@osu.cz, jan.sustek@seznam.cz

Radhakrishnan Nair
Mathematical Sciences
Peach Street
Liverpool L69 7ZL, U.K.
e-mail: nair@liverpool.ac.uk

Pavel Rucki
Department of Mathematics and Didactics
University of Ostrava
Dvořákova 7
701 03 Ostrava 1, Czech Republic
e-mail: pavel.rucki@seznam.cz

Dmitry Bodyagin
Department of Theory of Numbers
Institute of Mathematics
National Academy of Sciences of Belarus
Surganov str. 11
220072 Minsk, Belarus
e-mail: bodiagin@mail.ru

The paper was supported by the grants no. 201/04/0381, 201/07/0191,
and MSM6198898701