

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 65 (2010)

**Artikel:** A trigonometric double-inequality  
**Autor:** Alzer, Horst  
**DOI:** <https://doi.org/10.5169/seals-130688>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## A trigonometric double-inequality

Horst Alzer

Horst Alzer received his doctoral degree in mathematics from the University of Bonn, and his habilitation in mathematics from the University of Würzburg.

Trigonometric functions, in particular the sine and cosine functions, play a relevant rôle in physics and various branches of mathematics. In view of their importance the properties of these functions have been studied intensively. Of special interest are inequalities for trigonometric functions. In his classical book *Analytic Inequalities* D.S. Mitrinović dedicated a whole section to this subject. In the recent past, several authors discovered remarkable new inequalities for sine, cosine, and their relatives. Many of them appeared in the *Journal of Inequalities in Pure and Applied Mathematics*, in *Mathematical Inequalities and Applications*, and in the *Journal of Inequalities and Applications*.

In this note we present a double-inequality involving sine and cosine. Our work has been inspired by a short paper published by R.J. Webster [4] in 1984. He proved the elegant inequalities

$$|\sin(\cos x)| \leq |\cos x| \leq \cos(\sin x) \quad (x \in \mathbb{R}). \quad (1)$$

See also [1] and [2]. We offer the following counterpart of (1):

Seit Jahrzehnten spielen Ungleichungen für trigonometrische Funktionen eine wichtige Rolle in der Mathematik. D.S. Mitrinović hat dem Thema in seinem Buch „Analytic Inequalities“ ein ganzes Kapitel gewidmet. In diesen Zusammenhang gehört auch die elegante Ungleichung

$$|\sin(\cos x)| \leq |\cos x| \leq \cos(\sin x) \quad (x \in \mathbb{R}),$$

die im Jahr 1984 von R.J. Webster veröffentlicht wurde. In der vorliegenden Arbeit stellt der Autor für beliebige reelle Werte von  $x$  eine Variation der vorhergehenden Ungleichungen in der Form

$$\alpha |\sin(\cos x) + \sin(\sin x)| \leq |\cos x + \sin x| \leq \beta (\cos(\cos x) + \cos(\sin x))$$

vor mit den bestmöglichen Konstanten  $\alpha$ ,  $\beta$ , welche durch  $\alpha = (\sqrt{2} \sin(1/\sqrt{2}))^{-1}$  bzw.  $\beta = (\sqrt{2} \cos(1/\sqrt{2}))^{-1}$  gegeben sind.

**Theorem.** For all real numbers  $x$  we have

$$\alpha |\sin(\cos x) + \sin(\sin x)| \leq |\cos x + \sin x| \leq \beta (\cos(\cos x) + \cos(\sin x)) \quad (2)$$

with the best possible constant factors

$$\alpha = \frac{1}{\sqrt{2} \sin(1/\sqrt{2})} = 1.0884\dots \quad \text{and} \quad \beta = \frac{1}{\sqrt{2} \cos(1/\sqrt{2})} = 0.9301\dots \quad (3)$$

*Proof.* First, we consider the left-hand side of (2) with  $\alpha$  as given in (3). We establish

$$f(x) = |\cos x + \sin x| - \alpha |\sin(\cos x) + \sin(\sin x)| \geq 0 \quad \text{for all } x \in \mathbb{R}.$$

Since  $f(x + \pi) = f(x)$ , it suffices to show that  $f$  is nonnegative on  $[0, \pi]$ . Using the formulas

$$\cos a + \sin a = \sqrt{2} \sin\left(\frac{\pi}{4} + a\right), \quad \cos a - \sin a = \sqrt{2} \cos\left(\frac{\pi}{4} + a\right), \quad (4)$$

and

$$\sin a + \sin b = 2 \sin \frac{a+b}{2} \cos \frac{a-b}{2}$$

we obtain the representation

$$\frac{1}{2\alpha} f(x) = |\delta| \sin \frac{1}{\sqrt{2}} - \left| \sin \frac{\delta}{\sqrt{2}} \cos \sqrt{\frac{1-\delta^2}{2}} \right|$$

with

$$\delta = \delta(x) = \sin\left(\frac{\pi}{4} + x\right). \quad (5)$$

If  $0 \leq x \leq 3\pi/4$ , then  $0 \leq \delta \leq 1$ . This leads to

$$\frac{1}{2\alpha} f(x) = \delta \sin \frac{1}{\sqrt{2}} - \sin \frac{\delta}{\sqrt{2}} \cos \sqrt{\frac{1-\delta^2}{2}}.$$

Furthermore, if  $3\pi/4 \leq x \leq \pi$ , then  $0 \leq -\delta \leq 1/\sqrt{2}$ . Hence,

$$\frac{1}{2\alpha} f(x) = (-\delta) \sin \frac{1}{\sqrt{2}} - \sin \frac{-\delta}{\sqrt{2}} \cos \sqrt{\frac{1-(-\delta)^2}{2}}.$$

This implies that we have to prove that

$$0 \leq t \sin \frac{1}{\sqrt{2}} - \sin \frac{t}{\sqrt{2}} \cos \sqrt{\frac{1-t^2}{2}} \quad \text{for } t \in [0, 1].$$

We denote the expression on the right-hand side by  $g(t)$ . Differentiation gives

$$\sqrt{2}(1-t^2)^{3/2} g''(t) = \sqrt{2}t(t^2-1) \cos \frac{t}{\sqrt{2}} \sin \sqrt{\frac{1-t^2}{2}} + h\left(\sqrt{\frac{1-t^2}{2}}\right) \sin \frac{t}{\sqrt{2}}, \quad (6)$$

where

$$h(s) = s \cos s - \sin s.$$

Since  $h(s) \leq 0$  for  $s \in [0, \pi]$ , we conclude from (6) that  $g''(t) \leq 0$  for  $t \in [0, 1]$ . Thus,  $g$  is concave. We have  $g(0) = g(1) = 0$ . Hence,  $g(t) \geq 0$  for  $t \in [0, 1]$ .

Now, we prove the second inequality in (2) with  $\beta$  as given in (3). We show that

$$u(x) = \beta(\cos(\sin x) + \cos(\cos x)) - |\cos x + \sin x| \geq 0 \quad \text{for all } x \in \mathbb{R}.$$

Since  $u$  is periodic with period  $\pi$ , we may assume that  $x \in [0, \pi]$ . Applying (4) and

$$\cos a + \cos b = 2 \cos \frac{a+b}{2} \cos \frac{a-b}{2}$$

yields

$$\frac{1}{2\beta} u(x) = \cos \frac{\delta}{\sqrt{2}} \cos \sqrt{\frac{1-\delta^2}{2}} - |\delta| \cos \frac{1}{\sqrt{2}},$$

where  $\delta$  is defined in (5). If  $0 \leq x \leq 3\pi/4$ , then  $0 \leq \delta \leq 1$  and

$$\frac{1}{2\beta} u(x) = \cos \frac{\delta}{\sqrt{2}} \cos \sqrt{\frac{1-\delta^2}{2}} - \delta \cos \frac{1}{\sqrt{2}}.$$

Moreover, if  $3\pi/4 \leq x \leq \pi$ , then  $0 \leq -\delta \leq 1/\sqrt{2}$  and

$$\frac{1}{2\beta} u(x) = \cos \frac{-\delta}{\sqrt{2}} \cos \sqrt{\frac{1-(-\delta)^2}{2}} - (-\delta) \cos \frac{1}{\sqrt{2}}.$$

Therefore, we have to show that

$$0 \leq \cos \frac{t}{\sqrt{2}} \cos \sqrt{\frac{1-t^2}{2}} - t \cos \frac{1}{\sqrt{2}} \quad \text{for } t \in [0, 1]. \quad (7)$$

We define for  $t \in (0, 1]$ :

$$v(t) = \cos \frac{t}{\sqrt{2}} \quad \text{and} \quad w(t) = \frac{1}{t} \cos \sqrt{\frac{1-t^2}{2}}.$$

The function  $v$  is decreasing. To prove that  $w$  is also decreasing, we apply the inequality

$$\tan y \leq y + \frac{\tan \lambda - \lambda}{\lambda^3} y^3,$$

which holds for all  $\lambda$  and  $y$  with  $0 < \lambda < \pi/2$  and  $0 \leq y \leq \lambda$ . A proof is given in [3, pp. 245, 246]. We set  $\lambda = 1/\sqrt{2}$ . Since

$$\frac{\tan \lambda - \lambda}{\lambda^3} = 0.41\dots,$$

we get

$$\tan y \leq y + \frac{1}{2}y^3 \quad \text{for } y \in [0, 1/\sqrt{2}]. \quad (8)$$

Let

$$y(t) = \sqrt{\frac{1-t^2}{2}}.$$

Using (8) gives

$$\begin{aligned} \frac{2y(t)}{\cos y(t)} w'(t) &= \tan y(t) - \frac{2y(t)}{t^2} \leq y(t) + \frac{1}{2}y(t)^3 - \frac{2y(t)}{t^2} \\ &= -\frac{2y(t)^5 + 3y(t)^3 + 2y(t)}{2t^2} < 0. \end{aligned}$$

Thus,  $v$  and  $w$  are positive and decreasing on  $(0, 1]$ , so that we obtain

$$v(t)w(t) \geq v(1)w(1) = \cos \frac{1}{\sqrt{2}} \quad \text{for } t \in (0, 1].$$

This settles (7). The proof of double-inequality (2) is complete.

We define

$$P(x) = \left| \frac{\cos x + \sin x}{\sin(\cos x) + \sin(\sin x)} \right| \quad \text{and} \quad Q(x) = \frac{|\cos x + \sin x|}{\cos(\cos x) + \cos(\sin x)}.$$

Then we have

$$P(\pi/4) = \frac{1}{\sqrt{2} \sin(1/\sqrt{2})} \quad \text{and} \quad Q(\pi/4) = \frac{1}{\sqrt{2} \cos(1/\sqrt{2})}.$$

This implies that the constant factors given in (3) are the best possible.  $\square$

## References

- [1] Bensom, H.: On  $\sin \circ \cos$  and  $\cos \circ \sin$ . *Math. Gaz.* 67 (1983), 43–44.
- [2] Haigh, G.: Another proof that  $\cos(\sin x) > \sin(\cos x)$ . *Math. Gaz.* 68 (1984), 36.
- [3] Mitrinović, D.S.: *Analytic Inequalities*. Springer, New York 1970.
- [4] Webster, R.J.:  $\cos(\sin x) \geq |\cos x| \geq |\sin(\cos x)|$ . *Math. Gaz.* 68 (1984), 37.

Horst Alzer  
Morsbacher Str. 10  
D–51545 Waldbröl, Germany  
e-mail: H.Alzer@gmx.de