Zeitschrift: Elemente der Mathematik

Herausgeber: Schweizerische Mathematische Gesellschaft

Band: 65 (2010)

Artikel: On a remark of Makowski about perfect numbers

Autor: Gallardo Luis H.

DOI: https://doi.org/10.5169/seals-130696

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 27.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

On a remark of Makowski about perfect numbers

Luis H. Gallardo

Luis H. Gallardo received his Ph.D. from the University of Paris 7 in 1978. He is an associate professor at the University of Brest, Brittany, France. He works on arithmetic problems related to polynomials.

1 Introduction

A positive number n > 0 is called perfect if it is equal to the sum $\sigma(n) - n$ of its proper divisors. Some classical papers on perfect numbers were written by Peirce [3], Servais [5], and Sylvester [7] in the 19th century. More recently, Weisstein [9] lists many other properties of the perfect numbers including Makowski's result [2] that 28 is the only even perfect number that exceeds a cube by one. It is natural to wonder whether the result holds for all perfect numbers. More generally, we can ask whether 28 is the only perfect number n that is a sum of two non-negative cubes, say, $n = x^3 + a^3$ with $a \ge 0$ and x > 0. This seems to be a very difficult question.

The object of this paper is

- (a) to prove that 28 is the only even perfect number that is a sum of two positive integral cubes;
- (b) to prove (case a = 0) that there are no perfect numbers that are cubes;
- (c) to describe a one parameter family of sums s of two cubes such that s is odd and $\sigma(s) \equiv 2 \pmod{4}$.

Eine positive natürliche Zahl n heisst bekanntlich vollkommen, falls n gleich der Summe seiner echten positiven Teiler ist; die Zahlen 6 und 28 sind beispielsweise vollkommen. Sätze von Euklid und Euler besagen, dass eine gerade Zahl n genau dann vollkommen ist, wenn $n = 2^{p-1}(2^p - 1)$ gilt, wobei $2^p - 1$ und somit auch p Primzahlen sind. Ausgehend von der Gleichung $28 = 3^3 + 1$ bewies A. Makowski im Jahr 1961, dass die Zahl 28 die einzige gerade vollkommene Zahl ist, die eine Kubikzahl um eins übertrifft. In der vorliegenden Arbeit verallgemeinert der Autor dieses Ergebnis dahingehend, dass er nachweist, dass 28 die einzige gerade vollkommene Zahl ist, die Summe zweier Kubikzahlen ist.

122 L.H. Gallardo

Now we recall some basic results about perfect numbers. It is well known that any even perfect number n has only two prime factors, so that

$$n = t(2t - 1) \tag{1}$$

where $t = 2^{p-1}$ for some prime number p such that 2t - 1 is also prime. Observe that 2t - 1 > t and that gcd(t, 2t - 1) = 1. On the other hand, the most basic result about the form of a possible odd perfect number comes from Euler. Euler [1] proved that odd perfect numbers n have the form

$$n = v^{4k+1}z^2 \tag{2}$$

with $k \ge 0$ a non-negative integer, z > 0 a positive integer and y a prime number such that $\gcd(y,z) = 1$ and $y \equiv 1 \pmod 4$. This is indeed an easy consequence of the fact that $\sigma(n) \equiv 2 \pmod 4$; where $\sigma(n)$ denotes the sum of all positive divisors of n.

Let *n* be an odd perfect number. Touchard [8] proved that either

$$n \equiv 1 \pmod{12}$$
 or $n \equiv 9 \pmod{36}$. (3)

2 The only even perfect number that is also a sum of two cubes is 28

Assume that the even perfect number n is a sum of two cubes:

$$n = x^3 + a^3 = (x + a)(x^2 - ax + a^2)$$

for some integers x>0 and $a\geq 0$. In particular a and x have the same parity. Let δ be the discriminant of $x^2-(a+1)x+a^2-a$. One has $\delta=-(3a^2-6a-1)$. Observe that $x^2-ax+a^2>x+a$ for all a>2 since $\delta<0$ if and only if a>2. Let us assume now that a>2. The case $a\in\{0,1,2\}$ will be considered later. Thus, from (1) we see that $x+a=2^{p-1}$ and $x^2-ax+a^2=2^p-1$ so that

$$x^{2} - ax + a^{2} = 2(x+a) - 1. (4)$$

Since (4) has integral roots x and a+2-x the discriminant $\Delta=-3a(a-4)$ of the quadratic

$$x^2 - (a+2)x + (a-1)^2$$

must be a perfect square. It is easy to see that Δ is non-negative exactly when $a \in \{0, 1, 2, 3, 4\}$. So it remains to consider the cases $a \in \{3, 4\}$. If a = 3 then $x \in \{4, 1\}$. But n is even so x = 1. Thus, x = 1 and a = 3. In other words, we get the perfect number n = 28. If a = 4 then x = 3. This is not possible since a and x have the same parity.

Now we discuss the case $a \in \{1, 2\}$. Observe that if $x + a \ge x^2 - ax + a^2$ then we must have $x + a > x^2 - ax + a^2$ since the perfect number $n = (x + a)(x^2 - ax + a^2)$ is never a square. Take a = 2. Thus $x + 2 > x^2 - 2x + 4$. But this is not possible for an integer x. It remains only the case a = 1. In this case we have as before $x + 1 > x^2 - 2x + 1$. This is true only for x = 1. So a = 1 and x = 1. But $n = x^3 + a^3 = 2$ is not perfect. Finally, observe that a = 0 is not possible since 3 does not divide the exponent 1 of the prime 2t - 1 in n = t(2t - 1). Thus, n = t(2t - 1) is not a cube. This proves the result.

3 A perfect number cannot be a cube

We have just seen that an even perfect number cannot be a cube. We assume in the rest of the section that $n = x^3$ is an odd perfect number. We will get a contradiction by considering the equality

$$\sigma(n) = 2n \pmod{12}$$
.

It follows from (2) that

$$n = p^{12k+9}r^6$$

for some non-negative integer $k \geq 0$, for some prime number $p \equiv 1 \pmod 4$ and for some positive integer r > 0 such that $\gcd(p,r) = 1$. Assume that r can be factored as $r = q_1^{\alpha_1} \cdots q_m^{\alpha_m}$ with prime numbers q_1, \ldots, q_m . We have then

$$2n = \sigma(n) = \sigma(p^{12k+9})\sigma(q_1^{6\alpha_1})\cdots\sigma(q_m^{6\alpha_m}). \tag{5}$$

3.1 Case 1: gcd(3, n) = 1

From (3) we get $2n \equiv 2 \pmod{12}$. Now we compute the value of the right hand side of (5) modulo 12. First of all (3) implies $p \in \{1, 5\} \pmod{12}$ for the Euler prime p. So

$$\sigma(p^{12k+9}) = 1 + p + \dots + p^{12k+9} \equiv 12k + 10 \equiv -2 \pmod{12}$$

when $p \equiv 1 \pmod{12}$ and

$$\sigma(p^{12k+9}) = (1^0 + 5^2 + \dots + 5^{12k+10}) + 5(1^0 + 5^2 + \dots + 5^{12k+8})$$

$$\equiv (6k+6) + 5(6k+5) = 36k+31 \equiv 7 \pmod{12}$$

when $p \equiv 5 \pmod{12}$, since trivially $5^2 \equiv 1 \pmod{12}$.

Observe that $q_i > 3$ is an odd prime. So $q_i \in \{1, 5, 7, 11\} \pmod{12}$. Thus, $q_i^2 \equiv 1 \pmod{12}$. In other words $q_i^{2k+1} \equiv q_i \pmod{12}$ and $q_i^{2k} \equiv 1 \pmod{12}$ for any integer k. Thus, one has

$$\sigma(q_i^{6\alpha_i}) \equiv 6\alpha_i + 1 \pmod{12}$$

when $q_i \equiv 1 \pmod{12}$, and

$$\sigma(q_i^{6\alpha_i}) = 1^0 + q_i^2 + \ldots + q_i^{6q_i} + (q_i^1 + q_i^3 + \ldots + q_i^{6q_i - 1})$$

$$\equiv (3\alpha_i + 1) + (3\alpha_i)q_i \equiv 3\alpha_i(1 + q_i) + 1 \pmod{12}$$

when $q_i \not\equiv 1 \pmod{12}$.

Set $a_i = 3\alpha_i(1+q_i) + 1$. Observe that $a_i \equiv 1 \pmod{12}$ when $q_i \equiv 7 \pmod{12}$ or when $q_i \equiv 11 \pmod{12}$. We have also $a_i \equiv 6\alpha_i + 1 \pmod{12}$ when $q_i \equiv 5 \pmod{12}$.

Moreover, observe that for any integer x, either $6x + 1 \equiv 1 \pmod{12}$ or $6x + 1 \equiv 7 \pmod{12}$. Thus, for all i one has

$$\sigma(q_i^{6\alpha_i}) \in \{1, 7\} \pmod{12}.$$
 (6)

124 L.H. Gallardo

Observe that $z \in \{-2, 7\} \pmod{12}$ and $y \in \{1, 7\} \pmod{12}$ implies $zy \in \{-2, 1, 7\} \pmod{12}$. Thus,

$$2n = \sigma(n) = \sigma(p^{12k+9}) \prod_{i=1}^{m} \sigma(q_i^{6\alpha_i}) \in \{-2, 1, 7\} \pmod{12}.$$

This contradicts (3) that gives $2n \equiv 2 \pmod{12}$; thereby proving the result.

3.2 Case 2: gcd(3, n) = 0

Observe that

$$\sigma(3^{6\alpha_i}) = 1 + (3+3^2) + 3^2(3+3^2) + \dots \equiv 1 \pmod{12}$$

Hence, as before, we get

$$\sigma(n) \in \{-2, 1, 7\} \pmod{12}$$
.

But (3) implies that $2n \equiv 6 \pmod{12}$. So we get the contradiction $\sigma(n) \neq 2n$. This completes the proof that an odd perfect number cannot be a cube.

4 Our main result about a family of perfect candidates that are sums of two cubes

First of all a technical and useful lemma follows:

Lemma 4.1. If for some prime number p, for some non-negative integer $k \ge 0$ and for some positive integer $s \ge 1$ one has

$$p^{4k+1} = 2 + 3^{2s-1} (7)$$

then

$$k = 0$$
.

Proof. We choose r = p, A = 2, $B = 3^{2s-1}$, and x = 4k + 1 in order to have

$$A+B=r^{\chi}$$

We get PQ = 6 since PQ is the largest squarefree divisor of AB. Thus, in order to have the condition $\delta = (AB/P)^{1/2}$ is integral fulfilled, we are forced to take P = 6 and Q = 1. Hence, $\delta = 3^{s-1}$.

It then follows from [4, Theorem 3, p. 219] that

$$x < \frac{1}{2}QP^{1/2}\ln(P) = \sqrt{6}\,\frac{\ln(6)}{2} < 3.$$

But x is odd. So, x = 1 and k = 0. Another proof is to choose x = 2r - 1, y = 1, n = p, z = 4k + 1 in [4, Lemma 2, p. 228] so that we have

$$3^x + 2^y = n^z.$$

We get as before k = 0.

Our main result (whose proof is an immediate corollary of Lemma 4.1 and a check of the formulae) is then:

Theorem 4.2. Let r > 0 be a positive integer. Define $w(r) = 3^{4r-2}$. Let $k \ge 0$ be a non-negative integer such that the diophantine equation

$$p(r)^{4k+1} = 2 + 3^{4r-1},$$

has an integral solution p(r). Let also, $a(r) = 3w(r)^2 - 1$, $m(r) = 3w(r)^2 + 3w(r) + 1$, $n(r) = p(r)^{4k+1} \cdot 3^{4r} \cdot m(r)^2$, $x(r) = 3^{4r-1}p(r)^{4k+1} - a(r)$. Then,

$$k = 0$$

provided p(r) is a prime number. In all cases we have $w(r) \equiv 1 \pmod{4}$, so that $p(r) \equiv 1 \pmod{4}$, and

- a) $p(r)^{4k+1} = 3w(r) + 2$,
- b) x(r) = 2m(r) 1, $x(r) = 6w(r)^2 + 6w(r) + 1$,
- c) x(r) + a(r) = 3w(r)(3w(r) + 2),
- d) $x(r)^2 a(r)x(r) + a(r)^2 = 3(3w(r)^2 + 3w(r) + 1)^2$,
- e) $x(r)^3 + a(r)^3 = n(r)$.

In other words there is a one parameter family $\{n(r)\}$ of integers such that $n(r) \equiv 1 \pmod{4}$. Moreover if p(r) is prime then n(r) satisfies the necessary condition (2) to be a perfect number. Furthermore, n(r) is a sum of two cubes for each r.

It is then of some interest to obtain the r's for which p(r) is a prime number. We have

$$n(r) \equiv 1 \pmod{4}$$
 and $\sigma(n(r)) \equiv 2 \pmod{4}$.

A quick computer check gives

- a) if $r \equiv 0 \pmod{3}$ then $p(r) \equiv 0 \pmod{7}$;
- b) if $r \equiv 2 \pmod{5}$ then $p(r) \equiv 0 \pmod{11}$; etc.

Using a) we constructed a list L of r's for which $p(r) = 2 + 3^{4r-1}$ is a prime number. For the moment

$$L = \{1, 4, 16, 31, 35, 59, 61, 79, 91, 98, 283, 376, 1801, 10948, 11384, 26536\}$$
 (8)

contains sixteen elements. We have examined all possible r's up to 38338. This took some time, e.g., about 2 hours CPU to test each possible candidate r when r is close to, say, 30000. This was done on a 8 processor linux machine running command line cmaple 11. We do not know if $a_r = p(r)$ is a prime number for an infinity of r's. Compare with sequence A134753 in Sloane's database [6].

Observe that n(r) is perfect if and only if

$$2n(r) = 2 \cdot p(r) \cdot 3^{4r} \cdot m(r)^2 = (p(r) + 1) \cdot \sigma(3^{4r}) \cdot \sigma(m(r)^2) = \sigma(n(r)). \tag{9}$$

126 L.H. Gallardo

We deduce from (9) that if n(r) is perfect then the prime p(r) divides the product $\sigma(3^{4r}) \cdot \sigma(m(r)^2)$.

Indeed, we checked (in about only 15 seconds) that for all $r \in L$ the corresponding n(r) is not perfect since the prime p(r) does not divide $\sigma(m(r)^2)$.

In order to show that this suffices we claim that if n(r) is perfect then p(r) does not divide $\rho = \sigma(3^{4r})$. To prove the claim assume, to the contrary, that p(r) does divide ρ . Then from the definition of w(r) in Theorem 4.2 we obtain that

$$9w(r) = 3^{4r}$$
.

But from Theorem 4.2 part a) we get

$$27w(r) \equiv -18 \pmod{p(r)}$$
.

So

$$\rho = \frac{3(3^{4r}) - 1}{2} \equiv \frac{27w(r) - 1}{2} \equiv \frac{-19}{2} \pmod{p(r)}.$$

Thus, p(r) = 19. But 19, contrary to p(r), is not congruent to 1 (mod 4). This proves the claim.

References

- [1] Euler, L.: *De numeris amicabilibus*. Reprinted in: Opera posthuma, Euler archive [E798]. http://www.math.dartmouth.edu/~euler1 (1862), 85–100.
- [2] Makowski, A.: Remark on perfect numbers. Elem. Math. 17 (1962), 109.
- [3] Peirce, B.: On perfect numbers. New York Math. Diary 2, XIII (1832), 267–277.
- [4] Scott, R.; Styer, R.: On $p^x q^y = c$ and related three term exponential Diophantine equations with prime bases. J. Number Theory 105 (2004), 212–234.
- [5] Servais, C.: Sur les nombres parfaits. Mathesis 8 (1888), 135.
- [6] Sloane, N.J.A.: *The On-Line Encyclopedia of Integers Sequences*. Published online at: http://www.research.att.com/~njas/sequences, 2007.
- [7] Sylvester, J.J.: Sur les nombres parfaits. Comptes Rendus Paris 106 (1888), 403–405.
- [8] Touchard, J.: On prime numbers and perfect numbers. Scripta Math. 19 (1953), 35-39.
- [9] Weisstein, E.W.: Perfect Number. http://mathworld.wolfram.com/PerfectNumber.html

Luis H. Gallardo Mathematics University of Brest 6, Avenue Le Gorgeu, C.S. 93837 F-29238 Brest Cedex 3, France

e-mail: Luis.Gallardo@univ-brest.fr