Zeitschrift: Elemente der Mathematik

Herausgeber: Schweizerische Mathematische Gesellschaft

Band: 68 (2013)

Artikel: Structure of a group related to a Napoleon tesselation

Autor: Dlab, Vlastimil

DOI: https://doi.org/10.5169/seals-515891

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 29.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Elemente der Mathematik

Structure of a group related to a Napoleon tessellation

Vlastimil Dlab

Vlastimil Dlab started his mathematical career 1956 in algebra at the Charles University in Prague. Then, he had positions at the University of Khartoum, at the Institute of Advanced Studies in Canberra, Australia, and finally at Carleton University in Ottawa, Canada. He visited mathematics departments all around the world and was elected Fellow of the Royal Society of Canada in 1977. Presently he is Professor Emeritus and Distinguished Research Professor at Carleton University and Professor Hospitus at Charles University.

The presentation of a two-dimensional plane using the repetition of a geometric shape with no overlaps and no gaps, the so-called tessellation, or tiling, appears as a natural structure in the real world and is displayed frequently in arts since ancient times. It is the Moorish architecture in Alhambra, and, in modern times, the work of M. C. Escher that made this concept a familiar notion.

Here, we are interested in "periodic" tessellations, the beginnings of whose studies can be traced to their classification in terms of seventeen different groups of isometries by Evgraf Stepanovich Fedorov in [3]. This classification, rediscovered later by a number of algebraists, crystallographers and geometers, provides a beautiful bond between group theory and geometry of the plane. On the one hand, theory of groups furnishes an abstract description of all possible plane patterns in terms of their symmetries and on the other hand, a particular tessellation can render a complete structure of a group given by generators and relations. The present note relates to the latter category, viz. it describes the structure of a group given by a very simple presentation exploring its identification with the symmetry

In den kunstvoll gestalteten Ornamenten der Alhambra findet man alle Parkettierungsarten, welche den 17 ebenen kristallographischen Gruppen entsprechen. Der Zusammenhang zwischen diesen endlichen Gruppen und den entsprechenden Parkettierungen der Ebene ist schon lange bakannt und reflektiert die Maxime, dass Gruppen Symmetrien von Objekten darstellen. Inspiriert vom Satz von Napoleon werden Napoleon-Parkettierungen erzeugt von einem beliebigen Dreieck und den über dessen Seiten errichteten gleichseitigen Dreiecken. Sie liefern eine anschauliche Beschreibung der Gruppe $G=\langle a,b \mid a^3=b^3=(ab)^3=1\rangle$: Die Gruppe wird durch ein Bild ersetzt, wodurch technische Argumente visuell illustriert werden können.

group of a particular tessellation. The group in question is denoted by **p3** in Hermann-Mauguin notation that has been adopted by the International Union of Crystallography and used by Coxeter-Moser in their monograph [2].

The symmetry group **p3** relates to a tessellation generated by a general (scalene) triangle using the Napoleon theorem construction. In the final remark of this note, we briefly discuss the special cases of triangles with additional symmetries.

Thus, we wish to present a comprehensive description of the structure of the following (polyhedral) group

$$G = \langle a, b \mid a^3 = b^3 = (ab)^3 = 1 \rangle$$
 (*)

using a geometric representation, and thus demonstrating the maxim that groups are symmetries of objects. The paper translate the group into a picture that allows to replace technical arguments with visual demonstrations. First, let us formulate the main theorem.

Theorem 1. The group defined by (*) is isomorphic to the semidirect product $H \times K$, where H is the free abelian group of rank 2 and K the cyclic group of order 3: $H = \langle a^2b \rangle \times \langle ba^2 \rangle$ and $K = \langle a \rangle$. The semidirect product is provided by the isomorphism $\varphi: K \to \operatorname{Aut}(H)$ defined by $\varphi_a(a^2b) = aba$ and $\varphi_a(ba^2) = a^2b$.

Note that $aba \in H$ since $aba = a(ba)^{-2} = a(a^2b^2)^2 = (b^2a)(ab^2) = (a^2b)^{-1}(ba^2)^{-1}$. It is also easy to check that H is in fact abelian and normal in G. Moreover, $(a^2b)^na = a(a^2b)^{-n}(ba^2)^{-n}$ and $(ba^2)^na = a(a^2b)^n$. In view of these simple identities, one can define for every element $g = a^{e_1}b^{e_2}a^{e_3}b^{e_4}\dots a^{e_{d-1}}b^{e_d} \in G$ the weight w by $w(g) = \sum_{k=1}^d e_k$. Every word g with weight w(g) divisible by three is in H and the index of H in G is thus at most three.


The proof of Theorem 1 follows immediately from the representation of G as a symmetry group of a **general Napoleon tessellation**. This tessellation is related to the following theorem, often called the **Napoleon Theorem** (see e.g. Coxeter [1] or Rigby [4]): Given an arbitrary triangle $\Delta = ABC$, erect on its sides (externally) the equilateral triangles $\Delta_1 = BPC$, $\Delta_2 = CQA$ and $\Delta_3 = ARB$. Denote by U, V and W the centroids of these triangles. Then the triangle $\Delta_0 = UVW$ is equilateral and its centroid G coincides with the centroid of the original triangle Δ (see Figure 1).

The Napoleon Theorem provides immediately a (Napoleon) tessellation (wallpaper) indicated in Figure 2.

In particular, the centroids of the equilateral triangles define a new tessellation by equilateral triangles, all congruent to the triangle Δ_0 . This tessellation is indicated in Figure 2 by dotted lines. Note that six of such triangles with a common vertex (that is the centroid of one of the equilateral triangles of the Napoleon tessellation) form a regular hexagon.

In order to describe the geometric representation of the group G given by (*) as a **symmetry group** of this tessellation consider Figure 3, where the points U, V and W are described by the "coordinates" (0,1), (-1,1) and (0,0), respectively. Let us point out that we are using the centroids of the equilateral triangles of the Napoleon tessellation to define a coordinate system (indicated in Figure 3 by two bold dotted lines) that is not cartesian and has different scales in the two chosen directions. This system enables us to define translations of the plane that will represent some elements of our group.

24 V. Dlab



However, it will be rotations by $\frac{2\pi}{3}$ that will play a fundamental role in the geometric representation of the group. Indeed, some elements of the group (and, the generators a and b, in particular) will be represented by $\frac{2\pi}{3}$ rotations along the dotted hexagons of the previous Figure 2. Here we deal with the original angles on the plane, rather than with some formal angles in the new coordinates. This fact is emphasized in Figure 3 by long dotted lines at angle $\frac{\pi}{3}$.


Thus, denote by $\mathbf{R}_{(r,s)}$ the (anticlockwise) rotations of the plane around (r,s) by $\frac{2\pi}{3}$. These are, for all $r,s\in\mathbf{Z}$, symmetries (of order 3) of the considered Napoleon tessellation. Moreover, denote by $\mathbf{T}_{(0,0)\to(r,r+3t)}$ the translation of the plane mapping (0,0) to (r,r+3t). Observe that for $r,t\in\mathbf{Z}$, these are all the translations of the plane that provide symmetries of the tessellation. Finally, denote by \mathbf{S}_N the symmetry group of the given general Napoleon tessellation, i.e. the group generated by $\mathbf{R}_{(0,0)},\mathbf{R}_{(0,1)},\mathbf{R}_{(-1,0)},\mathbf{T}_{(0,0)\to(1,-2)},\mathbf{T}_{(0,0)\to(1,1)}$ and $\mathbf{T}_{(0,0)\to(2,-1)}$. Now, we can formulate the following theorem.

Theorem 2. The mapping $\psi: G \to \mathbf{S}_N$ given by

$$\psi(a) = \mathbf{R}_{(0,0)}$$
 and $\psi(b) = \mathbf{R}_{(0,1)}$

is an isomorphism of the groups G and S_N .

Proof. First, one can see easily that $\psi(a)\psi(b)=R_{(-1,1)}^2$ is of order three. Therefore, ψ is a well-defined group epimorphism. The generators of H are mapped to translations:



 $\psi(a^2b) = \mathbf{T}_{(0,0) \to (1,-2)}$ and $\psi(ba^2) = \mathbf{T}_{(0,0) \to (1,1)}$. More generally, $\psi[(a^2b)^t(ba^2)^r] = \mathbf{T}_{(0,0) \to (r+t,r-2t)} = \mathbf{T}_{(0,0) \to (1,-2)}^t \mathbf{T}_{(0,0) \to (1,1)}^r$. Thus ψ maps H bijectively onto the lattice of translations generated by the vectors (1,-2) and (1,1) (see Figure 3). The note following Theorem 1 together with the simple fact that the rotation $\mathbf{R}_{(0,0)}$ cannot be one of the translations imply that ψ is a bijection.

For completeness, we remark that the elements of G other than those from H belong to one of the following two cosets. The elements with $w(g) \equiv 1 \mod 3$ correspond to anticlockwise rotations by $\frac{2\pi}{3}$:

$$\psi[(a^2b)^r(ba^2)^{r+s}a] = \psi[a(a^2b)^s(ba^2)^{-r}] = \mathbf{R}(r,s),$$

while those with $w(g) \equiv 2 \mod 3$ are mapped to clockwise rotations:

$$\psi[(a^2b)^{-s}(ba^2)^r a^2] = \psi[a^2(a^2b)^{-r}(ba^2)^{-r-s}] = \mathbf{R}^2(r,s).$$

Observe that $\psi[(a^2b)^t(ba^2)^{r+t}] = \mathbf{T}_{(0,0)\to(r,r+3t)}$. All these statements are readily obtained by routine calculations facilitated by the fact that H is abelian, making use of the earlier mentioned identities $(a^2b)^na = a(a^2b)^{-n}(ba^2)^{-n}$, $(ba^2)^na = a(a^2b)^n$ and similar ones. Of course, the abelian normal subgroup H consists of all elements whose weight is a multiple of 3, and they correspond to all translations of \mathbf{S}_N .

Remark. Theorem 1 follows readily: H corresponds to the (normal) subgroup of all translations of \mathbf{S}_N and $K = \langle \psi^{-1}(\mathbf{R}_{(0,0)}) \rangle$.

V. Dlab




Figure 3

Final remark. The group G is the symmetry group of a **scalene** Napoleon tessellation: The sides of the triangle $\triangle = ABC$ have different lengths. Indeed, in the case that the triangle is an **isosceles** triangle (i.e. when $\overline{AC} = \overline{BC} \neq \overline{AB}$), the respective symmetry group $\overline{G} = \langle b, r \mid b^3 = r^2 = (rbrb^2)^3 = 1 \rangle$ is a semidirect product of the (normal)

subgroup $G = \langle b, rbrb^2 \rangle$ and the cyclic group $\langle r \rangle$ of order 2. Here, using earlier notation, $rbrb^2 = a$. Furthermore, r corresponds to the reflection with respect to the axis defined by the points W and C. Thus there is a tower of normal subgroups: $H = \langle (brb)^2, (b^2r)^2 \rangle \lhd G = \langle b, rbr \rangle \lhd \overline{G}$. Observe that H is contained in three subgroups of index 2 in \overline{G} . One of these subgroups, viz. $A = H \times \langle r \rangle$ is a maximal abelian subgroup of \overline{G} ; the other two $\langle H, brb^2 \rangle$ and $\langle H, rbrb^2 r \rangle$ are non-commutative. In other words, $\overline{G} \simeq H \rtimes S_3$, where $S_3 = \langle a, r \rangle$.

The group \overline{G} is standardly denoted by **p31m**. In case when the "defining" triangle is equilateral, the Napoleon tessallation is just the regular triangular tessellation, and its symmetry group is **p6m** (see [2]; the reader should beware that the notation in Table 3 differs from the standard one in that the group **p31m** is denoted by **p3m1** and vice versa).

Acknowledgement. The author wishes to acknowledge referee's suggestions, as well as fruitful discussions with Dr. Tomáš Pospíchal, that both lead to several improvements in the presentation of the paper.

References

- [1] H.S.M. Coxeter, Introduction to Geometry, John Wiley & Sons, New York London, 1961.
- [2] H.S.M. Coxeter and W.O. Moser, *Generators and Relations for Discrete Groups*, Ergebnisse der Math. Band 14, Springer-Verlag, New York 1965.
- [3] E.S. Fedorov, The Symmetry of Regular Systems of Figures, Proc. St. Petersburg Mineral. Soc. **28** (1891), 1–146
- [Eng. transl. in Symmetry of Crystals, Amer. Crystall. Assoc. Monograph 7 (1971), 50–131].
- [4] J.F. Rigby, Napoleon, Escher, and Tessellations, Math. Magazine, 64 (1991), 242-246.

Vlastimil Dlab
Katedra didaktiky matematiky
Matematicko-fyzikální fakulta Karlovy university
Praha, Czech Republic
e-mail: dlab@karlin.mff.cuni.cz
and
School of Mathematics and Statistics

Carleton University Ottawa, Canada

e-mail: vdlab@math.carleton.ca