

# On the mean length of the diagonals of an n-gon

Autor(en): **Lükő, Gábor**

Objektyp: **Article**

Zeitschrift: **Elemente der Mathematik**

Band (Jahr): **68 (2013)**

PDF erstellt am: **26.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-515892>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

---

---

## On the mean length of the diagonals of an $n$ -gon

---

---

Gábor Lükő

Gábor Lükő received his doctoral degree in mathematics from Loránd Eötvös University in Budapest in 1974. His interests include discrete geometry, theory of numbers, and functional analysis. Until 2002 he was working in several research institutes taking part in projects concerning industrial optimization.

### 1 Introduction

P. Erdős [1] conjectured that the mean length of the diagonals of a convex  $n$ -gon with perimeter  $L$  is maximal iff  $n/2$  vertices are concentrated in a point  $A$  and the remaining  $n/2$  vertices in a point  $B$  whose distance is  $L/2$  in case of  $n = 2k$ . If  $n = 2k + 1$ , then  $k$  and  $k + 1$  vertices are concentrated in  $A$  and  $B$ , respectively, in the extremal figure. The minimum is attained when  $n - 1$  vertices are concentrated in  $A$  and a single one in  $B$ .

These conjectures have been proved in the author's Msc. thesis [3] and dissertation [4] shortly after they had been stated. In this paper we revisit our solution of the problem.

### 2 Notations, definitions and a lemma

**Notations.**  $\overline{PQ}$  denotes the distance of two points  $P$  and  $Q$ .

Vertices of the  $n$ -gon are denoted by  $P_1, \dots, P_n$ , the whole  $n$ -gon by  $(P)$ .

Bei welcher geschlossenen Kurve gegebener Länge ist die mittlere Länge von Sehnen maximal? Der Autor des vorliegenden Artikels beantwortete diese Frage von Wilhelm Blaschke and Luis Santaló 1965 in seiner Masterarbeit: Es ist der Kreis. In der selben Arbeit betrachtete der Autor ein diskretes Analogon und zeigte: Unter allen gleichseitigen Polygonen zeichnet sich das reguläre Polygon durch maximale mittlere Diagonalenlänge aus. Bereits früher lösten Stephen Vincze und Karl Reinhardt unabhängig voneinander das Problem, unter allen konvexen gleichseitigen Polygonen diejenigen von minimalem Durchmesser zu identifizieren. Dabei treten Reuleaux-Polygone auf. Paul Erdős wollte daraufhin wissen, welche konvexen Polygone von gegebenem Umfang maximale respektive minimale mittlere Länge von Diagonalen besitzen. Diese Frage wird in der vorliegenden Arbeit beantwortet.

$V$  denotes the set  $\{P_1, P_2, \dots, P_n\}$ .

Multiple points (e.g.  $P_1 = P_2$ ) are allowed. If  $i$  is an arbitrary integer lying outside of the interval  $[1, n]$ , then  $P_i$  is defined to be equal to  $P_{i'}$ , where  $i' \equiv i \pmod{n}$ .

Vertices of  $(P)$  are considered to form a cycle  $C = (V, \Gamma)$ , where  $\Gamma : i \mapsto i + 1$ .

**Definitions.** The anterior vertex of  $P_i$  in  $(P)$  is the last  $P_j$  preceding  $P_i$  in  $C$  not coincident with it. It is denoted by  $A_j$ .

The successor of  $P_i$  in  $(P)$  is the first vertex  $P_j$  following  $P_i$  in  $C$  not coincident with it. It is denoted by  $S_j$ .

The angle  $\alpha_i$  at vertex  $P_i$  is called the angle between the segments  $P_i A_i$  and  $P_i S_i$ .

$P_i$  is called to be a break-point if  $\alpha_i \neq \pi$ .

**Notations.** The length of the diagonal  $P_i P_{i+l}$  is denoted by  $r_{i,l}$ .

$l$  is called the order of  $P_i P_{i+l}$ .

So the mean length of the diagonals of order  $l$  is

$$Q_l = \frac{1}{n} \sum_{i=1}^n r_{i,l}. \quad (1)$$

For the mean length  $Q$  of the diagonals of  $(P)$  we have obviously

$$Q = \begin{cases} \frac{1}{k-1} \sum_{l=2}^k Q_l & \text{if } n = 2k + 1, \\ \frac{1}{k-2+\frac{1}{2}} \left( \sum_{l=2}^{k-1} Q_l + \frac{Q_k}{2} \right) & \text{if } n = 2k. \end{cases}$$

Our results depend on the following

**Lemma.** Let  $l$  be a natural number from the interval  $[1, \lfloor \frac{n}{2} \rfloor]$ . Then we have

$$Q_l \leq \frac{l}{n} L. \quad (2)$$

Equality occurs in (2) iff at least  $l$  points are concentrated in every break-point of  $(P)$ .

### 3 Statement of the results

**Theorem 1.** The inequality

$$Q \leq \sigma L \quad (3)$$

holds, where

$$\sigma = \begin{cases} \frac{1}{4} \frac{n+3}{n} & \text{if } n = 2k + 1, \\ \frac{1}{4} \frac{n^2 - 8}{n(n-3)} & \text{if } n = 2k, \end{cases}$$

the sign of equality occurs only if at least  $\lceil \frac{n}{2} \rceil$  points are concentrated at every break-point  $P_i$  of  $(P)$ . Inequality (3) is sharp. If  $n = 2k$ , then equality holds iff  $k$  points are concentrated in each endpoint of a segment  $AB$  of length  $L/2$ . If  $n = 2k + 1$  then similarly, the remaining one point lying anywhere on  $AB$ .

Our next result is related to the minimum of  $\varrho$ .

**Theorem 2.** *If  $(P)$  is convex, then we have the relation*

$$\varrho \geq \frac{L}{n}. \quad (4)$$

*This inequality is sharp too. The sign of equality occurs iff  $n - 1$  vertices are concentrated in one of the endpoints  $A$  of a segment  $AB$  and one in  $B$ .*

## 4 Proofs

*Proof of the lemma.* Obviously

$$r_{i,l} \leq r_{i,1} + r_{i+1,1} + \dots + r_{i+l-1,1}. \quad (5)$$

Summing up (5) we get (2). The sign of equality occurs in (2) iff the same is true for (5) for every index  $i$ . This happens iff  $P_i, P_{i+1}, \dots, P_{i+l}$  are collinear and  $P_{i+1}, P_{i+2}, \dots, P_{i+l-1}$  lie on the segment  $[P_i, P_{i+l}]$  for every index  $i$ . But this is possible iff at least  $l$  vertices are concentrated at every break-point of  $(P)$ .  $\square$

*Proof of Theorem 1.* First we note that every  $(P)$  has at least two break-points – otherwise all the vertices were concentrated in one point – contradicting to the assumption  $L > 0$ .

Relation (3) follows at once from (2) by summation. Equality occurs in (3) iff we have “=” in (1) for every  $l$  ( $l = 1, \dots, \lceil \frac{n}{2} \rceil$ ).

Regarding the lemma, this case occurs iff at least  $l$  break-points are concentrated in every break-point of  $(P)$  ( $l = 1, \dots, \lceil \frac{n}{2} \rceil$ ). Strongest of these conditions is the last one implying that  $(P)$  may have at most two break-points.

Let us see which figures  $(P)$  satisfy this extremum condition.

If  $n = 2k$ , all the vertices of  $(P)$  have to be concentrated in two break-points  $A$  and  $B$  lying at distance  $L/2$  each from other.

If  $n = 2k + 1$ , then  $k$  vertices are concentrated both in  $A$  and  $B$ , the remaining one may lie anywhere on the segment  $AB$ .

In the extremal case we have for  $n = 2k + 1$

$$\varrho = \frac{2 + \dots + k}{k-1} \frac{L}{n} = \frac{1}{4} \frac{n+3}{n} L,$$

and for  $n = 2k$

$$\varrho = \frac{2 + \dots + k - 1 + \frac{k}{2}}{k - 2 + \frac{1}{2}} \frac{L}{n} = \frac{(n+2)(n-4) + 2n}{4n(n-3)} L. \quad \square$$

*Proof of Theorem 2.* Consider the quadrangle  $P_i P_{i+1} P_{i+l} P_{i+l+1}$ . It is convex in consequence of the convexity of  $(P)$  and so we have

$$r_{i,l} + r_{i+1,l} \geq r_{i+1,l-1} + r_{i,l+1}. \quad (6)$$

Summing up (6) we get

$$q_l \geq \frac{q_{l-1} + q_{l+1}}{2} \quad \left( l = 2, 3, \dots, \left[ \frac{n}{2} \right] \right) \quad (7)$$

i.e. the sequence  $\{q_l\}_1^n$  is concave. Taking into consideration that obviously

$$q_l = q_{n-l} \quad (l = 1, \dots, n)$$

(7) implies that

$$q_1 \leq q_2 \leq \dots \leq q_{\lfloor \frac{n}{2} \rfloor},$$

i.e.

$$q_l \geq q_1 \quad \left( l = 2, 3, \dots, \left[ \frac{n}{2} \right] \right)$$

implying

$$q \geq q_1 = \frac{L}{n},$$

proving (4). If  $n = 2k + 1$ , then in the extremal case we have

$$q = \frac{k-1}{k-1} \frac{L}{n} = \frac{L}{n}.$$

In the case  $n = 2k$  we obtain

$$q = \frac{k-2 + \frac{1}{2}}{k-2 + \frac{1}{2}} \frac{L}{n} = \frac{L}{n}. \quad \square$$

### Remarks

1. Theorem 1 remains valid in higher dimensions too – moreover in any metric space.
2. K. Böröczky gave another proof for Theorem 2 on L. Fejes Tóth's seminar.
3. If the sum of squared lengths of the sides of an  $n$ -gon is given, then the sum of squared lengths of its diagonals is maximal in case of the affine images of the regular  $n$ -gon [4].
4. Open questions:
  - What  $n$ -gons are extremal if the length of every side is given?
  - What can be said about the diameter when the perimeter or every length of side is fixed?
  - What is the minimum of the mean squared length of the diagonals of an  $n$ -gon if the mean squared length of the sides is given assuming convexity of the  $n$ -gon?

**Acknowledgements.** I am indebted to professors I. Vincze and E. Makai who have read the manuscript and called my attention on some inaccuracies.

### References

- [1] Erdős, P.: *Oral communication*, 1966.
- [2] Larcher, G.; Pillichshammer, F.: The sum of distances between vertices of a convex polygon with unit perimeter. *Amer. Math. Monthly* 115 (2008), 350–355.
- [3] Lükő, G.: On the mean length of the chords of a closed curve. Msc. thesis, Attila Jozsef University, 1965 (in Hungarian).
- [4] Lükő, G.: Extremum problems related to closed curves, polygons and polyhedra. Doctorial dissertation, Lorand Eötvös University, 1972 (in Hungarian).
- [5] Pillichshammer, F.: On the sum of squared distances in the Euclidean plane. *Arch. Math.* 74 (2000), 472–480.
- [6] Pillichshammer, F.: A note on the sum of distances under a diameter constraint. *Arch. Math.* 77 (2001), 195–199.
- [7] Vincze, S.: On a geometrical extremum problem. *Acta Sci. Math. (Szeged)* 12 (1950), 136–142.

Gábor Lükő  
Kankalin u. 29  
H–2000 Szentendre, Hungary  
e-mail: jgluko@gmail.com