

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 68 (2013)

Artikel: The Faà di Bruno formula revisited
Autor: [s.n.]
DOI: <https://doi.org/10.5169/seals-515893>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The Faà di Bruno formula revisited

Raymond Mortini

Raymond Mortini, geboren 1958 in Luxemburg, studierte Mathematik mit Nebenfach Physik in Karlsruhe. Nach der Promotion und Habilitation wurde er zum ausserplanmässigen Professor in Karlsruhe ernannt. 1995 folgte er einem Ruf auf eine ordentliche Professur in Metz. 1993 wurde er mit dem Carl Freudenberg Preis ausgezeichnet. Seine Hauptarbeitsgebiete umfassen die Funktionentheorie, Funktionalanalysis und Operatorentheorie. Verschiedene Forschungsaufenthalte in den USA, Spanien und an seinem Lieblingssort, dem Mathematischen Forschungsinstitut Oberwolfach, prägen seinen Werdegang.

1 How one may (re)discover oneself this formula

Let f and g be two functions on \mathbb{R} whose n -th derivatives exist. The first 4 derivatives of $f \circ g$ are easy to calculate:

$$\begin{aligned}(f \circ g)' &= (f' \circ g)g' \\(f \circ g)^{(2)} &= (f^{(2)} \circ g)(g')^2 + (f' \circ g)g^{(2)} \\(f \circ g)^{(3)} &= (f^{(3)} \circ g)(g')^3 + 3(f^{(2)} \circ g)g^{(2)}g' + (f' \circ g)g^{(3)} \\(f \circ g)^{(4)} &= (f^{(4)} \circ g)(g')^4 + 6(f^{(3)} \circ g)g^{(2)}(g')^2 + 4(f^{(2)} \circ g)g^{(3)}g' + 3(f^{(2)} \circ g)(g^{(2)})^2 + (f' \circ g)g^{(4)}\end{aligned}$$

In der Leibnizschen Regel für die höheren Ableitungen eines Produktes $f \cdot g$ von zwei Funktionen treten die Binomialkoeffizienten auf. Bei Produkten von mehr als zwei Faktoren sind es entsprechend die Multinomialkoeffizienten. Die Kettenregel für die höheren Ableitungen der Zusammensetzung $f \circ g$ zweier Funktionen auf \mathbb{R} weist hingegen eine kompliziertere Struktur auf. Die entsprechende Formel wurde erstmals vom vielseitigen italienischen Mathematiker Francesco Faà di Bruno 1855 publiziert und seither immer wieder untersucht. Die klassische Variante der Formel für die n -te Ableitung benützt eine etwas unbequeme Summation über die nichtnegativen Lösungen der diophantischen Gleichung $b_1 + 2b_2 + \dots + nb_n = n$, die bekanntlich in der Kombinatorik eine wichtige Rolle spielt. In der vorliegenden Arbeit wird Faà di Brunos Formel auf eine intuitive Weise hergeleitet und eine einfachere Summationsreihenfolge vorgeschlagen.

The expressions get rapidly longer and more complicated. For example we have 42 summands for $n = 10$. Note that the number of terms equals the partition number $p(n)$, that is the number of ways (without order) of writing the integer n as a sum of strictly positive integers; by the Hardy-Ramanujan formula we have $p(n) \sim \frac{1}{4n\sqrt{3}} e^{\pi\sqrt{2n/3}}$.

Let

$$\mathbf{M}^j = \{\mathbf{k} = (k_1, \dots, k_j) \in (\mathbb{N}^*)^j, k_1 \geq k_2 \geq \dots \geq k_j \geq 1\}$$

be the set of ordered multi-indices in $\mathbb{N}^* = \{1, 2, \dots\}$. If g is a function defined on \mathbb{R} , and if $g^{(n)}$ is the n -th derivative of g , then we denote by $g^{(\mathbf{k})}$ the function $\prod_{i=1}^j g^{(k_i)}$, where $\mathbf{k} = (k_1, \dots, k_j) \in \mathbf{M}^j$. Also, $g^{(0)}$ is, by convention, equal to the function g itself.

The classical Faà di Bruno formula from ca. 1850 gives an explicit formula for $(f \circ g)^{(n)}$:

$$(f \circ g)^{(n)}(x) = \sum \frac{n!}{b_1! b_2! \dots b_n!} f^{(j)}(g(x)) \prod_{i=1}^n \left(\frac{g^{(i)}(x)}{i!} \right)^{b_i},$$

where the sum is taken over all different solutions in nonnegative integers b_1, b_2, \dots, b_n of

$$b_1 + 2b_2 + \dots + nb_n = n \text{ and } j := b_1 + \dots + b_n.$$

A nice historical survey on this appeared in [2]. See also [1].

Without being aware of that formula, I developed around 1976–1980 the following formula:

$$(f \circ g)^{(n)}(x) = \sum_{j=1}^n f^{(j)}(g(x)) \left(\sum_{\substack{\mathbf{k} \in \mathbf{M}^j \\ |\mathbf{k}|=n}} C_{\mathbf{k}}^n g^{(\mathbf{k})}(x) \right), \quad (1.1)$$

where

$$C_{\mathbf{k}}^n = \frac{\binom{n}{\mathbf{k}}}{\prod_i N(\mathbf{k}, i)}.$$

Here $\binom{n}{\mathbf{k}}$ is the multinomial coefficient defined by $\binom{n}{\mathbf{k}} = \frac{n!}{k_1! k_2! \dots k_j!}$, where $|\mathbf{k}| := k_1 + \dots + k_j = n$, and $N(\mathbf{k}, i)$ is the number of times the integer i appears in the j -tuple \mathbf{k} ($i \in \mathbb{N}^*$ and $\mathbf{k} \in (\mathbb{N}^*)^j$).

For example, the coefficient $C_{(4,1,1)}^6$ of the term $g^{(4)}(g')^2$ when looking at the 6-th order derivative of $f \circ g$ is

$$C_{(4,1,1)}^6 = \frac{1}{2!} \frac{6!}{4! \cdot 1! \cdot 1!} = 15$$

and the coefficient $C_{(4,2,1,1,1,1)}^{10}$ of the term $g^{(4)}g''(g')^4$ in the 10-th derivative is

$$C_{(4,2,1,1,1,1)}^{10} = \frac{1}{4!} \frac{10!}{4! \cdot 2! \cdot 1! \cdot 1! \cdot 1! \cdot 1!} = 3150.$$

The difference between our formula and the Faà di Bruno formula is that we use a simpler summation order and do not consider exponents of the form $b_j = 0$. In particular, we do not need summation over those (b_1, \dots, b_n) satisfying (the difficult to grasp) condition $\sum_{i=1}^n i b_i = n$. That these two formulas are equivalent though, immediately follows from a direct comparison of the coefficients. Indeed, for fixed j and $\mathbf{k} = (k_1, \dots, k_j) \in \mathbf{M}^j$, $|\mathbf{k}| = n$, we have:

$$\begin{aligned} C_{\mathbf{k}}^n &= \left(\frac{n!}{k_1! \cdots k_j!} \right) \frac{1}{N(\mathbf{k}, 1)!} \cdots \frac{1}{N(\mathbf{k}, n)!} \\ &= \left(n! \underbrace{\frac{1}{i_1! \cdots i_1!}}_{b_{i_1} \text{ times}} \cdots \underbrace{\frac{1}{i_\ell! \cdots i_\ell!}}_{b_{i_\ell} \text{ times}} \right) \frac{1}{b_{i_1}! \cdots b_{i_\ell}!} = \frac{n!}{b_1! \cdots b_n!} \prod_{i=1}^n \left(\frac{1}{i!} \right)^{b_i} \end{aligned}$$

where the b_{i_m} are those exponents that are different from zero and where \mathbf{k} has been represented in the canonical form $\mathbf{k} = (\underbrace{i_1, \dots, i_1}_{b_{i_1} \text{ times}}, \dots, \underbrace{i_\ell, \dots, i_\ell}_{b_{i_\ell} \text{ times}})$ in decreasing order. Note

that

$$\sum_{i=1}^n i b_i = \sum_{s=1}^{\ell} i_s b_{i_s} = \sum_{p=1}^j k_p = n$$

and that

$$\sum_{i=1}^n b_i = b_{i_1} + \cdots + b_{i_\ell} = j.$$

Next I would like to present the (intuitive) steps that led me to the discovery of the formula (1.1) above, at pre-PC times; the first (non-programmable) slide rule calculator SR50 had just appeared.

1) I calculated explicitly the derivatives $(f \circ g)^{(n)}$ up to the order 10 and wrote them down in a careful chosen order (see Figure 1);

2) An immediate guess is that

$$(f \circ g)^{(n)}(x) = \sum_{j=1}^n f^{(j)}(g(x)) \left(\sum_{\substack{\mathbf{k} \in \mathbf{M}^j \\ |\mathbf{k}|=n}} c_{\mathbf{k}}^n g^{(\mathbf{k})}(x) \right),$$

for some coefficients $c_{\mathbf{k}}^n$ to be determined.

3) Next I gave an inductive proof that this representation is correct; that needs the main step of the construction: where does the factor $g^{(k_1)} \cdots g^{(k_j)}$ with $k_1 + \cdots + k_j = n + 1$ comes from? So let us look at the ordered j -tuple (k_1, \dots, k_j) . This tuple is generated, through anti-differentiation, by the j j -tuples

$$(k_1 - 1, k_2, \dots, k_j), (k_1, k_2 - 1, k_3, \dots, k_j), \dots, (k_1, \dots, k_j - 1);$$

that is,

$$\begin{aligned} \left(g^{(k_1-1)} g^{(k_2)} \dots g^{(k_j)} \right)' &= g^{(k_1)} g^{(k_2)} \dots g^{(k_j)} + \dots, \\ \left(g^{(k_1)} g^{(k_2-1)} \dots g^{(k_j)} \right)' &= g^{(k_1)} g^{(k_2)} \dots g^{(k_j)} + \dots, \end{aligned}$$

etc.

The main difficulty being that the components k_j are not pairwise distinct. So their ‘‘multiplicities’’ had to be taken into account. This led to the guess that one may have

$$c_1^1 = 1$$

$$c_{\mathbf{k}}^{n+1} = \sum_{i=1}^j \frac{N(\mathbf{k} - \mathbf{e}_i^j, k_i - 1)}{N(\mathbf{k}, k_i)} c_{\mathbf{k} - \mathbf{e}_i^j}^n$$

where for $i = 1, \dots, j$, $\mathbf{e}_i^j = (0, \dots, 0, \underbrace{1}_{i\text{-th}}, 0, \dots, 0)$, $\mathbf{e}_i^j \in \mathbb{N}^j$, $\mathbf{k} \in \mathbf{M}^j$, $|\mathbf{k}| = n + 1$

and $1 \leq j \leq n + 1$. Note that the j -tuple $\mathbf{k} - \mathbf{e}_i^j$ is not necessarily represented in the canonical form with decreasing coordinates. Also, if the i -th coordinate of \mathbf{k} is one, then the i -th coordinate of $\mathbf{k} - \mathbf{e}_i^j$ is 0 and we identify $\mathbf{k} - \mathbf{e}_i^j$ with the associated $(j - 1)$ -tuple. For example in the case $\mathbf{k} = (3, 1, 1, 1)$ we have

$$20 = C_{(3,1,1,1)}^6 = \frac{1}{1} C_{(2,1,1,1)}^5 + \frac{1}{3} C_{(3,0,1,1)}^5 + \frac{1}{3} C_{(3,1,0,1)}^5 + \frac{1}{3} C_{(3,1,1,0)}^5,$$

where $(3, 0, 1, 1)$, $(3, 1, 0, 1)$ and $(3, 1, 1, 0)$ are identified with $(3, 1, 1)$.

With these recursion formula the inductive proof went through.

4) Next one has to guess the explicit value of $c_{\mathbf{k}}^n$, $|\mathbf{k}| = n$. Now there are $\binom{n}{\mathbf{k}} = \frac{n!}{k_1! \dots k_j!}$ ways to choose k_1 objects out of n , then k_2 objects of the remaining ones, and so on. Due to the multiplicity, one has again to divide by $N(\mathbf{k}, i)!$.

This gives the guess that $c_{\mathbf{k}}^n = \frac{\binom{n}{\mathbf{k}}}{\prod_i N(\mathbf{k}, i)!}$.

5) These coefficients $c_{\mathbf{k}}^n$ actually satisfy the recursion relation above. Since $c_1^1 = 1$, and the fact that the recursion relation determines uniquely the next coefficients, we are done: $C_{\mathbf{k}}^n = c_{\mathbf{k}}^n$.

2 Further formulas and questions

Applying formula (1.1) for the function $f(x) = \log x$, $x > 0$ and $g(x) = e^x$ gives

$$\sum_{j=1}^n (-1)^{j-1} (j-1)! \sum_{\substack{\mathbf{k} \in \mathbf{M}^j \\ |\mathbf{k}|=n}} C_{\mathbf{k}}^n = 0, \quad n \geq 2;$$

There are actually infinitely many pairs of columns that coincide (just use that $C_{(2,2,2,i)}^{6+i} = C_{(4,2,i)}^{6+i}$ for every $i \geq 5$.)

Are there triples (or a higher number) of columns that coincide?

Acknowledgement. I thank Jérôme Noël for his help in finding the exact $\text{T}_{\text{E}}\text{X}$ -commands for creating the tabular form above. I also thank Claude Jung for valuable discussions around 1980.

References

- [1] Craik, Alex. Prehistory of Faà di Bruno's formula, Amer. Math. Monthly 112 (2005), 119–130.
- [2] Johnson, Warren. The curious history of Faà di Bruno's formula, Amer. Math. Monthly 109 (2002), 217–234.

Raymond Mortini
Département de Mathématiques
LMAM, UMR 7122, Université de Lorraine
Ile du Saulcy
F-57045 Metz, France
e-mail: mortini@univ-metz.fr