Zeitschrift: Elemente der Mathematik

Herausgeber: Schweizerische Mathematische Gesellschaft

Band: 69 (2014)

Artikel: Some hidden harmonies between new and old geometric loci

Autor: Ritelli, Daniele / Scimone, Aldo

DOI: https://doi.org/10.5169/seals-515868

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 29.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Elemente der Mathematik

Some hidden harmonies between new and old geometric loci*

Daniele Ritelli and Aldo Scimone

Daniele Ritelli teaches mathematical analysis and financial mathematics at the School of Economics, Management and Statistics at the University of Bologna. His research interests are classical analysis (special functions in analytic number theory) and applied mathematics (mathematical modeling through nonlinear ordinary differential equations and minimum cost models in production planning).

Aldo Scimone, PhD (Mathematics Didactics) has been a supervisor for mathematics teachers at SISSIS (University of Palermo). He is author of several books and articles on the history of mathematics. His current research interests centre on elementary number theory, geometry, the history of mathematics and mathematics didactics.

1 Introduction

Geometric loci have always fascinated mathematicians. Many of the most beautiful and famous curves represent many geometric loci which have given a remarkable push to the creation of most of the mathematics we know. At once a geometric locus is determined, mathematicians often wonder if it can also be derived in a different way, because, if it is possible, the geometric locus gains a deeper meaning. We consider the map $H(u, v) \mapsto$

Die Geschichte der Mathematik ist gespickt mit Entdeckungen von immer neuen geometrischen Örtern. Darunter sind solche, welche der Untersuchung bekannter Fragestellungen entsprangen, etwa den drei klassischen durch Zirkel und Lineal unlösbaren geometrischen Problemen der griechischen Antike. Bekannte Beispiele sind die Archimedische Spirale, die Konchoide von Nikomedes, die Zissoide von Diokles oder die Quadratrix des Hippias. Während Jahrhunderten blieben geometrische Örter beliebte Studienobjekte, und seit dem XVII. Jahrhundert erlauben die Werkzeuge der analytischen Geometrie und der Analysis die Entdeckung neuer oder die neue Interpretation altbekannter geometrischer Örter. Mit jeder neuen Methode oder Fragestellung ergeben sich auch heute noch immer wieder erhellende Zusammenhänge, so auch im vorliegenden Beitrag.

^{*}Dedicated to the memory of the unforgettable friend and Master Prof. Filippo Spagnolo




Fig. 1 The map $H(u, v) \mapsto P(x, y)$.

P(x, y), a composition of a rotation of an angle θ of a point H(u, v) upon the x-axis and a vertical projection of H' up to the point P(x, y) (Figure 1), given (by Pythagoras and Thales) by the relations

$$\begin{cases} x = \sqrt{u^2 + v^2} \\ y = \frac{v}{u}x = \frac{v}{u}\sqrt{u^2 + v^2}. \end{cases}$$
 (1)

In this paper we aim to show how, by this composition of a rotation and a vertical projection, it is possible to get already-known results, which sheds a new light on geometric objects.

2 The locus

Let r be a straight line referred to a cartesian coordinate system with A(a, 0), B(0, b) and consider the line t through the origin O of equation y = mx (Figure 2). Let H be the point of intersection between r and t. Since the equation of the straight line through A and B is bx + ay - ab = 0, the point H has the coordinates:

$$\begin{cases} u = \frac{ab}{ma + b} \\ v = \frac{mab}{ma + b}. \end{cases}$$

Now if we apply the map (1) to the point H, we obtain the point P whose coordinates are:

$$\begin{cases} x = \frac{ab\sqrt{1+m^2}}{ma+b} \\ y = \frac{mab\sqrt{1+m^2}}{ma+b}. \end{cases}$$
 (2)

Applying map (1) to the point H' of intersection between t and the straight line s parallel to r and passing through A' = (-a, 0) and B' = (0, -b) we obtain the point P'. Our




Fig. 2 The generation of the locus.

purpose is to study the geometric locus described by P and P' varying the straight line t. By symmetry we can study the case of point P. Since m = y/x, substituting into the first or second one of (2) we obtain

$$axy + bx^2 = ab\sqrt{x^2 + y^2}$$

from which, squaring, one has the quartic

$$b^{2}x^{4} + 2abx^{3}y + a^{2}x^{2}y^{2} - a^{2}b^{2}x^{2} - a^{2}b^{2}y^{2} = 0.$$
 (3)

It is symmetric to the point O which is an isolated double point. It has (see Figure 3) two vertical asymptotes whose equations are $x = \pm b$, while the oblique asymptotes are:

$$y = -\frac{b}{a}x \pm \frac{b\sqrt{a^2 + b^2}}{a}.$$

If in (3) we divide by a^2 and we take the limit as a goes to infinity we obtain the curve of equation

$$x^2y^2 - b^2x^2 - b^2y^2 = 0 (4)$$

which is the equation of a famous quartic named for its form: cross curve (Figure 4). In general, the equation of the cross curve is

$$x^2v^2 - a^2x^2 - b^2v^2 = 0$$

which, for a=b, is said to be *equilateral*, and it assumes the form of (4). It is also represented by the form

$$\frac{a^2}{y^2} + \frac{b^2}{x^2} = 1.$$




Fig. 3 The quartic (3) with a = 3 and b = 4.




Fig. 4 The cross curve with a = 3 and b = 4.

The origin O is an isolated double point and the points at infinity on the axes are knots of inflection. The real points of the curve are external to the strips limited by the equations $x = \pm a$ and $y = \pm b$, which, in our case, (the equilateral one) are $x = \pm b$ and $y = \pm b$. This curve has been studied since 1847 in a question proposed by the French mathematician Olry Terquem in [3], and, thirteen years later, in a paper by the Italian mathematician Angelo Francesco Siacci, who demonstrated a fine property of the cross curve: the sum of the areas limited by the four branches of the curve and by the respective asymptotes is equal to the area of the rectangle formed by the same asymptotes. Siacci's proof of this




Fig. 5 The proof of the property of the Siacci cross curve.

property is sharp, [2] pp. 199–200, and makes use of calculus. Here we present a simpler geometric argument, brought to our attention by the anonymous referee whom we acknowledge. It is carried out as follows (Figure 5): multiply $x^2 - u^2 = b^2$ by the constant HO^2/u^2 to obtain, by Thales, $PO^2 - HO^2 = GO^2$. Hence, for any infinitesimally thin slice between two half-lines through O, the areas between PH and GO are the same. Therefore, the integrals are also the same.

The curve was also studied in [1] by the great Italian mathematician Ernesto Cesàro who named it *stauroid* from the Greek word $\sigma\tau\alpha\nu\rho\sigma\epsilon\iota\delta\eta\zeta$ which means crux-form, just for its form. Now if we divide (3) by b^2 and take the limit for $b\to\infty$ we obtain the quartic

$$x^4 - a^2x^2 - a^2y^2 = 0 (5)$$

whose graph is illustrated in Figure 6.




Fig. 6 The *kampyle* curve with a = 3.

The curve intersects the x-axis at (-a, 0) and (a, 0) and it is the famous kampyle curve $(\chi \alpha \mu \pi v \lambda \eta \gamma \rho \alpha \mu \mu \eta)$ which, according to Paul Tannery, was used by the great Greek mathematician Eudoxus of Cnidus to perform on the plane the construction devised by his teacher Archytas of Tarentum in order to solve the famous problem duplication of cube, known as the Delian problem. We recall the statement of the problem: given the edge of a cube, construct with compasses and a straight edge alone the edge of a second cube having double the volume of the first. Archytas of Tarentum devised a solution without compasses and a straight edge which is briefly as follows: his construction is not on a plane but a bold construction in three dimensions, determining a certain point as the intersection of three surfaces of revolution – a right cone, a cylinder, and a torus. The intersection of the two latter surfaces gives (says Archytas) a certain curve (which is, in fact, a curve of double curvature), and the point required is found where the cone meets this curve. Eudoxus of Cnidus was searching for a way to perform the construction devised by Archytas on a plane, and he obtained the kampyle curve.

We can obtain the kampyle curve using the map (1) again. In fact, let us consider two parallel straight lines r and s of equations x = a and x = -a respectively, and a straight line y = mx. Let us apply map (1) to the points H(u, v) and H'(-u, -v) (Figure 7).




Fig. 7 A new generation of the kampyle curve.

By symmetry we consider only the point H, so for the coordinates of P one has:

$$\begin{cases} x = a\sqrt{m^2 + 1} \\ y = ma\sqrt{m^2 + 1}. \end{cases}$$
 (6)

Substituting y = mx into (6) we obtain

$$x^2 = a\sqrt{x^2 + y^2}$$

so that by squaring we get the quartic

$$x^4 - a^2x^2 - a^2y^2 = 0$$

that is the equation of the kampyle curve.

The cross curve and the kampyle are reduced to the two isotropic straight lines arising from the origin both when one divides (4) by b^2 (with $b \to \infty$) and when we divide (5) by a^2 (with $a \to \infty$).

3 The case of the ellipse

Now we map (1) to the point H(u, v) of intersection between the ellipse with semi-axes a and b of the equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

and the straight line r of the equation y = mx and the point P(x, y).




Fig. 8 The case of the ellipse.

Since:

$$\begin{cases} u = \frac{ab}{\sqrt{a^2m^2 + b^2}} \\ v = \frac{mab}{\sqrt{a^2m^2 + b^2}} \end{cases}$$

the action of map (1) gives

$$\begin{cases} x = ab\sqrt{\frac{1+m^2}{a^2m^2+b^2}} \\ y = mab\sqrt{\frac{1+m^2}{a^2m^2+b^2}}. \end{cases}$$
 (7)

Since y = mx, substituting into the one of the two equations in (7) we obtain the quartic

$$b^{2}x^{4} + a^{2}x^{2}y^{2} - a^{2}b^{2}x^{2} - a^{2}b^{2}y^{2} = 0$$
 (8)




Fig. 9 The quartic (8) with a = 3 and b = 4.

which has, at the origin, an isolated double point with two vertical asymptotes of the general equation $x = \pm b$, whose graph is represented in Figure 9.

If in (8) we divide by a^2 and $a \to \infty$, we get the curve of equation

$$x^2y^2 - b^2x^2 - b^2y^2 = 0$$

that is, again, the equilateral cross curve. So, starting from simple geometric considerations, two famous curves historically obtained by other methods, both appear now as particular cases of general quartic curves. This offers a unitary vision and shows how often in mathematics subjects with no apparent linkage can find common significance with a mere change of the viewpoint.

Acknowledgement. We wish to thank the anonymous referee for his fruitful suggestions which have improved our article.

References

- Cesàro, E. (1901), Sull'uso delle condizioni di immobilità in geometria intrinseca, Le matematiche pure ed applicate, Città di Castello: I, 241–254.
- [2] Siacci, F. (1860), Intorno ad una linea di quart'ordine, Giornale arcadico di Scienze Lettere ed Arti, Roma, Tomo CLXVII, 181–201.
- [3] Terquem, O. (1847), Question 165, Nouvelles Annales des mathématiques, 1re série, Paris: tome 6, 394.

Daniele Ritelli Dipartimento di scienze statistiche via Belle Arti, 41 I-40126 Bologna Italy daniele.ritelli@unibo.it Aldo Scimone Via Nigra, 30 I-Palermo Italy aldo.scimone@libero.it