

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 69 (2014)

Artikel: Some remarks about Descartes' rule of signs
Autor: Albouy, Alain / Fu, Yanning
DOI: <https://doi.org/10.5169/seals-515869>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Some remarks about Descartes' rule of signs

Alain Albouy and Yanning Fu

Alain Albouy got his doctorate in mathematics at Paris 7 university in 1992. Since this date he is a full-time researcher at CNRS and Paris Observatory. His research work is mainly about geometrical aspects of classical celestial mechanics.

Yanning Fu got his doctorate in celestial mechanics at Nanjing University in 1997. He is a researcher in astronomy at Purple Mountain Observatory and a professor of celestial mechanics at the University of the Chinese Academy of Sciences. His research interests include the n -body problem and ephemeris astronomy.

Introduction

What can we deduce about the roots of a real polynomial in one variable by only considering the signs of its coefficients?

If we restrict our attention to the positive roots, the answer is: nothing more than what the classical Descartes rule of sign states. This is a precise mathematical claim that needs a proof. We give it in Section 3, after some due preliminaries.

If we are interested in the negative roots also, and concentrate on the combined numbers of positive and negative roots, there is nothing as simple to tell. In Section 5, we show how the situation complicates with the degree, by giving a complete study up to degree 6.

Section 4 is a little interlude about discriminants and trinomials.

Die Vorzeichenregel von Descartes besagt: Die Anzahl P der positiven Nullstellen (gezählt mit ihrer Vielfachheit) eines reellen Polynoms $p(x)$ ist gleich der Zahl der Vorzeichenwechsel seiner Koeffizientenfolge oder um eine gerade natürliche Zahl kleiner als diese. So hat insbesondere ein Polynom mit nur einem Vorzeichenwechsel genau eine positive Nullstelle. Durch Betrachten von $p(-x)$ ergibt sich eine entsprechende Aussage für die Anzahl N negativer Nullstellen von $p(x)$. Umgekehrt kann man fragen, ob in jedem Fall ein Polynom mit gegebener Vorzeichenfolge existiert, welches eine damit kompatible Anzahl (P, N) positiver respektive negativer Nullstellen aufweist. Die Autoren behandeln diese und andere Fragen und verallgemeinern zudem die Regel von Descartes auf Linearkombinationen beliebiger reeller Potenzen.

1 Coefficients and positive roots

It is convenient to consider that an analytic function in one variable has a *multiset* of roots. A multiset is defined just as a set, except that identical elements are allowed. The *cardinality* of a multiset is the number of its elements. The *multiplicity* of an element is the cardinality of the sub-multiset of the elements identical to it.

Consider expressions of the type

$$Y = a_0x^{\alpha_0} + a_1x^{\alpha_1} + \cdots + a_nx^{\alpha_n}, \quad (1)$$

with real exponents $\alpha_0 < \alpha_1 < \cdots < \alpha_n$ and non-zero real coefficients a_0, \dots, a_n . The *sequence of signs* of an expression (1) is the ordered list $\sigma_0, \dots, \sigma_n$ of the signs $\sigma_i = |a_i|^{-1}a_i$. A *variation* occurs in such a sequence when $\sigma_i = -\sigma_{i+1}$. For example the sequence 1, -1, 1, 1 has two variations.

Proposition 1. *On one hand, let us prescribe the number of terms, the exponents and the signs of the coefficients in an expression (1), and denote by k the number of variations of the sign sequence. On the other hand, let us give arbitrarily a multiset of positive numbers, and call p its cardinality. There exists an expression (1), as prescribed, whose multiset of positive roots is the given one, if and only if $k - p$ is even and non-negative.*

This statement gives an “if and only if” form to the classical Descartes rule of signs. It continues previous works. In 1998, Anderson, Jackson and Sitharam [1] proposed examples of polynomials with any sequence of signs, and with any number p of positive roots, provided that $k - p$ is even and non-negative. In 1999, Grabiner [5] extended this result by giving also examples of polynomials with missing terms (i.e., zero coefficients for some of the intermediate integer powers of x). Our Proposition 1 strengthens this statement, by extending these previous results to real exponents, and by showing that when a given sequence of signs allows p positive roots, it indeed allows any multiset of positive roots of cardinality p . Surprisingly the proof, as presented below, remains very elementary.

2 Proof of the “only if” part

This proof is well known and we give it only for completeness. We assume that the given multiset is the multiset of positive roots of Y .

- (i) The parity of k decides if the sign of a_0 is or is not the sign of a_n . The parity of p decides if the sign of Y is or is not the same at the two ends, i.e., at 0^+ and at $+\infty$. Obviously these boundary signs are respectively the signs of a_0 and of a_n . Thus $k - p$ is even.
- (ii) Let m and m' be the cardinalities of the respective multisets of positive roots of a function f and of its derivative f' . Then $m \leq m' + 1$.
- (iii) One multiplies Y by $x^{-\alpha}$, computes the derivative in x and observes the resulting sequence of signs. The signs corresponding to the exponents $\alpha_i < \alpha$ are changed, while those corresponding to $\alpha_i > \alpha$ are not changed. We can delete any given variation of sign without touching the others by choosing α in some interval of the form $] \alpha_i, \alpha_{i+1} [$.

To conclude, we prove that $p \leq k$ by an induction on k . If $k = 0$ then $p = 0$. If Y has k variations, argument (iii) shows that $(x^{-\alpha}Y)'$ has $k - 1$ variations for some choice of α . Argument (ii) gives the required estimate on the respective numbers of roots. \square

These arguments are actually older than usually thought. Argument (ii) was given in 1741 by de Gua and again in 1798 by Lagrange. Argument (iii) and the statement of the “only if” part are given by de Gua in the case of integer exponents, by Laguerre in the case of real exponents¹. Laguerre is explicit about the parity conclusion (i), which is obvious. If the statement of the “only if” part omits this conclusion, then it remains correct if the roots are counted without multiplicity².

3 Proof of the “if” part

We call $(\sigma_0, \dots, \sigma_n)$ the prescribed sequence of signs (with k variations). We assume that the cardinality p of the given multiset of positive numbers is such that $k - p$ is non-negative and even. We look for an expression (1) with $\sigma_i a_i > 0$ for all i , such that the given multiset is the multiset of its positive roots. We will be able to find such an expression with the additional constraint that $a_i = a_{i+1}$ when $\sigma_i = \sigma_{i+1}$. We collect the terms in (1) according to this constraint. We set $i_0 = 0$ and call i_1, i_2, \dots, i_k the integers i_j such that $i_{j-1} < i_j$, $\sigma_{i_{j-1}} \neq \sigma_{i_j}$, $\sigma_{i_k} = \sigma_n$. We call b_0, b_1, \dots, b_k the common values of the coefficients:

$$b_j = a_{i_j} = a_{i_{j+1}} = \dots = a_{i_{j+1}-1}.$$

We set

$$\varphi_j(x) = x^{a_{i_j}} + x^{a_{i_j+1}} + \dots + x^{a_{i_{j+1}-1}}.$$

Now we look for a (b_0, b_1, \dots, b_k) such that $\sigma_0 b_0 > 0$, $b_i b_{i+1} < 0$ and the expression

$$Y = b_0 \varphi_0(x) + b_1 \varphi_1(x) + \dots + b_k \varphi_k(x)$$

has the required multiset of positive roots.

We begin with the case where all the positive roots x_1, \dots, x_p are simple. The column vector (b_0, \dots, b_k) should be in the kernel of the matrix

$$\Phi = \begin{pmatrix} \varphi_0(x_1) & \varphi_1(x_1) & \dots & \varphi_k(x_1) \\ \varphi_0(x_2) & \varphi_1(x_2) & \dots & \varphi_k(x_2) \\ & & \dots & \\ \varphi_0(x_p) & \varphi_1(x_p) & \dots & \varphi_k(x_p) \end{pmatrix}.$$

This $p \times (k + 1)$ matrix has a non-trivial (b_0, \dots, b_k) in its kernel since $p \leq k$.

¹See [6] pp. 89–92, [10] p. 195, [11]. Jensen [8] noticed the similarity of de Gua’s and Laguerre’s arguments. Instead of derivation, de Gua used Johan Hudde’s operation, which is described as a term by term multiplication of the polynomial and an arithmetic progression. The modern reader will simply compare this with the operation $Y \mapsto x^{-m+1}(x^m Y)'$.

²Authors of the 18th century used to count the roots with their multiplicity and to omit the parity conclusion. Incidentally, many authors following Cajori [2] attribute the parity conclusion to Gauss [4] in 1828. This is very strange as firstly, this conclusion is obvious, secondly, Fourier published it in 1820 (see [3] p. 294) and thirdly, it is absent from Gauss’ paper.

If $p = k$, the “only if” part shows that the coefficients b_i are non-zero, that their signs alternate and that there is no other positive root. This makes an (a_0, \dots, a_n) such that Y has the prescribed set of positive roots.

If $1 \leq p < k$, observe that the $p \times p$ submatrix formed by the first p columns of Φ is invertible. If its determinant were zero we could find a non-trivial column vector (c_0, \dots, c_{p-1}) in its kernel. But $\sum_0^{p-1} c_i \varphi_i(x)$ would have a root at each x_j . It would have p roots and at most $p - 1$ variations, which contradicts what we proved in the “only if” part.

So for any given (b_p, \dots, b_k) there is a unique (b_0, \dots, b_{p-1}) , computed by Cramer's rule, such that (b_0, \dots, b_k) is in the kernel of Φ . We introduce a real parameter ϵ and take $b_i(\epsilon) = (-1)^{k-i} \sigma_n \epsilon$, $p \leq i < k$ and $b_k(\epsilon) = \sigma_n$. The unique column vector $(b_0(\epsilon), \dots, b_{p-1}(\epsilon), b_p(\epsilon), \dots, b_k(\epsilon))$ in the kernel of Φ is such that each of its $k + 1$ components is a polynomial in ϵ , of first degree.

Set $Y_\epsilon(x) = b_0(\epsilon)\varphi_0(x) + \dots + b_k(\epsilon)\varphi_k(x)$. We have $Y_\epsilon(x_j) = 0$ for $j = 1, \dots, p$. Since $Y_0(x) = b_0(0)\varphi_0(x) + \dots + b_{p-1}(0)\varphi_{p-1}(x) + b_k(0)\varphi_k(x)$, the $p + 1$ coefficients in this expression are non-zero and have alternating signs, according to the “only if” part, which at the same time establishes that Y_0 cannot have any other positive root than the x_j 's. As $k - p$ is even and b_k has the prescribed sign, the $b_i(0)$'s, $i < p$, have the prescribed signs, and the same is true of $b_i(\epsilon)$ for any small enough ϵ . Now, the $b_i(\epsilon)$'s, $p \leq i < k$, have the prescribed signs for any positive ϵ . Thus Y_ϵ has the prescribed sign sequence for any sufficiently small positive ϵ .

Consider Y_ϵ/Y_0 . This quotient is analytic in $x \in]0, +\infty[$, as shown by expanding numerator and denominator in Taylor series around $x = x_j$ and by simplifying the factors $(x - x_j)$. As $\epsilon \rightarrow 0$, $Y_\epsilon/Y_0 \rightarrow 1$ for any $x \in]0, +\infty[$. This convergence is indeed uniform, the behavior of the fraction at zero and at infinity being controlled by the leading terms of the numerator and the denominator. Thus Y_ϵ has no other positive root for a sufficiently small ϵ . The problem is solved in the case of simple roots.

In the cases with multiple roots the construction and the proofs are exactly the same, except that Φ has, e.g., the row $(\varphi'_0(x_1), \dots, \varphi'_k(x_1))$ if x_1 is a multiple root, the row $(\varphi''_0(x_1), \dots, \varphi''_k(x_1))$ if it is at least a triple root, etc.

If finally $p = 0$, k is even and we can build a Y without positive root. We simply choose a sufficiently small positive ϵ and take $b_0(\epsilon) = \sigma_0$, $b_k(\epsilon) = \sigma_0$ and $b_i(\epsilon) = (-1)^i \epsilon \sigma_0$. \square

4 Further information about trinomials

Etymologically a trinomial is simply the sum of three monomials, and there is no restriction on the degree. Trinomials with unprescribed degree were studied early in the history of algebraic equations (see [14], pp. 11 and 24). They are natural objects in the context of Laguerre's extension of Descartes' rule to real exponents, as well as in the context of Khovanskky's theory of fewnomials [9], and are consequently the object of recent studies. In 2002, Haas [7], Li, Rojas & Wang [12] proved that the optimal upper bound on the number of roots in the positive quadrant of a system of two trinomials in two variables is five.

We were not able to find the following elegant and elementary formula in any of these old or recent studies.

Proposition 2. A trinomial $ax^\alpha + bx^\beta + cx^\gamma$, $\alpha < \beta < \gamma$, $a > 0$, $c > 0$, $b < 0$, is positive on $]0, \infty[$ if and only if

$$\left(\frac{a}{\gamma - \beta}\right)^{\gamma - \beta} \left(\frac{b}{\alpha - \gamma}\right)^{\alpha - \gamma} \left(\frac{c}{\beta - \alpha}\right)^{\beta - \alpha} > 1.$$

Remark. The discriminant of $ax^\alpha + bx^\beta + cx^\gamma$ is $b^2 - 4ac$ if $(\alpha, \beta, \gamma) = (0, 1, 2)$. It is $-c(4b^3 + 27ca^2)$ if $(\alpha, \beta, \gamma) = (0, 1, 3)$. It is $c^2(-27b^4 + 256ca^3)$ if $(\alpha, \beta, \gamma) = (0, 1, 4)$ and $16ac(4ca - b^2)^2$ if $(\alpha, \beta, \gamma) = (0, 2, 4)$. These formulas are quite familiar. The general formula in the proposition gives in each case the main factor.

Proof. The formula may be obtained by a direct computation, where there occur unexpected simplifications. We will present a short matrix algebra argument, basically the same as in the previous section. Consider the trinomial $Y = ax^\alpha + bx^\beta + cx^\gamma$. We determine (a, b, c) such that Y has a double root at x_1 . We find that (a, b, c) should be in the kernel of

$$\Phi = \begin{pmatrix} x_1^\alpha & x_1^\beta & x_1^\gamma \\ \alpha x_1^{\alpha-1} & \beta x_1^{\beta-1} & \gamma x_1^{\gamma-1} \end{pmatrix},$$

which means

$$\frac{ax_1^\alpha}{\gamma - \beta} = \frac{bx_1^\beta}{\alpha - \gamma} = \frac{cx_1^\gamma}{\beta - \alpha}.$$

We set

$$A = \frac{a}{\gamma - \beta}, \quad B = \frac{b}{\alpha - \gamma}, \quad C = \frac{c}{\beta - \alpha},$$

which are positive numbers according to the hypotheses of the proposition, and continue the computation by eliminating x_1 , which gives the expected condition

$$A^{\gamma - \beta} B^{\alpha - \gamma} C^{\beta - \alpha} = 1$$

for a double root. Now a mere study of $A^{\gamma - \beta} B^{\alpha - \gamma} C^{\beta - \alpha}$ as a function of b gives the proposition. \square

For completeness, note that under the condition $A^{\gamma - \beta} B^{\alpha - \gamma} C^{\beta - \alpha} = 1$ the double root is located at

$$x_1 = \left(\frac{C}{B}\right)^{1/(\beta - \gamma)} = \left(\frac{A}{C}\right)^{1/(\gamma - \alpha)} = \left(\frac{B}{A}\right)^{1/(\alpha - \beta)}.$$

5 Combining positive and negative roots

If we just focus on the positive roots of a given polynomial Y , Proposition 1 tells us that there is nothing we can add to Descartes' rule of signs. All that can be deduced from the sign sequence of Y is the upper bound and the parity of P , the cardinality of the multiset of positive roots. By changing x to $-x$, the same can be said on N , the cardinality of the multiset of negative roots of Y . But let us consider the constraints on (P, N) .

Consider a polynomial $Y(x)$ of degree 4 with sign sequence $+, -, -, -, +$. By Descartes' rule $P = 0$ or 2 . If we change x to $-x$ the sign sequence becomes $+, +, -, +, +$. Thus $N = 0$ or 2 . Grabiner [5] points out that $(P, N) = (0, 2)$ is *impossible* for such a $Y(x)$.

The proof does not require any computation. Because the constant term $Y(0)$ is positive, $P = 0$ implies $Y > 0$ for $x > 0$. But the odd part of Y is negative when $x > 0$. So the even part is positive. For $x < 0$ the odd and the even parts are then positive. We must have $N = 0$. \square

Consider an expression (1) as in Proposition 1, where the α_i 's are non-negative integers, i.e., $Y(x)$ is a polynomial. Given the sequence of signs $\sigma_0, \dots, \sigma_n$, Grabiner proves³ that all the (P, N) 's produced as follows are *possible*. One chooses any subset of $\{1, \dots, n-1\}$ and "erases" the corresponding σ_i 's from the sequence of signs. The number of variations in the resulting sequence gives P . One then considers the modified sequence of signs $\sigma_0(-1)^{\alpha_0}, \sigma_1(-1)^{\alpha_1}, \dots, \sigma_n(-1)^{\alpha_n}$ and erases from it the signs with index in the same subset. The new number of variations gives N .

Let us call this construction of possible (P, N) 's Grabiner's erasing term rule. Grabiner observes that it gives all the possible (P, N) 's for polynomials up to degree 4, and conjectures that the same is true for higher degrees.

Here is a counterexample. Consider a polynomial of 5th degree with sign sequence $+, +, -, +, +, -$. Here $(P, N) = (3, 0)$ is possible, as shown by

$$\begin{aligned} & (20 + 37x + 18x^2)(1 - x)(2 - x)(3 - x) \\ &= 120 + 2x - 179x^2 + 4x^3 + 71x^4 - 18x^5. \end{aligned}$$

This combination is not obtained by Grabiner's erasing term rule. Indeed, after changing x in $-x$ the sequence becomes $+, -, -, -, +, +$. As $N = 0$ we should erase terms and obtain a sequence without variation. We should erase either the three $-$, or the three $-$ and the internal $+$. But erasing the corresponding signs in $+, +, -, +, +, -$ gives in both cases only one variation, while $P = 3$.

In contrast $(P, N) = (3, 0)$ is impossible for the sequence $+, +, -, +, -, -$. Descartes' rule predicts all the impossibilities for degree 5 polynomials without gaps, except this one and its trivial analogues.

To prove this impossibility, let us specify the ordering of sign sequences by associating the first sign to the constant term. We notice that the odd part \mathcal{O} has sequence of signs $+, +, -$. Its derivative of course has the same sequence and consequently a unique positive root r . On $]0, r[$, \mathcal{O} is positive, and for $x > r$, \mathcal{O} is decreasing. The even part \mathcal{E} has sequence of signs $+, -, -$, thus it decreases for $x > 0$. Our polynomial $\mathcal{E} + \mathcal{O}$ decreases and has at most 1 root on $]r, +\infty[$. As $N = 0$, $\mathcal{E} - \mathcal{O}$ is positive on $]0, +\infty[$. On $]0, r[$, \mathcal{O} and thus $\mathcal{E} + \mathcal{O}$ are positive. There is one positive root on $]r, +\infty[$, thus $P = 3$ is impossible. \square

Let us pass to degree 6 polynomials.

³The idea is that small terms can be neglected before applying Descartes' rule. Estimates on how small these terms should be are discussed in [13]. It is interesting to compare these estimates to the first lemma in [6].

The exhaustive list of non-Descartes impossibilities is, up to trivial transformations:

$$\begin{aligned} +, +, -, +, -, +, + & \text{ is incompatible with } (P, N) = (2, 0) \text{ or } (4, 0), \\ +, +, +, +, -, +, + & \text{ is incompatible with } (P, N) = (2, 0), \\ +, +, -, -, -, -, + & \text{ is incompatible with } (P, N) = (0, 4). \end{aligned}$$

The argument we gave in Grabiner's example proves the first and second statements.

To prove the third statement, we write a polynomial with $(P, N) = (0, 4)$ as

$$p = (c - bx + x^2)(x + x_1)(x + x_2)(x + x_3)(x + x_4),$$

where $x_i > 0, i = 1, \dots, 4$,

$$4c > b^2 > 0. \quad (2)$$

Expanding, $p = a_0 + a_1x + \dots + a_5x^5 + x^6$, with in particular

$$a_2 = c\beta - b\gamma + \delta < 0, \quad (3)$$

$$a_5 = -b + \alpha < 0, \quad (4)$$

the positive numbers $\alpha, \beta, \gamma, \delta$ being defined as

$$\begin{aligned} \alpha &= x_1 + x_2 + x_3 + x_4, \\ \beta &= x_1x_2 + x_1x_3 + x_2x_3 + x_1x_4 + x_2x_4 + x_3x_4, \\ \gamma &= x_1x_2x_3 + x_1x_2x_4 + x_1x_3x_4 + x_2x_3x_4, \\ \delta &= x_1x_2x_3x_4. \end{aligned} \quad (5)$$

By (2) and (4),

$$\sqrt{c} > b/2 > \alpha/2 > 0, \quad (6)$$

and, by (2) and (3),

$$2\sqrt{c} > b > \frac{c\beta + \delta}{\gamma},$$

which gives

$$\beta\sqrt{c}^2 - 2\gamma\sqrt{c} + \delta < 0. \quad (7)$$

We expand the discriminant $4\gamma^2 - 4\beta\delta$ of this expression using (5). All the terms are positive. Now (7) implies in particular that

$$\sqrt{c} < \frac{\gamma + \sqrt{\gamma^2 - \beta\delta}}{\beta}.$$

Combining with (6) gives $\beta\alpha/2 < \beta\sqrt{c} < \gamma + \sqrt{\gamma^2 - \beta\delta}$ or

$$\alpha\beta/2 - \gamma < \sqrt{\gamma^2 - \beta\delta}. \quad (8)$$

After expanding and canceling, we see that all the terms of $\alpha\beta/2 - \gamma$ are positive. We square both sides of (8). This gives us $\alpha^2\beta - 4\alpha\gamma + 4\delta < 0$. But the left-hand side, after expanding and canceling, has only positive terms. This is a contradiction. \square

To check that our list of impossibilities is complete at degree 6, it is enough to consider the possibilities established by Grabiner's erasing term rule and the two following polynomials, which after changing x into $-x$, x into $1/x$ or Y into $-Y$ provide examples for all the other possibilities:

$$\begin{aligned} & (31 + 11x + x^2)(1 - x)(2 - x)(3 - x)(4 - x) \\ &= 744 - 1286x + 559x^2 + 25x^3 - 44x^4 + x^5 + x^6, \\ & (9 + 8x + 2x^2)(1 - x)(2 - x)(3 - x)(4 - x) \\ &= 216 - 258x - 37x^2 + 90x^3 - x^4 - 12x^5 + 2x^6. \end{aligned}$$

These possibilities and impossibilities do not seem to organize themselves in simple classes.

Acknowledgement

We wish to thank Alain Chenciner, David Grabiner, Dieter Schmidt and the referee for their precious remarks. This work is supported by the Chinese Academy of Sciences visiting professorship for senior international scientists grant no. 2009J2-11, and NSFC grant no. 10833001.

References

- [1] Bruce Anderson, Jeffrey Jackson, Meera Sitharam, *Descartes' rule of signs revisited*, The American Mathematical Monthly 105 (1998) pp. 447–451
- [2] Florian Cajori, *A history of the arithmetical methods of approximation to the roots of numerical equations of one unknown quantity*, Colorado College Publication, Science series 12–7 (1910) pp. 171–215
- [3] Joseph Fourier, *Sur l'usage du théorème de Descartes dans la recherche des limites des racines*, Bulletin des sciences par la Société philomatique de Paris (1820) pp. 156–165, 181–187; *œuvres* 2, pp. 291–309, Gauthier-Villars, 1890
- [4] Carl Friedrich Gauss, *Beweis eines algebraischen Lehrsatzes*, Journal für die reine und angewandte Mathematik 3 (1828) pp. 1–4; *Werke* 3, pp. 67–70, Göttingen, 1866
- [5] David J. Grabiner, *Descartes' Rule of Signs: Another Construction*, The American Mathematical Monthly 106 (1999) pp. 854–856
- [6] Jean-Paul de Gua de Malves, *Démonstrations de la Règle de Descartes, Pour connaître le nombre des Racines positives & négatives dans les Équations qui n'ont point de Racines imaginaires*, Mémoires de Mathématique et de Physique tirés des registres de l'Académie Royale des Sciences (1741) pp. 72–96
- [7] Bertrand Haas, *A Simple Counterexample to Kouchnirenko's Conjecture*, Beiträge zur Algebra und Geometrie, Contributions to Algebra and Geometry 43 (2002) pp. 1–8
- [8] J.L.W.V. Jensen, *Recherches sur la théorie des équations*, Acta Mathematica 36 (1913) pp. 181–195
- [9] A.G. Khovanskii, *Fewnomials*, Translations of mathematical monographs 88, American Math. Society, 1991
- [10] Joseph Louis Lagrange, *Traité de la résolution des équations numériques de tous les degrés, avec des Notes sur plusieurs points de la Théorie des équations algébriques*, Courcier, Paris 1798 and 1808; *œuvres* 8, Gauthier-Villars, 1879
- [11] Edmond Laguerre, *Sur la théorie des équations numériques*, Journal de Mathématiques pures et appliquées, s. 3, t. 9 (1883) pp. 99–146; *œuvres* 1, Paris, 1898, Chelsea, New York, 1972, pp. 3–47

- [12] Tien-Yien Li, J. Maurice Rojas, Xiaoshen Wang, *Counting real connected components of trinomial curve intersections and m -nomial hypersurfaces*, Discrete Comput. Geom. 30 (2003) pp. 379–414
- [13] Christian Marchal, *Une amélioration commode du critère de Descartes*, Sciences, AFAS, fasc. 4 (2006) pp. 3–6
- [14] Jacqueline Stedall, *From Cardano's great art to Lagrange's reflections: filling a gap in the history of algebra*, European Mathematical Society, 2011

Alain Albouy
Purple Mountain Observatory
Chinese Academy of Sciences
2 West Beijing road
Nanjing 210008, China

and

CNRS-UMR8028, IMCCE
Observatoire de Paris
77 avenue Denfert-Rochereau
F-75014 Paris, France
albouy@imcce.fr

Yanning Fu
Purple Mountain Observatory
Chinese Academy of Sciences
2 West Beijing road
Nanjing 210008, China
fyn@pmo.ac.cn