

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 69 (2014)

**Artikel:** An explicit family of Um-numbers  
**Autor:** Chaves, Ana Paula / Marques, Diego  
**DOI:** <https://doi.org/10.5169/seals-515853>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

---

## An explicit family of $U_m$ -numbers

---

Ana Paula Chaves and Diego Marques

Diego Marques was awarded a PhD degree at the Universidade de Brasília (2009) in less than four months. Currently, he is a professor in the department of mathematics at the same university. His field of interest is elementary and transcendental number theory.

Ana Paula Chaves received her B.S. and M.S. degrees in mathematics from the Universidade Federal do Ceará, and her PhD from Universidade de Brasília in 2013. Currently, she is a professor of mathematics and statistics at the Universidade Federal de Goiás. Her interests are transcendental number theory and diophantine equations.

### 1 Introduction

Transcendental number theory began in 1844 with Liouville's proof [7] that if an algebraic number  $\alpha$  has degree  $n > 1$ , then there exists a constant  $C > 0$  such that  $|\alpha - p/q| > Cq^{-n}$ , for all  $p/q \in \mathbb{Q} \setminus \{0\}$ . Using this result, Liouville gave the first explicit examples of transcendental numbers, the so-called Liouville numbers: a real number  $\xi$  is called a *Liouville number*, if for any positive real number  $\omega$  there exist infinitely many rational numbers  $p/q$ , with  $q \geq 1$ , such that

$$0 < \left| \xi - \frac{p}{q} \right| < \frac{1}{q^\omega}.$$

Im Jahre 1844 gab Joseph Liouville ein erstes Beispiel einer transzendenten Zahl, nämlich die Liouville-Zahl  $\ell = \sum_{n \geq 1} 10^{-n!}$ . Fast ein Jahrhundert später schlug Kurt Mahler vor, die reellen Zahlen in vier Kategorien einzuteilen, je nachdem wie gut sie sich bei der Approximation durch algebraische Zahlen verhalten. Insbesondere zerfielen dabei die transzendenten Zahlen in drei Klassen, nämlich die  $S$ -,  $T$ - und  $U$ -Zahlen. 1952 bemerkte LeVeque, dass die  $U$ -Zahlen sich noch weiter, bezüglich ihrer Approximierbarkeit durch algebraische Zahlen vom Grad  $m$ , in die unendlich vielen disjunkten Klassen der  $U_m$ -Zahlen unterteilen lassen. LeVeque zeigte insbesondere, dass  $\sqrt[m]{(3 + \ell)/2}$  eine  $U_m$ -Zahl ist. In der vorliegenden Arbeit konstruieren die Autoren  $U_m$ -Zahlen für alle  $m$  auf besonders transparente Weise, nämlich als Produkt von  $\ell$  mit gewissen algebraischen Zahlen vom Grad  $m$ .

A classical example of a Liouville number is the *Liouville constant*  $\ell$ , defined as a decimal with a 1 in each decimal place corresponding to  $n!$  and 0 otherwise. It can be represented by the fast convergent series  $\ell = \sum_{n=1}^{\infty} 10^{-n!} = 0.1100010\dots$

In 1962, Erdős [4] proved that every nonzero real number can be written as the sum and the product of two Liouville numbers. Since the set of the Liouville numbers has null Lebesgue measure, one may interpret this as saying that in spite of being an “invisible” set, the Liouville numbers are strategically disposed along the real line.

There exist several classifications of the transcendental numbers in the literature. One attempt towards a classification was made in 1932 by Mahler [8], who proposed to subdivide the set of real numbers into four classes (one of them being the class of algebraic numbers) according to their properties of approximation by algebraic numbers. For instance, he split the set of transcendental numbers into three disjoint sets named *S*-, *T*- and *U-numbers*. Particularly, the *U*-numbers generalize the concept of Liouville numbers.

We denote by  $\omega_n^*(\xi)$  as the supremum of the real numbers  $\omega^*$  for which there exist infinitely many real algebraic numbers  $\alpha$  of degree  $n$  satisfying

$$0 < |\xi - \alpha| < \mathcal{H}(\alpha)^{-\omega^*-1},$$

where  $\mathcal{H}(\alpha)$  (so-called the *height* of  $\alpha$ ) is the maximum of absolute values of coefficients of the minimal polynomial<sup>1</sup> of  $\alpha$ . The number  $\xi$  is said to be a  *$U_m^*$ -number* (according to LeVeque [6]) if  $\omega_m^*(\xi) = \infty$  and  $\omega_n^*(\xi) < \infty$  for  $1 \leq n < m$  ( $m$  is called the *type* of the *U*-number). We point out that we actually have defined a Koksma  *$U_m^*$ -number* instead of a Mahler  *$U_m$ -number*. However, it is well known that they are the same [3, cf. Theorem 3.6] and [1]. We remark that the set of  *$U_1$ -numbers* is precisely the set of Liouville numbers.

The existence of  *$U_m$ -numbers* for all  $m \geq 1$ , was first proved by LeVeque [6]. Indeed, he was able to exhibit such examples as the  $m$ th root of some convenient Liouville numbers, e.g.,  $\sqrt[m]{(3 + \ell)/4}$  is a  *$U_m$ -number*, for all  $m \geq 1$ .

In this note, we use the Gütting method [5] to prove that we can find explicit  *$U_m$ -numbers* in a more natural way: the product of certain  $m$ -degree algebraic numbers by  $\ell$ . Moreover, we obtain an upper bound for  $\omega_n^*$ . More precisely, our result is the following

**Theorem 1.** *Let  $\alpha$  be an algebraic number of degree  $m$ . Suppose that the minimal polynomial  $P$  of  $\alpha$  has a leading coefficient of the form  $2^a \cdot 5^b > 1$ , and  $p \nmid P(0)$ , for  $p = 2, 5$ , and let  $\ell$  be the Liouville constant. Then  $\alpha\ell$  is a  *$U_m$ -number*, with*

$$\omega_n^*(\alpha\ell) \leq 2m^2n + m - 1, \text{ for } n = 1, \dots, m - 1. \quad (1)$$

For example,  $\sqrt[m]{3/2} \cdot \ell$  is a  *$U_m$ -number* for all  $m \geq 1$ .

---

<sup>1</sup>Throughout the paper, a polynomial is said to be minimal if it is a primitive minimal polynomial over  $\mathbb{Z}$ .

## 2 Auxiliary Results

Before starting the proof of the Theorem, two technical results are needed.

**Lemma 1.** *Given  $P(x) \in \mathbb{Z}[x]$  with degree  $m$  and  $a/b \in \mathbb{Q} \setminus \{0\}$ . If  $Q(x) = a^m P(bx/a)$ , then*

$$\mathcal{H}(Q) \leq \max\{|a|, |b|\}^m \mathcal{H}(P),$$

where, as usual,  $\mathcal{H}(P)$  denotes the maximum of absolute values of coefficients of  $P$  (the so-called height of  $P$ ).

*Proof.* If  $P(x) = \sum_{j=0}^m a_j x^j$ , then  $Q(x) = \sum_{j=0}^m a_j b^j a^{m-j} x^j$ . Supposing, without loss of generality, that  $|a| \geq |b|$ , we have  $|a|^m |a_j| \geq |a|^{m-j} |a_j| |b|^j$  for  $0 \leq j \leq m$ . Hence, we are done.  $\square$

In addition to Lemma 1, we use the fact that algebraic numbers are not well approximable by algebraic numbers.

**Lemma 2** (Cf. Corollary A.2 of [3]). *Let  $\alpha$  and  $\beta$  be two distinct nonzero algebraic numbers of degree  $n$  and  $m$ , respectively. Then we have*

$$\begin{aligned} |\alpha - \beta| &\geq (n+1)^{-m/2} (m+1)^{-n/2} \max \left\{ \frac{(n+1)^{-(m-1)/2}}{2^{-n}}, \frac{(m+1)^{-(n-1)/2}}{2^{-m}} \right\} \\ &\quad \times H(\alpha)^{-m} H(\beta)^{-n}. \end{aligned}$$

*Proof.* A sketch of the proof can be found in Appendix A of [3].  $\square$

## 3 Proof of the Theorem

For  $k \geq 1$ , set

$$p_k = 10^{k!} \sum_{j=1}^k 10^{-j!}, \quad q_k = 10^{k!} \quad \text{and} \quad \alpha_k = \frac{p_k}{q_k}.$$

We observe that  $\mathcal{H}(\alpha_{k-1}) < \mathcal{H}(\alpha_k) = 10^{k!} = \mathcal{H}(\alpha_{k-1})^k$  and

$$|\ell - \alpha_k| < \frac{10}{9} \mathcal{H}(\alpha_k)^{-k-1}. \tag{2}$$

Thus, setting  $\gamma_k = \alpha \alpha_k$ , we obtain of (2)

$$|\alpha \ell - \gamma_k| \leq c \mathcal{H}(\alpha_k)^{-k-1}, \tag{3}$$

where  $c = 10|\alpha|/9$ . It follows by Lemma 1 that  $\mathcal{H}(\alpha_k)^m \geq \mathcal{H}(\alpha)^{-1} \mathcal{H}(\gamma_k)$  and thus we conclude that

$$|\alpha \ell - \gamma_k| \leq c \mathcal{H}(\alpha)^{(k+1)/m} \mathcal{H}(\gamma_k)^{-(k+1)/m}. \tag{4}$$

Consequently,  $\alpha \ell$  is a  $U$ -number with type at most  $m$  (since  $\gamma_k$  has degree  $m$ ).

We claim that  $H(\alpha_k) \leq H(\gamma_k)$ , for all  $k \geq 1$ . In fact, let  $P(x) = \sum_{j=0}^m a_j x^j$  be the minimal polynomial of  $\alpha$ . In particular,  $P(\alpha) = 0$  and a simple calculation gives  $Q(\gamma_k) = 0$ , where  $Q(x) = \sum_{j=0}^m a_j p_k^{m-j} q_k^j x^j \in \mathbb{Z}[x]$ . Note that  $\deg Q = m$  and  $\gamma_k$  is an  $m$ -degree algebraic number. Thus, in order to prove that  $Q$  is the minimal polynomial of  $\gamma_k$ , we need to prove that  $Q$  is primitive. In other words, we must prove that

$$\gcd(a_0 p_k^m, a_1 p_k^{m-1} q_k, \dots, a_m q_k^m) = 1.$$

This follows immediately from the facts that  $\gcd(a_0, \dots, a_m) = 1$  and the hypotheses on  $a_0$  and  $a_m$  (yielding  $\gcd(a_0, q_k) = \gcd(a_m, p_k) = 1$ ), we leave the details to the reader. Thus, in particular, we have that

$$H(\gamma_k) \geq \max\{|a_0||p_k|^n, |a_n||q_k|^n\} \geq \max\{|p_k|, |q_k|\} = H(\alpha_k)$$

as desired.

Now we use this together with Lemma 1 to obtain

$$\mathcal{H}(\gamma_{k+1}) \leq \mathcal{H}(\alpha) \mathcal{H}(\alpha_{k+1})^m = \mathcal{H}(\alpha) \mathcal{H}(\alpha_k)^{(k+1)m} \leq \mathcal{H}(\alpha) \mathcal{H}(\gamma_k)^{(k+1)m}. \quad (5)$$

Now, let  $\gamma$  be an  $n$ -degree real algebraic number, with  $n < m$  and  $\mathcal{H}(\gamma) \geq \mathcal{H}(\gamma_1)$ . Thus, there exists a sufficient large  $k$  such that

$$\mathcal{H}(\gamma_k) < \mathcal{H}(\gamma)^{2m^2} < \mathcal{H}(\gamma_{k+1}) \leq \mathcal{H}(\alpha) \mathcal{H}(\gamma_k)^{(k+1)m}. \quad (6)$$

On the other hand, Lemma 2 yields

$$|\gamma_k - \gamma| \geq f(m, n) \mathcal{H}(\gamma)^{-m} \mathcal{H}(\gamma_k)^{-n}, \quad (7)$$

where  $f(m, n)$  is a positive number which does not depend on  $k$  and  $\gamma$  (see Lemma 2). Therefore by the chain of inequalities in (6)

$$|\gamma_k - \gamma| \geq f(m, n) \mathcal{H}(\alpha)^{-1/2m} \mathcal{H}(\gamma_k)^{-(k+1)/2-n}. \quad (8)$$

By taking  $\mathcal{H}(\gamma)$  large enough, the index  $k$  satisfies

$$\mathcal{H}(\gamma_k)^{(k+1)/2-n} \geq 2cf(m, n)^{-1} \mathcal{H}(\alpha)^{k+1/2m}. \quad (9)$$

Thus (4), (8) and (9) yield that  $|\gamma_k - \gamma| \geq 2|\alpha\ell - \gamma_k|$ . Therefore, for all  $n$ -degree algebraic numbers with a sufficiently large weight, we have

$$\begin{aligned} |\alpha\ell - \gamma| &\geq |\gamma_k - \gamma| - |\alpha\ell - \gamma_k| \geq \frac{1}{2} |\gamma_k - \gamma| \\ &\geq \frac{f(m, n)}{2} \mathcal{H}(\gamma)^{-m} \mathcal{H}(\gamma_k)^{-n} > \frac{f(m, n)}{2} \mathcal{H}(\gamma)^{-2m^2n-m}, \end{aligned}$$

where we used the left-hand side of (6). In conclusion,  $\alpha\ell$  is a  $U_m$ -number with  $\omega_n^*(\alpha\ell) \leq 2m^2n + m - 1$ . This finishes the proof.  $\square$

We finish by pointing out that Alniaçik et al. [2] showed the existence of  $U_m$ -numbers  $\xi$  with sharper upper bounds for  $\omega_n^*(\xi)$ , where  $n = 1, \dots, m-1$ . However, in their method  $\xi$  is constructed as the limit of a rapidly converging sequence of  $m$ -degree algebraic numbers and therefore could not be made explicit.

### Acknowledgement

The authors are grateful to CAPES, FEMAT, DPP-UnB and CNPq for the financial support. They also thank Yann Bugeaud for nice discussions on the subject. Part of this work was made during a postdoctoral position of the second author in the Department of Mathematics at the University of British Columbia. He thanks the UBC for excellent working conditions.

### References

- [1] Alnacı̄k, K.: On the subclasses  $U_m$  in Mahler's classification of the transcendental numbers, *Istanbul Univ. Fen Fak. Mecm. Ser.* **44**, 39–82 (1972).
- [2] Alnacı̄k, K., Avcı, Y., Bugeaud, Y.: On  $U_m$ -numbers with small transcendence measures. *Acta. Math. Hungar.* **99**, 271–277 (2003).
- [3] Bugeaud, Y.: *Approximation by Algebraic Numbers*, Cambridge Tracts in Mathematics Vol. **160**, Cambridge University Press, New York (2004).
- [4] Erdős, P.: Representations of real numbers as sums and products of Liouville numbers, *Michigan Math. J.* **9**, 59–60 (1962).
- [5] Gütting, R.: Zur Berechnung der Mahlerschen Funktionen  $w_n$ , *J. Reine Angew. Math.* **232**, 122–135 (1968).
- [6] LeVeque, W.J.: On Mahler's  $U$ -Numbers, *J. of the London Math. Soc.* **28**, 220–229 (1953).
- [7] Liouville, J.: Sur des classes très-étendue de quantités dont la valeur n'est ni algébrique, ni même réductible à des irrationnelles algébriques, *C. R.* **18**, 883–885 (1844).
- [8] Mahler, K.: Zur Approximation der Exponentialfunktion und des Logarithmus I, *J. Reine Angew. Math.* **166**, 118–136 (1932).

Ana Paula Chaves  
 Instituto de Matemática e Estatística  
 Universidade Federal de Goiás  
 Goiânia, Go, Brazil  
 e-mail: apchaves@ufg.br

Diego Marques  
 Departamento de Matemática  
 Universidade de Brasília  
 Brasília, DF, Brazil  
 e-mail: diego@mat.unb.br