

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 69 (2014)

Artikel: Limit shape of iterated Kiepert triangles
Autor: Nicollier, Grégoire / Stadler, Albert
DOI: <https://doi.org/10.5169/seals-515859>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 12.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Limit shape of iterated Kiepert triangles

Grégoire Nicollier and Albert Stadler

Grégoire Nicollier is a mountain guide and teaches mathematics in French and in German to engineering students at the University of Applied Sciences of Western Switzerland in Sion (Valais). He studied mathematics (together with Albert Stadler) at ETH, obtained a PhD in homological algebra in 1984 under Professor Stambach, and remained at ETH till 1991 as a lecturer.

Albert Stadler obtained his PhD in mathematics from ETH Zürich in 1986, under the supervision of Professor Chandrasekharan. Since 1988 he works for a global financial services firm. His main interests focus on analytic number theory, in particular on the Riemann zeta function. He is an active contributor to various problem solving forums.

We consider the following discrete dynamical system: start with a base triangle $\Delta_0 = ABC$ having at least two different vertices; erect externally on all sides of Δ_0 similar isosceles triangles with apex angle $\pi - 2\theta_0$; get the *Kiepert triangle* $\Delta_1 = A_1B_1C_1$ by taking the apices opposite to A , B , and C , respectively. Note that $\theta_0 = 0$ gives the medial triangle Δ_1 , which is directly similar to Δ_0 . We iterate the process with Δ_1 and $\pi - 2\theta_1$, Δ_2 and $\pi - 2\theta_2$, and so on. It is shown in [2] that this sequence of triangles has an equilateral limit shape when the apex angles are all equal and nonstraight. We give here a very short proof of this result and we determine the convergence behavior for any other choice of the apex angles.

Aus einem Dreieck kann ein neues Dreieck auf viele Arten entstehen. Ein Beispiel ist das Napoleon-Dreieck: Gleichschenklige Dreiecke mit einem Scheitelwinkel von 120° , die über den Seiten eines Dreiecks errichtet werden, ergeben immer ein gleichseitiges Dreieck. Oder die aufeinanderfolgenden Höhenfußpunktdreiecke eines Dreiecks: ihre Formenfolge kann beinahe jedes Verhalten aufweisen. Erwähnenswert ist auch die fraktale Struktur der Folge der Spiegelungsdreiecke, wo die Eckpunkte jeweils an der gegenüberliegenden Seite gespiegelt werden (G. Nicollier, *Iterated Reflection Triangles*, Forum Geometricorum 12 (2012), 83–129). Der Themenkreis der iterierten Dreiecke (und Polygone) ist sehr fruchtbar und bedient sich oft zyklischer Matrizen und der diskreten Fourier-Transformation. Die Autoren des vorliegenden Artikels verallgemeinern mit Hilfe einer Formfunktion einen Teil der Arbeit *Iterative Triangle Transformations* von D. Vartziotis und S. Huggenberger, die hier vor kurzem erschienen ist.

We use for any triangle $\Delta = UVW$ in the complex plane (with at least two different vertices) the shape function [1]

$$\sigma(\Delta) = \frac{U + V\zeta + W\zeta^2}{U + V\zeta^2 + W\zeta} \quad \text{with } \zeta = e^{2\pi i/3}.$$

One has $\sigma(\Delta) = \sigma(\Delta')$ if and only if Δ and Δ' are directly similar for the given ordered vertices, and $\sigma(\Delta) = 0$ or ∞ means that Δ is a positively or negatively oriented equilateral triangle, respectively. The image of σ is the extended complex plane: one has $\sigma(\Delta) = \zeta$ for the vertices $0, 0, 1$ in order, and $\sigma(\Delta) = s \neq \zeta$ for the vertices $0, 1, \frac{\zeta s - 1}{\zeta - s}$.

Theorem 1. *If the successive angles $\theta_0, \theta_1, \dots \in]0, \frac{\pi}{2}[$ are bounded away from 0 and from $\frac{\pi}{2}$, the above sequence of triangles converges to an equilateral limit shape.*

Proof. Suppose that $\Delta_0 = ABC$ is positively oriented. The vertices of Δ_1 are then

$$C_1 = \frac{A+B}{2} + i \frac{A-B}{2} \tan \theta_0,$$

and cyclically. A simple computation gives

$$\sigma(\Delta_1) = \frac{1 - \sqrt{3} \tan \theta_0}{1 + \sqrt{3} \tan \theta_0} \cdot \sigma(\Delta_0)$$

(note that $\theta_0 = 30^\circ$ proves Napoleon's theorem). Since Δ_1 is also positively oriented, one gets

$$\sigma(\Delta_2) = \frac{1 - \sqrt{3} \tan \theta_1}{1 + \sqrt{3} \tan \theta_1} \cdot \frac{1 - \sqrt{3} \tan \theta_0}{1 + \sqrt{3} \tan \theta_0} \cdot \sigma(\Delta_0),$$

and so on. Since each factor $\left| \frac{1 - \sqrt{3} \tan \theta}{1 + \sqrt{3} \tan \theta} \right|$ is smaller than some fixed bound $b < 1$ (Figure 1), one has $\lim_{n \rightarrow \infty} \sigma(\Delta_n) = 0$. \square

Theorem 2. *If one excepts the trivial case of an equilateral base triangle Δ_0 , the above sequence of iterated Kiepert triangles converges to an equilateral limit shape if and only if $\lim_{n \rightarrow \infty} \prod_{k=0}^n \frac{1 - \sqrt{3} \tan \theta_k}{1 + \sqrt{3} \tan \theta_k} = 0$, i.e., if and only if the successive angles $\theta_0 = \frac{\pi}{2}x_0, \theta_1 = \frac{\pi}{2}x_1, \dots \in [0, \frac{\pi}{2}[$ are such that $\theta_k = \frac{\pi}{6}$ for some k or $\sum_{k=0}^{\infty} \|x_k\| = \infty$, where $\|x\| = \min(x, 1-x)$ denotes the distance from $x \in [0, 1]$ to the nearest integer 0 or 1.*

If the sequence of iterated Kiepert triangles has no equilateral limit shape (and if Δ_0 is positively oriented), the sequence $(\sigma(\Delta_n))$ has the nonzero limit

$$\sigma(\Delta_0) \prod_{k=0}^{\infty} \frac{1 - \sqrt{3} \tan \theta_k}{1 + \sqrt{3} \tan \theta_k}$$

when $\lim_{k \rightarrow \infty} \theta_k = 0$, and has otherwise exactly two accumulation points given by

$$\pm \sigma(\Delta_0) \prod_{k=0}^{\infty} \left| \frac{1 - \sqrt{3} \tan \theta_k}{1 + \sqrt{3} \tan \theta_k} \right|.$$




Figure 1

Proof. After continuous extension at $x = 1$, the function

$$f(x) = \frac{1 - \sqrt{3} \tan(\frac{\pi}{2}x)}{1 + \sqrt{3} \tan(\frac{\pi}{2}x)} = \frac{2}{1 + \sqrt{3} \tan(\frac{\pi}{2}x)} - 1$$

falls from 1 to -1 on $[0, 1]$ with $f(\frac{2}{3}) = -\frac{1}{2}$ (Figure 1), and its derivative

$$f'(x) = \frac{-\sqrt{3} \pi}{4 \sin^2(\frac{\pi}{2}(x + \frac{1}{3}))}$$

grows on $[0, \frac{2}{3}]$ from $-\sqrt{3} \pi \approx -5.44$ at $x = 0$ to $-\frac{\sqrt{3}}{4} \pi$ at $x = \frac{2}{3}$, with $f'(\frac{1}{3}) = -\frac{\sqrt{3}}{3} \pi$, before falling to $\frac{\sqrt{3}}{3} \pi$ on $[\frac{2}{3}, 1]$. One has thus (Figure 1)

$$1 - 6\|x\| \leq |f(x)| \leq 1 - \|x\| \leq e^{-\|x\|}, \quad \forall x \in [0, 1],$$

and $\prod_{k=0}^{\infty} |f(x_k)| \leq e^{-\sum_{k=0}^{\infty} \|x_k\|}$: the infinite product is 0 or diverges to 0 when $x_k = \frac{1}{3}$ for some k or $\sum_{k=0}^{\infty} \|x_k\| = \infty$.

Conversely, suppose that all x_k are $\neq \frac{1}{3}$ and that $\sum_{k=0}^{\infty} \|x_k\| < \infty$: one has then $\|x_k\| < \frac{1}{10}$ as soon as k is large enough, say for all $k \geq K$, and one gets $\prod_{k=0}^{K-1} |f(x_k)| \neq 0$; further, since

$$1 - 6\|x\| \geq e^{-10\|x\|} \text{ for } \|x\| \leq \frac{1}{10},$$

one obtains $\prod_{k=K}^{\infty} |f(x_k)| \geq e^{-10 \sum_{k=K}^{\infty} \|x_k\|} > 0$. The full product $\prod_{k=0}^{\infty} |f(x_k)|$, whose factors lie in $]0, 1]$, has thus a limit $\lambda > 0$ (and any $\lambda \in]0, 1]$ can be obtained by an appropriate choice of the x_k 's): by taking Δ_0 nonequilateral and positively oriented, $\sigma(\Delta_n) = \sigma(\Delta_0) \prod_{k=0}^{n-1} f(x_k)$ has a limit $\pm \lambda \sigma(\Delta_0) \neq 0$ if all $f(x_k)$ are eventually

positive; otherwise, the infinite subsequences of the $\sigma(\Delta_n)$ with positive and negative $\prod_{k=0}^{n-1} f(x_k)$ have opposite limits $\lambda\sigma(\Delta_0)$ and $-\lambda\sigma(\Delta_0)$, respectively. Note that $\pm\lambda\sigma(\Delta_0)$ are, when $\neq -\sigma(\Delta_0)$, shapes of Kiepert triangles of Δ_0 . \square

References

- [1] Hajja, M.; Martini, H.; Spirova, M.: On converses of Napoleon's theorem and a modified shape function. *Beitr. Algebra Geom.* 47 (2006), 363–383.
- [2] Vartziotis, D.; Huggenberger, S.: Iterative geometric triangle transformations. *Elem. Math.* 67 (2012), 68–83.

Grégoire Nicollier
University of Applied Sciences of Western Switzerland
Route du Rawyl 47
CH-1950 Sion, Switzerland
e-mail: gregoire.nicollier@hevs.ch

Albert Stadler
Buchenrain 61
CH-8704 Herrliberg, Switzerland
e-mail: albert.stadler@ubs.com