

Zeitschrift:	Elemente der Mathematik
Herausgeber:	Schweizerische Mathematische Gesellschaft
Band:	71 (2016)
Heft:	3
Artikel:	Alternative proofs for Kocik's geometric diagram for relativistic velocity addition
Autor:	Sasane, Amol / Ufnarovski, Victor
DOI:	https://doi.org/10.5169/seals-630621

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Alternative proofs for Kocik's geometric diagram for relativistic velocity addition

Amol Sasane and Victor Ufnarovsky

Amol Sasane was trained as an Electrical Engineer at the Indian Institute of Technology, Mumbai. Subsequently, he received a PhD in Mathematics from the University of Groningen, the Netherlands, in 2001. He joined the London School of Economics in 2004, and is currently a Professor of Mathematics there. His research interests lie in Applicable Analysis.


Victor Ufnarovsky graduated from Moscow State University in 1977, and received his PhD in Mathematics from the Steklov Mathematical Institute in 1980. He joined the Institute of Mathematics and Computer Science in Moldova. Later he moved to Sweden, and currently he is a Professor of Mathematics at the Centre for Mathematical Sciences in Lund. His main interest lies in Computer Algebra and Algebra. For the last 15 years, he has been a deputy leader of the Swedish team at the International Mathematics Olympiad.

1 Introduction

Imagine a train moving at speed u with respect to the ground (as reckoned by someone sitting on the ground), and further that a person P is running with a speed v on the train (as reckoned by somebody sitting in the train). Before 1905, Newtonian physics dictated that the speed of the person P as observed by someone on the ground is $u + v$, while we


In einem Brief an Hendrik Lorentz vom Mai 1905 beschrieb Henri Poincaré die Formel für die relativistische Addition von Geschwindigkeiten in einer Dimension, nämlich $u \oplus v = \frac{u+v}{1+uv}$, wenn man für die Lichtgeschwindigkeit $c = 1$ setzt. Dieses Additionstheorem entspricht einer geometrischen Konstruktion, die Jerzy Kocik 2012 im American Journal of Physics veröffentlichte. Sein Originalbeweis verwendet kartesische Koordinaten. In der vorliegenden Arbeit geben die Autoren gleich drei kurze und elegante Beweise für die Korrespondenz der Poincaré-Formel und der Kocik-Konstruktion: Einer ist trigonometrischer Natur, ein zweiter ist in der euklidischen Geometrie angesiedelt, und der dritte verwendet Argumente der projektiven Geometrie. Alle drei Beweise werfen ein erhellendes Licht auf die relativistische Addition von Geschwindigkeiten.

now know better; the relativistic formula for velocity addition says that the speed should be $u \oplus v := (u + v)/(1 + uv)$, in units in which the speed of light is 1.




In [1], a geometric diagram for the construction of $u \oplus v$ from u and v was given. We recall it below.

Theorem 1.1 ([1]). *Draw a circle with center O and radius 1. Mark points U, V at distances u, v from O along the radius OC perpendicular to a diameter AB . Let the line joining B to V meet the circle at V' , and let the line joining A to U meet the circle at U' . Then $u \oplus v = OW$, where W the point of intersection of $U'V'$ with the radius OC .*



This construction allows visual justification of the following properties of \oplus . For all $u, v \in [0, 1]$, $u \oplus v \in [0, 1]$, $v \oplus 1 = 1$, $v \oplus 0 = v$, and when $0 \leq u, v \ll 1$, then $u \oplus v \approx u + v$. For example, let us justify this last fact geometrically. If $u, v \ll 1$, then $\angle OBV \approx 0$, and AV' is almost parallel to OV .



So ΔBOV is almost similar to $\Delta BAV'$, giving

$$AV' \approx \frac{AB}{OB} \cdot OV = \frac{2}{1} \cdot OV = 2v.$$


Since AV' is almost parallel to OC , $\Delta U'UW$ is almost similar to $\Delta U'AV'$. Moreover, as $u, v \ll 1$, $U'V' \approx AB = 2$, and $U'W \approx OB = 1$. Hence

$$UW \approx \frac{U'W}{U'V'} \cdot AV' \approx \frac{1}{2} \cdot 2v = v.$$

Thus if $w := OW$, then $w - u = UW \approx v$, that is, $w \approx u + v$.

In [1], Theorem 1.1 was proved using Cartesian coordinate geometry. In the next three sections, we give three alternative proofs of this result. (The more proofs, the merrier!)

2 A trigonometric proof



We refer to the picture above, calling

$$\angle BAU' = \angle OAU =: \alpha \quad \text{and} \quad \angle ABV' = \angle OBV =: \beta.$$

Let W be the point of intersection of $U'V'$ and OC , and set $OW =: w$. Then by looking at the right triangles ΔBOV and ΔAOU , we see that

$$\tan \beta = v \quad \text{and} \quad \tan \alpha = u.$$

Using the Sine Rule in $\Delta OWU'$, we have

$$\frac{1}{\sin \angle OWU'} = \frac{OU'}{\sin \angle OWU'} = \frac{OW}{\sin \angle OU'W} = \frac{w}{\sin \angle OU'W},$$

giving

$$w = \frac{\sin \angle OU'W}{\sin \angle OWU'}. \tag{1}$$

The proof will be completed by showing (below) that $\angle OU'W = \alpha + \beta$ and $\angle OWU' = 90^\circ + (\alpha - \beta)$, so that (1) yields

$$\begin{aligned} w &= \frac{\sin(\alpha + \beta)}{\sin(90^\circ + (\alpha - \beta))} = \frac{\sin(\alpha + \beta)}{\cos(\alpha - \beta)} = \frac{\sin \alpha \cos \beta + \cos \alpha \sin \beta}{\cos \alpha \cos \beta + \sin \alpha \sin \beta} \\ &= \frac{\tan \alpha + \tan \beta}{1 + \tan \alpha \tan \beta} = \frac{u + v}{1 + uv}, \end{aligned}$$

as desired.

First we will show $\angle OU'W = \alpha + \beta$. Note that $\Delta OAU'$ is isosceles with $OA = OU' = 1$ and so $\angle OU'U = \angle OAU = \alpha$. The chord AV' subtends equal angles at B and U' , and so $\angle UU'W = \angle ABV = \beta$. Hence

$$\angle OU'W = \angle OU'U + \angle UU'W = \alpha + \beta.$$

Next, let us show that $\angle OWU' = 90^\circ + (\alpha - \beta)$. To this end, note that $\angle OUU'$ is the common exterior angle for ΔOAU and $\Delta OU'U$, and using the fact that this equals the sum of the opposite interior angles in each triangle, we obtain

$$90^\circ + \alpha = \angle OUU' = \beta + \angle UWU',$$

so that $\angle OWU' = \angle UWU' = 90^\circ + (\alpha - \beta)$, completing the proof.

Yet another trigonometric proof can be obtained by focussing on $\Delta U'CW$, determining all its angles, and the side length $U'C$ (using the isosceles triangle $\Delta OU'C$), enabling the determination of WC ($= 1 - w$). The details are as follows. In the isosceles triangle $\Delta OU'C$, we have

$$\angle U'OC = 90^\circ - \angle BOU' = 90^\circ - 2\angle BAU' = 90^\circ - 2\alpha.$$

As $OU' = OC = 1$, we obtain $\angle OCU' = 45^\circ + \alpha$ and $U'C = 2\cos(45^\circ + \alpha)$. Also $\angle WU'C = \angle V'U'C = \angle V'BC = \angle ABC - \angle ABV' = 45^\circ - \beta$. This yields $\angle U'WC = 180^\circ - (\angle WU'C + \angle U'CW) = 90^\circ + (\beta - \alpha)$. Again, by the Sine Rule, this time in $\Delta U'WC$, we have

$$\frac{1 - w}{\sin \angle WU'C} = \frac{WC}{\sin(45^\circ - \beta)} = \frac{U'C}{\sin \angle U'WC} = \frac{2\cos(45^\circ + \alpha)}{\sin(90^\circ + (\beta - \alpha))},$$


that is,

$$\begin{aligned} 1 - w &= \frac{2\cos(45^\circ + \alpha)\sin(45^\circ - \beta)}{\sin(90^\circ + (\beta - \alpha))} = \frac{(\cos \alpha - \sin \alpha)(\cos \beta - \sin \beta)}{\cos \alpha \cos \beta + \sin \alpha \sin \beta} \\ &= \frac{(1 - \tan \alpha)(1 - \tan \beta)}{1 + \tan \alpha \tan \beta} = \frac{(1 - u)(1 - v)}{1 + uv}, \end{aligned}$$

which, upon solving for w , gives

$$w = \frac{u + v}{1 + uv}.$$

3 A Euclidean geometric proof



As $\angle AOU = 90^\circ = \angle AU'B$ and $\angle OAU = \angle U'AB$ (common), by the AA Similarity Rule, $\triangle AOU \sim \triangle AU'B$. So

$$\frac{AU'}{2} = \frac{AU'}{AB} = \frac{AO}{AU} = \frac{1}{\sqrt{1+u^2}},$$

giving $AU' = 2/\sqrt{1+u^2}$. Hence

$$UU' = AU' - AU = \frac{2}{\sqrt{1+u^2}} - \sqrt{1+u^2} = \frac{1-u^2}{\sqrt{1+u^2}}.$$

Proceeding similarly, $BV' = 2/\sqrt{1+v^2}$ and $VV' = (1-v^2)/\sqrt{1+v^2}$. Let W be the point of intersection of $U'V'$ and OC , and set $OW =: w$. Let the extension of $U'V'$ meet the extension of AB at O' . Menelaus' Theorem applied to ΔAOU with the line $O'U'$ gives

$$\frac{w-u}{w} \cdot \frac{OO'}{OO' - 1} \cdot \frac{2/\sqrt{1+u^2}}{(1-u^2)/\sqrt{1+u^2}} = \frac{UW}{OW} \cdot \frac{OO'}{AO'} \cdot \frac{AU'}{UU'} = 1.$$

This yields

$$\frac{1}{\partial \partial'} = 1 - \frac{2}{1-\mu^2} \cdot \frac{w-u}{w}. \quad (2)$$

Similarly, Menelaus' Theorem applied to ΔBOV with the line $O'U'$ gives

$$\frac{w-v}{w} \cdot \frac{OO'}{OO'+1} \cdot \frac{2/\sqrt{1+v^2}}{(1-v^2)/\sqrt{1+p^2}} = \frac{VW}{OW} \cdot \frac{OO'}{BO'} \cdot \frac{BV'}{VV'} = 1.$$

This yields

$$\frac{1}{OO'} = \frac{2}{1-v^2} \cdot \frac{w-v}{w} - 1. \quad (3)$$

Equating the right-hand sides of (2) and (3) gives $w = \frac{u+v}{1+uv}$.

4 A projective geometric proof

We recall the notion of the cross ratio in projective geometry. If A, B, C, D are collinear points that are projected along four concurrent lines meeting at P , to the collinear points A', B', C', D' , respectively, then we know that the cross ratio is preserved, that is,

$$(A, B; C, D) := \frac{AC}{AD} / \frac{BC}{BD} = \frac{A'C'}{A'D'} / \frac{B'C'}{B'D'} =: (A', B'; C', D').$$


Recall that this is an immediate consequence of the Sine Rule for triangles, using which one can see that

$$\begin{aligned} \frac{AC}{AP} &= \frac{\sin \angle APC}{\sin \angle PCA}, & \frac{AD}{AP} &= \frac{\sin \angle APD}{\sin \angle PDA}, \\ \frac{BD}{BP} &= \frac{\sin \angle BPD}{\sin \angle PDB}, & \frac{BC}{BP} &= \frac{\sin \angle BPC}{\sin \angle PCB}, \end{aligned}$$


and so

$$(A, B; C, D) = \frac{\sin \angle APC}{\sin \angle APD} / \frac{\sin \angle BPC}{\sin \angle BPD}.$$


In light of this invariance, we refer to the cross ratio of the four concurrent lines instead of particular collinear points on the lines.



We also recall Chasles' Theorem, which says that if A_1, A_2, A_3, A_4 are four fixed points on a circle, and P is a movable point, then the cross ratio of the lines PA_1, PA_2, PA_3, PA_4 is a constant. This is an immediate consequence of the fact that a chord of a circle subtends equal angles at any point on its major (or minor) arc.



We refer to the geometric diagram for relativistic velocity addition below, with the labelling of points shown. X is the point of intersection of $P'A_3$ with OA_4 .



As $\Delta OP'X$ is a right angled triangle, it follows that

$$OX = \cos \angle P'OX = \sin \angle POP' = \sin(2\alpha) = \frac{2 \tan \alpha}{1 + (\tan \alpha)^2} = \frac{2u}{1 + u^2}.$$

Hence

$$\begin{aligned} UX &= OX - OU = \frac{2u}{1 + u^2} - u = u \cdot \frac{1 - u^2}{1 + u^2} \quad \text{and} \\ WX &= OX - OW = \frac{2u}{1 + u^2} - w. \end{aligned}$$

By Chasles' Theorem, we have

$$\frac{u}{1} / \frac{u-v}{1-v} = \frac{OU}{OA_4} / \frac{VU}{VA_4} = \frac{UX}{UA_4} / \frac{WX}{WA_4} = \frac{u \cdot \frac{1-u^2}{1+u^2}}{1-u} / \frac{u \cdot \frac{2u}{1+u^2} - w}{1-w}.$$


Solving for w , this yields $w = \frac{u+v}{1+uv}$.

5 A few remarks

We remark that the projective perspective also sheds light on the (algebraically easily verified) formula

$$u \oplus v = \frac{1}{u} \oplus \frac{1}{v}.$$

Indeed, let us see the picture below, where U' , V' are the images of the points U , V , respectively, under inversion in the circle.



Let $OS =: 1/w$. By the preservation of the cross-ratio for the four collinear lines AP , AQ , AR , AS , we obtain

$$(P, Q; R, S) = (O, V; U, S) = \frac{u}{1/w} / \frac{u - v}{(1/w) - v}.$$

On the other hand, by the preservation of the cross-ratio for the four collinear lines BP , BQ , BR , BS , we obtain

$$(P, Q; R, S) = (O, V'; U', S) = \frac{1/u}{1/w} / \frac{(1/v) - (1/u)}{(1/v) - (1/w)}.$$

Thus

$$\frac{u}{1/w} / \frac{u - v}{(1/w) - v} = (P, Q; R, S) = \frac{1/u}{1/w} / \frac{(1/v) - (1/u)}{(1/v) - (1/w)},$$

which gives $w = \frac{u + v}{1 + uv}$.

We also mention that although we have been considering $u, v \in [0, 1]$ for our pictures, one may in fact take $u, v \in [-1, 1]$ without any essential change in our derivations. The operation \oplus is associative and the set $(-1, 1)$ is a group with the operation \oplus .

Reference

- [1] Jerzy Kocik. Geometric diagram for relativistic addition of velocities. *American Journal of Physics*, volume 80, number 8, page 737, 2012.

Amol Sasane
Department of Mathematics
London School of Economics
Houghton Street
London WC2A 2AE, United Kingdom
e-mail: sasane@lse.ac.uk

Victor Ufnarovski
Center for Mathematical Sciences
Lund University
SE-221 00 Lund, Sweden
e-mail: victor.ufnarovski@math.lth.se