Zeitschrift: Elemente der Mathematik

Herausgeber: Schweizerische Mathematische Gesellschaft

Band: 71 (2016)

Heft: 4

Artikel: A lower bound for the isoperimetric deficit

Autor: Cufí, Julià / Reventós, Agustí

DOI: https://doi.org/10.5169/seals-630628

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 30.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

A lower bound for the isoperimetric deficit

Julià Cufí and Agustí Reventós¹

Agusti Reventós is professor of Geometry at the Universitat Autònoma de Barcelona. He has translated to Catalan and commented the Disquisitiones generales circa superficies curvas by C.F. Gauss, with Carlos Rodriguez. He is editor, with A.M. Naveira, of the Selected Works of Luis Antonio Santaló. He is interested in affine maps, Euclidean motions and quadrics.

Julià Cufí received his Ph.D. in Mathematics from Barcelona University. He is emeritus professor at the Department of Mathematics of the Universitat Autònoma de Barcelona. His main field of interest is the theory of functions of complex variables. He has published several research papers in this field and a textbook on Complex Analysis with J. Bruna.

1 Introduction

Let K be a plane compact convex set of area F with boundary curve $C = \partial K$ of length L. As it is well known, the isoperimetric inequality states

$$F \leq L^2/4\pi$$
,

with equality only for discs.

Für jede Figur K in der Ebene mit Umfang L und Fläche F gilt die isoperimetrische Ungleichung $\Delta := L^2 - 4\pi F \ge 0$. Gleichheit gilt genau für Kreise. Hurwitz gelang 1902 nicht nur ein eleganter Beweis der isoperimetrischen Ungleichung mit Hilfe von Fourier-Reihen, er bewies zudem eine obere Schranke für das isoperimetrische Defizit Δ , indem er die Evolute der Kurve ins Spiel brachte. 1920 fand Bonnesen eine untere Schranke für Δ , nämtlich $\pi (R-r)^2 \le \Delta$, wobei R und r den Um- respektive den Inkreisradius der Randkurve C der betrachteten Figur K bezeichnen. In der vorliegenden Arbeit wird eine andere untere Schranke für Δ bewiesen: Diese ergibt sich aus der Differenz der von C umrandeten Fläche und der Fläche welche die Pedalkurve von C bezüglich des Steiner-Punktes von C einschliesst. Das Resultat verbessert damit Abschätzungen z.B. von Groemer. Es wird zudem bestimmt, für welche Kurven die neue Abschätzung scharf ist.

¹The work was partially supported by grants MTM2012-36378 and MTM2012-34834 (MEC).

Introducing the quantity $\Delta = L^2 - 4\pi F$, called the *isoperimetric deficit*, the above inequality can be written as $\Delta \geq 0$. In some sense Δ measures the extend to which the convex set is away from a disc. It is interesting to know upper and lower bounds for Δ in terms of quantities associated to K.

Hurwitz, in his paper about the use of Fourier series in some geometrical problems [3], proves the following inequality, which is a sort of reverse isoperimetric inequality and provides an upper bound for Δ ,

$$0 \le \Delta \le \pi |F_e|,\tag{1}$$

where F_e is the algebraic area enclosed by the evolute of C. Equality holds when C is a circle or parallel to an astroid.

Recall that the *evolute* of a plane curve is the locus of its centers of curvature or, equivalently, the *envelope* of all the normals to this curve (i.e., the tangents to the evolute are the normals to the curve).

As for lower bounds, along the 1920s Bonnesen provided some inequalities of the type $\Delta \ge B$, where B is a non-negative quantity associated to the convex set vanishing only for circles. Moreover these quantities have a relevant geometrical meaning (see [4]).

In this note we prove a Bonnesen-style inequality which gives a lower bound for the isoperimetric deficit in terms of the difference between the area enclosed by the pedal curve of C with respect to the Steiner point of K, and the area enclosed by C.

The *pedal curve* of a plane curve C with respect to a fixed point O is the locus of points X so that the line OX is perpendicular to the tangent to C passing through X. The *Steiner point* of a plane convex set K, or the curvature centroid of K, is the center of mass of ∂K with respect to the density function that assigns to each point of ∂K its curvature.

Let A be the area enclosed by the pedal curve of $C = \partial K$ with respect to the Steiner point of K. In Theorem 3.1 it is proved that

$$\Delta \ge 3\pi (A - F). \tag{2}$$

So, the quantity $3\pi(A - F)$ is a lower estimate of the isoperimetric deficit. Since $A \ge F$, this inequality implies the isoperimetric one. Moreover A = F only for circles and so $\Delta = 0$ implies C is a circle.

We point out that inequality (15) shows that Theorem 3.1 improves Theorem 4.3.1 in [2]. Moreover our lower bound has a very clear geometric significance.

For the special case of convex sets of constant width we obtain the inequality

$$\Delta \ge \frac{32}{9}\pi(A - F),\tag{3}$$

which in turn improves an inequality on page 144 of [2], as inequality (17) shows.

We also consider when equality holds in (2) and (3). In Corollaries 4.2 and 4.4 it is shown that this is so for circles and curves which are parallel to an astroid or to a hypocycloid of three cusps, respectively.

Finally in Propositions 5.1 and 5.3 we prove that for convex curves C parallel to an astroid or a hypocycloid of three cusps, the evolute of C is similar, with ratio 2 or 3 respectively, to the corresponding astroid or hypocycloid.

2 Preliminaries

Support function

A straight line G in the plane is determined by the angle ϕ that the direction perpendicular to G makes with the positive x-axis and the distance $p = p(\phi)$ of G from the origin. The equation of G then takes the form

$$x\cos\phi + y\sin\phi - p = 0. (4)$$

Equation (4), when $p = p(\phi)$ varies with ϕ , is the equation of a family of lines. If we assume that the 2π -periodic function $p(\phi)$ is differentiable, the envelope of the family is obtained from (4) and the derivative of its left-hand side, as follows:

$$-x\sin\phi + y\cos\phi - p' = 0, \quad p' = dp/d\phi. \tag{5}$$

From (4) and (5) we arrive at a parametric representation of the envelope of the lines (4):

$$x = p\cos\phi - p'\sin\phi, \qquad y = p\sin\phi + p'\cos\phi.$$

If the envelope is the boundary ∂K of a convex set K and the origin is an interior point of K, then $p(\phi)$ is called the *support function* of K (or the support function of the convex curve ∂K).

Since $dx = -(p + p'') \sin \phi \, d\phi$ and $dy = (p + p'') \cos \phi \, d\phi$ (we here assume that the function p is of class C^2), arclength measure on ∂K is given by

$$ds = \sqrt{dx^2 + dy^2} = |p + p''| \, d\phi \tag{6}$$

and the radius of curvature ρ by

$$\rho = \frac{ds}{d\phi} = |p + p''|.$$

It is well known (see for instance [5], page 3) that a necessary and sufficient condition for a periodic function p to be the support function of a convex set K is that p+p''>0. Finally, it follows from (6) that the length of a closed convex curve that has support function p of class C^2 is given by

$$L = \int_0^{2\pi} p \, d\phi. \tag{7}$$

The area of the convex set K is expressed in terms of the support function by

$$F = \frac{1}{2} \int_{\partial K} p ds = \frac{1}{2} \int_{0}^{2\pi} p(p + p'') d\phi = \frac{1}{2} \int_{0}^{2\pi} p^{2} d\phi - \frac{1}{2} \int_{0}^{2\pi} p'^{2} d\phi.$$
 (8)

For any curve C given by $(x(\phi), y(\phi))$, convex or not, we will say that $p(\phi)$ is the generalized support function of C when

$$x(\phi) = p(\phi)\cos(\phi) - p'(\phi)\sin(\phi),$$

$$y(\phi) = p(\phi)\sin(\phi) + p'(\phi)\cos(\phi).$$

Note that $p(\phi)$ is not necessarily a distance, as it happens when we define the support function of a convex set. In fact, $|p(\phi)|$ is the distance from the origin to the tangent to C at the point $(x(\phi), y(\phi))$.

It is easy to see that the generalized support function $p_e(\phi)$ of the evolute of $C = \partial K$ is $p_e(\phi) = -p'(\phi + \pi/2)$, where $p(\phi)$ is the support function of C, see [1]. Hence, assuming $p(\phi)$ is a C^3 -function, the algebraic area F_e enclosed by the evolute of C is given by

$$F_e = \frac{1}{2} \int_0^{2\pi} p'(p' + p''') d\phi = \frac{1}{2} \int_0^{2\pi} p'^2 d\phi - \frac{1}{2} \int_0^{2\pi} p''^2 d\phi.$$

Steiner point

The Steiner point of a convex set K of the Euclidean plane is defined by

$$S(K) = \frac{1}{\pi} \int_0^{2\pi} (\cos \phi, \sin \phi) p(\phi) d\phi,$$

where $p(\phi)$ is the support function of ∂K (see [2]).

Thus, if

$$p(\phi) = a_0 + \sum_{n \ge 1} a_n \cos n\phi + b_n \sin n\phi, \tag{9}$$

is the Fourier series of the 2π -periodic function $p(\phi)$, the Steiner point is

$$S(K) = (a_1, b_1).$$

The Steiner point of K is also known as the curvature centroid of K because under appropriate smoothness conditions it is the center of mass of ∂K with respect to the density function that assigns to each point of ∂K its curvature.

The relation between the support function $p(\phi)$ of a convex set K and the support function $q(\phi)$ of the same convex set but with respect to a new reference with origin at the point (a, b), and axes parallel to the previous x- and y-axes, is given by

$$q(\phi) = p(\phi) - a\cos\phi - b\sin\phi$$
.

Hence, taking the Steiner point as a new origin, we have

$$q(\phi) = a_0 + \sum_{n>2} a_n \cos n\phi + b_n \sin n\phi.$$

We recall that the *Steiner ball* of *K* is the ball whose center is the Steiner point and whose diameter is the mean width of *K*.

Pedal curve

If the curve C is given in Cartesian coordinates as the envelope of the lines $x \cos \phi + y \sin \phi - p(\phi) = 0$, then the pedal curve $P = P(\phi)$ of C with respect to the origin, is given by

$$\mathcal{P}(\phi) = (p(\phi)\cos\phi, p(\phi)\sin\phi),$$

or, in polar coordinates, by $r = p(\phi)$.

In particular, if C is closed, the area enclosed by \mathcal{P} is

$$A = \frac{1}{2} \int_0^{2\pi} p^2 \, d\phi. \tag{10}$$

If F is the area enclosed by C, we obviously have $A \ge F$ with equality if and only if C is a circle.

3 A lower bound for the isoperimetric deficit

We proceed now to provide a lower bound for the isoperimetric deficit.

Theorem 3.1. Let K be a compact convex set of area F with boundary curve $C = \partial K$ of class C^2 and length L. Let A be the area enclosed by the pedal curve of C with respect to the Steiner point S(K). Then

$$\Delta \ge 3\pi (A - F),\tag{11}$$

where $\Delta = L^2 - 4\pi F$ is the isoperimetric deficit.

Proof. Let $p(\phi)$ be the support function of C, with respect to an orthonormal reference with origin in the Steiner point, and axes parallel to the x- and y-axes.

We know that the Fourier series of $p(\phi)$, is

$$p(\phi) = a_0 + \sum_{n \ge 2} a_n \cos n\phi + b_n \sin n\phi.$$

By Parseval's identity we have

$$\frac{1}{2\pi} \int_0^{2\pi} p^2 d\phi = a_0^2 + \frac{1}{2} \sum_{n \ge 2} (a_n^2 + b_n^2), \tag{12}$$

and similar expressions for p' and p''. Concretely we have

$$\int_0^{2\pi} p'^2 d\phi = \pi \sum_{n \ge 2} n^2 (a_n^2 + b_n^2), \qquad \int_0^{2\pi} p''^2 d\phi = \pi \sum_{n \ge 2} n^4 (a_n^2 + b_n^2). \tag{13}$$

Hence, the isoperimetric deficit $\Delta = L^2 - 4\pi F$, according to (7) and (8), is given by

$$\Delta = \left(\int_0^{2\pi} p \, d\phi\right)^2 - 2\pi \int_0^{2\pi} p^2 \, d\phi + 2\pi \int_0^{2\pi} p'^2 \, d\phi$$
$$= 2\pi^2 \sum_{n \ge 2} (n^2 - 1)(a_n^2 + b_n^2) \ge \frac{3\pi^2}{2} \sum_{n \ge 2} n^2 (a_n^2 + b_n^2) = \frac{3\pi}{2} \int_0^{2\pi} p'^2 \, d\phi.$$

But it follows from (8) and (10) that

$$\frac{1}{2} \int_0^{2\pi} p'^2 d\phi = \frac{1}{2} \int_0^{2\pi} p^2 d\phi - F = A - F,$$

and hence

$$\Delta \geq 3\pi (A - F)$$
.

The above proof shows that $\Delta = 0$ if and only if $p(\phi) = a_0$, that is, when C is a circle.

Although the constant 3π appearing in Theorem 3.1 cannot be improved for general convex sets, it is possible to obtain a stronger inequality for special types of convex sets. For instance, for convex sets of constant width we have the following result.

Proposition 3.2. Let K be a compact convex set of constant width in the hypothesis of Theorem 3.1. Then

$$\Delta \geq \frac{32}{9}\pi(A-F).$$

Proof. Since constant width means $p(\phi) + p(\phi + \pi)$ is constant and

$$p(\phi) + p(\phi + \pi) = 2\sum_{n=0}^{\infty} (a_{2n}\cos 2n\phi + b_{2n}\sin 2n\phi)$$

it follows that $a_n = b_n = 0$ for all even n > 0.

Introducing this in the proof of Theorem 3.1 the result follows.

Relationship with the L^2 metric

Consider now the quantity $\delta_2(K)$ equal to the distance in $L^2(S^1)$, where S^1 is the unit circle, between the support function of K and the support function of the Steiner ball of K. It is known, see [2] Theorem 4.3.1, that

$$\Delta \ge 6\pi \,\delta_2(K)^2. \tag{14}$$

We can state now the following inequality

$$3\pi (A - F) \ge 6\pi \delta_2(K)^2. \tag{15}$$

To prove this we first observe that

$$\delta_2(K)^2 = \pi \sum_{n=2}^{\infty} (a_n^2 + b_n^2),$$

where a_n , b_n are Fourier coefficients of the support function of K, $p(\phi)$, as in (9) (see page 142 of [2]). Moreover the proof of Theorem 3.1 shows that

$$3\pi (A - F) = \frac{3\pi^2}{2} \sum_{n>2} n^2 (a_n^2 + b_n^2),$$

hence inequality (15) follows.

So we have

$$\Delta \ge 3\pi (A - F) \ge 6\pi \delta_2(K)^2$$

which improves the inequality (14).

For compact convex sets of constant width it is known that

$$\Delta \ge 16\pi \,\delta_2(K)^2 \tag{16}$$

(see page 144 of [2]).

We can state now the following inequality

$$\frac{32}{9}\pi(A-F) \ge 16\pi\,\delta_2(K)^2. \tag{17}$$

The proof is the same as for (15) taking into account that now the even coefficients vanish. So we have

$$\Delta \ge \frac{32}{9}\pi (A - F) \ge 16\pi \delta_2(K)^2$$

which improves the inequality (16).

4 Equality of the lower bound with the isoperimetric deficit

Now we study the case of equality in Theorem 3.1. It is clear from the proof that $\Delta = 3\pi (A - F)$ if and only if

$$p(\phi) = a_0 + a_2 \cos 2\phi + b_2 \sin 2\phi.$$

In order to characterize the curves with this type of support function we recall that the parametric equations of an *astroid* (a 4-cusped hypocycloid) are

$$x(\phi) = 2a\sin^3(\phi),$$

$$y(\phi) = 2a\cos^3(\phi),$$

for some constant $a \in \mathbb{R}$, $a \neq 0$, with $0 \leq \phi \leq 2\pi$. From this it is easy to see that the generalized support function $p(\phi)$ of the astroid is $p(\phi) = a \sin(2\phi)$, where ϕ is the angle between the normal $(-y'(\phi), x'(\phi))$ and the positive x-axis.

This implies that the curves with generalized support function given by

$$q(\phi) = b + p(\phi) = b + a\sin(2\phi),$$

where $b \in \mathbb{R}$, are parallel to an astroid. The distance between these curves and the astroid is |b|.

We have the following result.

Proposition 4.1. *Let*

$$p(\phi) = a_0 + a_2 \cos(2\phi) + b_2 \sin(2\phi)$$

be the support function of a closed convex curve C of length L, with $a_2^2 + b_2^2 \neq 0$. Then the interior parallel curve to C at distance $L/2\pi$ is an astroid.

Proof. We make the change of variable $u = \phi - \phi_0 + \frac{\pi}{4}$, where

$$\tan 2\phi_0 = \frac{b_2}{a_2}.$$

Then

$$\sin 2u = \cos 2(\phi - \phi_0) = \cos 2\phi \cos 2\phi_0 + \sin 2\phi \sin 2\phi_0$$
$$= \cos 2\phi \frac{a_2}{\pm \sqrt{a_2^2 + b_2^2}} + \sin 2\phi \frac{b_2}{\pm \sqrt{a_2^2 + b_2^2}}.$$

Hence

$$p(u) = a_0 \pm a \sin 2u$$

where $a = \sqrt{a_2^2 + b_2^2}$. This shows that the given curve is parallel to an astroid at distance $|a_0|$. By the condition of convexity, $p + p'' = a_0 \mp 3a \sin 2u > 0$, and so a_0 is positive. Since $L = \int_0^{2\pi} p(\phi) d\phi = 2\pi a_0$, the proposition is proved.

Corollary 4.2. Equality in Theorem 3.1 holds if and only if C is a circle or a curve parallel to an astroid.

Proof. As we have said, equality in (11) holds when

$$p(\phi) = a_0 + a_2 \cos 2\phi + b_2 \sin 2\phi$$
.

If $a_2 = b_2 = 0$, $p(\phi) = a_0$ is the support function of a circle. If $a_2^2 + b_2^2 \neq 0$, the result follows from Proposition 4.1.

Now we study the case of equality in Proposition 3.2.

It is clear from the proof of this proposition that equality holds if and only if

$$p(\phi) = a_0 + a_3 \cos 3\phi + b_3 \sin 3\phi$$
.

In order to characterize the curves with this type of support function we recall that the parametric equations of a *hypocycloid of three cusps*, with respect to a suitable orthogonal system, are

$$x(t) = -2a\cos t - a\cos 2t$$

$$v(t) = -2a\sin t + a\sin 2t$$

with $a \in \mathbb{R}$, $a \neq 0$, $t \in [0, 2\pi]$.

The relationship between the parameter t and the angle $\phi(t)$ between the normal vector (-y'(t), x'(t)) and the positive x-axis is

$$\phi(t) = \alpha(t) - \frac{\pi}{2},$$

where $\alpha(t)$ denotes the angle between the tangent vector (x'(t), y't) and the positive x-axis.

Hence

$$\tan \phi(t) = -\cot \alpha(t) = \frac{\sin t + \sin 2t}{\cos t - \cos 2t} = \cot \frac{t}{2}$$

and so

$$t = \pi - 2\phi(t)$$
.

On the other hand, the generalized support function $p(\phi)$ of the hypocycloid must verify

$$\begin{pmatrix} x(\phi) \\ y(\phi) \end{pmatrix} = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} p(\phi) \\ p'(\phi) \end{pmatrix},$$

so

$$\begin{pmatrix} p(\phi) \\ p'(\phi) \end{pmatrix} = \begin{pmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} -2a\cos(\pi - 2\phi) - a\cos 2(\pi - 2\phi) \\ -2a\sin(\pi - 2\phi) + a\sin 2(\pi - 2\phi) \end{pmatrix}.$$

Then, using standard addition trigonometric formulas, it follows

$$p(\phi) = a\cos(3\phi)$$
.

Proposition 4.3. *Let*

$$p(\phi) = a_3 \cos 3\phi + b_3 \sin 3\phi$$

be the generalized support function of a closed curve C, with $a_3^2 + b_3^2 \neq 0$. Then C is a hypocycloid of three cusps.

Proof. We make the change of variable given by $u = \phi - \phi_0$, where

$$\tan 3\phi_0 = \frac{b_3}{a_3}.$$

Then, an easy computation gives

$$p(u) = a\cos(3u),$$

where $a = \frac{a_3}{\cos \phi_0}$, and the proposition follows.

Corollary 4.4. Equality in Proposition 3.2 holds if and only if C is a circle or a curve parallel to a hypocycloid of three cusps.

Proof. We have seen that equality in Proposition 3.2 holds when

$$p(\phi) = a_0 + a_3 \cos 3\phi + b_3 \sin 3\phi$$
.

If $a_3 = b_3 = 0$, $p(\phi) = a_0$ is the support function of a circle. If $a_3^2 + b_3^2 \neq 0$, the result follows from Proposition 4.3.

Remark 4.5. As it is well known (see for instance page 8 of [5]) the area F_r enclosed by the interior parallel at distance r to a closed curve is given by

$$F_r = F - Lr + \pi r^2$$

where L and F are respectively the length and the area corresponding to the given curve. In particular, if $r=L/2\pi$, we get

$$F_{L/2\pi} = F - \frac{L^2}{4\pi},$$

or, equivalently

$$\Delta = -4\pi \, F_{L/2\pi},$$

which gives a geometrical interpretation of the isoperimetric deficit.

In particular the isoperimetric inequality $\Delta \geq 0$ is equivalent to $F_{L/2\pi} \leq 0$, a fact that suggests a more geometric proof of the isoperimetric inequality, by showing that in the process of collapsing, the curve reverses orientation. Moreover, $F_{L/2\pi} = 0$ holds only for circles.

Remark 4.6. Combining Theorem 3.1 with Hurwitz' inequality (1) we have the relation

$$A - F \le \frac{1}{3} |F_e|,\tag{18}$$

with equality for circles or curves parallel to an astroid.

5 Parallel curves and evolutes

We have seen the role played by the convex curves parallel to an astroid or to a hypocycloid of three cusps. For such curves C we show that there is a quite surprising relationship between the parallel curve at distance $L/2\pi$ to C and the evolute of C.

Proposition 5.1. *Let*

$$p(\phi) = a_0 + a_2 \cos(2\phi) + b_2 \sin(2\phi)$$

be the support function of a closed convex curve C of length L. Then the evolute of C and the interior parallel curve to C at distance $L/2\pi$, are similar with ratio 2.

Proof. We shall see that there is a similarity, composition of a rotation with a homothecy, applying the parallel curve on the evolute. We may assume, by the proof of Proposition 4.1, $p(\phi) = a_0 + a \sin(2\phi)$. The generalized support function of the parallel curve to C at distance $L/2\pi = a_0$ is $q(\phi) = a \sin(2\phi)$ and the corresponding one to the evolute of C is

$$p_e(\phi) = -p'\left(\phi + \frac{\pi}{2}\right) = 2a\cos(2\phi).$$

The generalized support function of the rotated $3\pi/4$ parallel curve to C is

$$\tilde{p}(\phi) = q\left(\phi - \frac{3\pi}{4}\right) = a\cos(2\phi).$$

Hence this rotated curve is homothetic, with ratio 2, to the evolute.

Remark 5.2. In particular, the area of the evolute of such a curve is four times the area of the parallel curve at distance $L/2\pi$. The reciprocal is also true. In fact, since Hurwitz' inequality, by Remark 4.5, is equivalent to

$$4|F_{L/2\pi}| - |F_e| \le 0$$
,

the curves for which the area of the evolute is four times the area of the parallel curve at distance $L/2\pi$, are *exactly* circles or curves parallel to an astroid.


Proposition 5.3. *Let*


$$p(\phi) = a_0 + a_3 \cos(3\phi) + b_3 \sin(3\phi)$$


be the support function of a closed convex curve C of length L. Then the evolute of C and the interior parallel curve to C at distance $L/2\pi$, are similar with ratio 3.

Proof. Analogous to that of Proposition 5.1 with $q(\phi) = a\cos(3\phi)$, according to the proof of Proposition 4.3.

The next figures show convex curves with support functions $p(\phi) = 5 + \sin(2\phi)$ and $p(\phi) = 8 + \sin(3\phi)$, their parallel interior curves at distance $L/2\pi = 5$ and $L/2\pi = 8$ respectively, and the corresponding evolutes.







 $p(\phi) = 8 + \sin(3\phi)$

References

- [1] C.A. Escudero and A. Reventós. An interesting property of the evolute. Amer. Math. Monthly, 114(7):623-628, 2007.
- [2] H. Groemer. Geometric applications of Fourier series and spherical harmonics, volume 61 of Encyclopedia of Mathematics and its Applications. Cambridge University Press, 1996.
- [3] A. Hurwitz. Sur quelques applications géométriques des séries de Fourier. Annales scientifiques de l'É.N.S., 19(3e série):357-408, 1902.
- [4] Robert Osserman. Bonnesen-style isoperimetric inequalities. Amer. Math. Monthly, 86(1):1–29, 1979.
- [5] L.A. Santaló. Integral Geometry and Geometric Probability. Cambridge University Press, 2004. Second edition.

Julià Cufí and Agustí Reventós Departament de Matemàtiques Universitat Autònoma de Barcelona E-08193 Bellaterra Barcelona, Catalonia e-mails: jcufi@mat.uab.cat

agusti@mat.uab.cat