

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 72 (2017)
Heft: 2

Artikel: Triangles with sides in arithmetic progression
Autor: Mikami, Kentaro / O'Hara, Jun / Sugawara, Kunio
DOI: <https://doi.org/10.5169/seals-730834>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Triangles with sides in arithmetic progression

Kentaro Mikami, Jun O'Hara and Kunio Sugawara

Kentaro Mikami is a differential geometer and former professor of the Graduate School of Engineering and Resource Science, Akita University, Japan. He graduated from Tohoku University in 1970, where he also earned his Ph.D.

Jun O'Hara is a geometer and professor of the Department of Mathematics and Informatics, Faculty of Science, Chiba University, Japan. He was born in 1963 and graduated from the University of Tokyo where he obtained his Ph.D. in 1991.

Kunio Sugawara is a differential geometer and former professor of the Department of Mathematical Sciences, Osaka Kyoiku University, Japan. He was born in 1960 and graduated from Kyoto University, where he earned his Ph.D.

In this note we study geometric properties of triangles $\triangle ABC$ the sides of which are in arithmetic progression, $CA - BC = AB - CA$. As far as the authors know, this topic does not seem to have been in the spotlight¹. This is why we would like to integrate some known results and (possibly) new results here.

Theorem 1. *Let $\triangle ABC$ be a triangle with edge lengths $a = BC, b = CA, c = AB$. Then, each of the following three conditions is a necessary and sufficient condition so that the sequence of sides a, b, c is an arithmetic progression.*

(i) *The line joining the centroid G and the incenter I is parallel to the edge AC .*

¹with the exception of the study in the case of Heronian triangles, i.e., triangles such that the side lengths and area are all integers ([BG]).

Im Zentrum dieses Artikels stehen Dreiecke, deren Seiten a, b, c eine arithmetische Progression bilden, d.h. $b = (a + c)/2$. Derartige Dreiecke sind weder besonders selten noch besonders häufig anzutreffen: In der Menge der Äquivalenzklassen ähnlicher Dreiecke bilden sie einen Raum der Kodimension eins, genau wie gleichschenklige oder rechtwinklige Dreiecke. Im Gegensatz zu den eben genannten Klassen scheinen aber Dreiecke, deren Seiten eine arithmetische Progression bilden, wenig untersucht worden zu sein, obwohl sie, wie die Autoren der vorliegenden Arbeit zeigen, hübsche geometrische Eigenschaften aufweisen, über die sie sich charakterisieren lassen.




Fig. 1

Fig. 2

- (ii) The line joining the incenter I and the circumcenter O is perpendicular to the line BI (see Figure 1).
- (iii) The midpoint of BC and the excenter opposite to B lies on the edge AC (see Figure 2).

The condition (i) is given in an exercise in Section 8.6.7 on page 120 of a note by Paul Yiu which is available through the web ([Y]). The authors found the conditions (ii) and (iii) accidentally when they were drawing figures with softwares such as GeoGebra, Maple, and Mathematica. Later, they got informed that the condition (ii) has been given in Problem 1 of the Indian National Mathematical Olympiad 2006. Three proofs of (ii), using Stewart's theorem, Ptolemy's theorem, and Euler's theorem and the formula of $\cos(B/2)$ respectively, can be found in [I]. We give a (possibly) new proof of (ii), a little bit more elementary, but still using Euler's theorem. As for the condition (iii), the authors do not know references.

Let us first prepare formulae which are needed in our proof. Let r, R denote the radii of the incircle and the circumcircle respectively, S the area of $\triangle ABC$, and $s = (a + b + c)/2$. Then, Heron's formula states

$$S = \sqrt{s(s-a)(s-b)(s-c)}. \quad (1)$$

Since $|\triangle ABC| = |\triangle IAB| + |\triangle IBC| + |\triangle IAC|$ we have

$$r = \frac{2S}{a+b+c} = \frac{S}{s}. \quad (2)$$

The law of sines, $R = a/(\sin A)$, implies

$$R = \frac{abc}{4S}. \quad (3)$$

Finally, Euler's theorem (see, for example, [J, p. 186]) states

$$IO^2 = R^2 - 2rR. \quad (4)$$

Proof. (i) The line GI is parallel to the line AC if and only if the heights of G and I above the line AC are the same, which should be one third of the height of the vertex B . Therefore the condition (i) is equivalent to saying that $|\triangle IAC| = |\triangle ABC|/3$, which, by (2), is equivalent to $br = (a + b + c)r/3$.

(ii) Let D be the foot of the perpendicular to the edge BC from I (see Figure 1). Since D is the tangent point of the incircle with the side BC , we have $BD = (a - b + c)/2 = s - b$. Therefore,

$$\begin{aligned} OI \perp BI &\iff BI^2 + IO^2 = OB^2 \\ &\stackrel{(4)}{\iff} (BD^2 + r^2) + (R^2 - 2rR) = R^2 \\ &\stackrel{(2)}{\iff} 2rR = (s - b)^2 + \frac{s(s - a)(s - b)(s - c)}{s^2} \\ &\stackrel{(2),(3)}{\iff} \frac{abc}{2s} = \frac{s - b}{s} (s(s - b) + (s - a)(s - c)) \\ &\iff abc = 2(s - b)(2s^2 - (a + b + c)s + ac) = 2(s - b)ac \\ &\iff b = a - b + c, \end{aligned}$$

which completes the proof.

(iii) Let J be the excenter opposite to B , M the intersection of BJ and AC . We show that $2b = a + c$ if and only if $BJ = 2BM$.

Let us compute BJ first. Let H be the foot of perpendicular to the line BC from J (see Figure 1). Since H is the tangent point of the excircle and the line BC , we have $BH = (a + b + c)/2 = s$. On the other hand, since JH is the radius of the excircle, $JH = 2S/(a - b + c) = S/(s - b)$. Therefore,

$$\begin{aligned} BJ^2 &= BH^2 + JH^2 \\ &\stackrel{(1)}{=} s^2 + \frac{s(s - a)(s - b)(s - c)}{(s - b)^2} \\ &= \frac{s}{(s - b)} (s(s - b) + (s - a)(s - c)) \\ &= ac \frac{a + b + c}{a - b + c}. \end{aligned}$$

Next we compute BM . Let E be the intersection of the circumcircle of the triangle ABC and the line BM . Put $m = CM$, $n = AM$, $x = BM$, and $y = ME$. Since the triangle




Fig. 3

BEC is similar to the triangle BAM (see Figure 3), we have $(x + y) : a = c : x$, i.e., $x^2 = ac - xy$. On the other hand, the secant theorem implies $xy = mn$. Since $m : n = a : c$,

$$x^2 = ac - mn = ac - \frac{ab}{a+c} \cdot \frac{cb}{a+c} = \frac{ac}{(a+c)^2} \left((a+c)^2 - b^2 \right),$$

which implies

$$BM^2 = \frac{ac(a+b+c)(a-b+c)}{(a+c)^2}. \quad (5)$$

Therefore,

$$\begin{aligned} BJ = 2BM &\iff (a+c)^2 = 4(a-b+c)^2 \\ &\iff 2b = a+c, \end{aligned}$$

which completes the proof. \square

Remark 2. We can also show the formula (5) by applying the law of cosines to $\triangle BAC$ and $\triangle BAM$.

Acknowledgement

The authors thank the anonymous referee of the previous version of this note for the references [CG] and [I].

References

- [BG] H. Bailey and W. Gosnell, *Heronian triangles with sides in arithmetic progression: An inradius perspective*, *Mathematics Magazine* 85 (2012), 290–294.

- [CG] D. Coxeter and S. Greitzer, *Geometry Revisited*, Mathematical Association of America, Washington, DC, USA, (1967).
- [I] <http://olympiads.hbcse.tifr.res.in/uploads/inmo-2006>
- [J] Roger A. Johnson, *Advanced Euclidean Geometry*, Dover Publ., (2007) [1929]
- [Y] P. Yiu, <http://math.fau.edu/Yiu/EuclideanGeometryNotes.pdf>

Kentaro Mikami
Graduate School of Engineering and Resource Science
Akita University
1-1 Tegata Gakuen-machi
Akita City, 010-8502 JAPAN
e-mail: mikami@math.akita-u.ac.jp

Jun O'Hara
Department of Mathematics and Informatics
Faculty of Science
Chiba University
1-33 Yayoi-cho, Inage
Chiba, 263-8522, JAPAN
e-mail: ohara@math.s.chiba-u.ac.jp

Kunio Sugawara
Department of Mathematical Sciences
Osaka Kyoiku University
4-698-1 Asahigaoka
Kashiwara, Osaka 582-8582 JAPAN
e-mail: sugahara@cc.osaka-kyoiku.ac.jp