

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 72 (2017)
Heft: 4

Artikel: Gerretsen and the Finsler-Hadwiger inequality
Autor: Wanner, Gerhard
DOI: <https://doi.org/10.5169/seals-730844>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 21.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

**Short note Gerretsen and the
Finsler–Hadwiger inequality**

Gerhard Wanner

The inequalities. The beautiful paper [2] (1953) by J.C.H. Gerretsen, written in Dutch language, is of difficult access¹. It starts by stating Weitzenböck's inequality from 1919 [4]

$$a^2 + b^2 + c^2 \geq 4\mathcal{A}\sqrt{3}, \quad (1)$$

where \mathcal{A} denotes the area of a triangle with sides a, b, c . Next follows its sharpening by P. Finsler and H. Hadwiger [1] from 1938

$$a^2 + b^2 + c^2 \geq 4\mathcal{A}\sqrt{3} + Q, \quad (2)$$

where

$$Q = (a - b)^2 + (b - c)^2 + (c - a)^2 \quad (3)$$

can be seen as a measure for the “*ongelijkzijdigheidsgraad*”² of the triangle. In both inequalities we have equality only for the equilateral triangle. Gerretsen then elaborates in Art. 2 the original proof of Finsler and Hadwiger (“*Het bewijs ... is nogal listig*”) and gives simpler proofs for (1) and (2) in Arts. 3 and 4.

A further consequence. In Art. 5 Gerretsen continues as follows: From the Cosine Theorem

$$a^2 = b^2 + c^2 - 2bc \cos \alpha = b^2 + c^2 - 4\mathcal{A} \cot \alpha$$

we obtain by cyclic permutations and addition³

$$\sum a^2 = 4\mathcal{A} \sum \cot \alpha$$

¹The author is grateful to W. Hundstorfer, Amsterdam, for providing him with a copy (“The scanner is not used often anymore, everything is on the internet. Well ... *almost* everything”).

²literally “degree of unequilaterality”; Gerretsen adds: “as long as one does not find this word to crazy [wanneer men dit woord niet te gek vindt]”.

³Gerretsen leaves the meaning of the notations $\sum a^2 = a^2 + b^2 + c^2$ or $\sum \cot \alpha = \cot \alpha + \cot \beta + \cot \gamma$ or $\sum ab = ab + bc + ca$ to the intelligence of the reader.


so that

$$\sum a^2 - Q = 2 \sum ab - \sum a^2 = 4A \sum \left(\frac{1}{\sin \alpha} - \cot \alpha \right) = 4A \sum \tan \frac{\alpha}{2}, \quad (4)$$

thus the inequality of Finsler and Hadwiger means that

$$\tan \frac{\alpha}{2} + \tan \frac{\beta}{2} + \tan \frac{\gamma}{2} \geq \sqrt{3} \quad (5)$$

for all angles α, β, γ of a triangle.



$$\text{Proof of } \tan \frac{\alpha}{2} + \tan \frac{\beta}{2} + \tan \frac{\gamma}{2} \geq \sqrt{3}$$

Backward proof. However, we believe the inequality (5) at once: the angles α, β, γ of a triangle satisfying $\alpha + \beta + \gamma = \pi$ form in \mathbb{R}^3 an equilateral triangle for which (α, β, γ) are the barycentric coordinates (see the Figure above). The three terms of (5) are convex functions on this triangle, each one strictly convex in another direction, therefore their sum is strictly convex with a 120° -symmetry. Thus its minimal point, which must be unique, can only lie in the center where $\alpha = \beta = \gamma = \frac{\pi}{3}$ and the sum is $\sqrt{3}$. Reading now the proof in (4) backwards, we obtain another nice proof of the Finsler–Hadwiger inequality.

Epilog. In Art. 7 Gerretsen proves “his” famous inequalities with the standard proof from the Euler distances of the “bijzondere punten” (see, e.g., [3]), and, inversely, Lukarevski [3] proves the Finsler–Hadwiger inequality from Gerritsen’s inequalities.

References

- [1] P. Finsler, H. Hadwiger, *Einige Relationen im Dreieck*, Comm. Math. Helv. 10 (1938) 316–326.
- [2] J.C.H. Gerretsen, *Ongelijkheden in de Driehoek*, Nieuw Tijdschrift voor Wiskunde 41 (1953) 1–7.
- [3] M. Lukarevski, *An alternate proof of Gerretsen’s inequalities*, Elemente der Math. 72 (2017) 2–8.
- [4] R. Weitzenböck, *Über eine Ungleichung in der Dreiecksgeometrie*, Math. Zeitschr. 5 (1919) 137–146.

Gerhard Wanner
gerhard.wanner@unige.ch