

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 73 (2018)  
**Heft:** 3

**Artikel:** 3 in 1 : a simple way to prove that  $\pi$ ,  $\ln(r)$  and 2 are irrational  
**Autor:** Müller, Jürgen / Müller, Tom  
**DOI:** <https://doi.org/10.5169/seals-780944>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**Short note**

**3 in 1: A simple way to prove that  $e^r$ ,  
 $\ln(r)$  and  $\pi^2$  are irrational**

Jürgen Müller and Tom Müller

In 1947, apparently inspired by the classical Hermite method to show the transcendence of  $e$  (see, e.g., [3, Chapter 2]), Ivan Niven published his famous note “A simple proof that  $\pi$  is irrational” [6]. The key to this proof is the use of sums of different derivatives of special polynomials in order to construct a sequence  $(P_n)$  of polynomials with integer coefficients and of degree  $\leq n$  fulfilling  $0 < |P_n(\pi)| < 1/n!$ . If, in this setting, we suppose  $\pi = a/b$  to be a rational number, then  $b^n P_n(\pi)$  would be an integer number with  $0 < |b^n P_n(\pi)| < 1$  for all large  $n$ . This is of course impossible and hence  $\pi$  is irrational.

Soon after, Iwamoto [4] and Butlewski [2] exploited variations of Niven’s method and constructed other approximation polynomials in order to get simple irrationality proofs for  $\pi^2$ , resp.  $e^k$  for any integer  $k \neq 0$ . In all cases the used polynomials seem to appear from nowhere and to show that they are actually in  $\mathbb{Z}[x]$  is more or less tricky.

In this note we take a new look at the classic analytic irrationality proofs for  $\pi^2$  and the integer powers of  $e$ , showing that the required approximation polynomials are generated by one single integral expression. Our approach makes it obvious how the polynomials come into existence, why they have integer coefficients and that the irrationality proofs for  $\pi$ ,  $\pi^2$  and  $e^k$  are only different special cases derived from the same general formula.

We start by studying the integral

$$I_k(z) := z^{k+1} \int_0^1 t^k e^{zt} dt$$

for  $z \in \mathbb{C}$  and  $k \in \mathbb{N}_0$ . With integration by parts we get the recursive formula

$$I_k(z) = z^k e^z - k I_{k-1}(z) \tag{1}$$

and considering  $I_0(z) = e^z - 1$  we see (by induction over  $k$ ) that for every  $k \in \mathbb{N}_0$  there is a polynomial  $r_k(z) \in \mathbb{Z}[z]$  of degree  $k$  such that

$$I_k(z) = r_k(z)e^z - (-1)^k k!.$$

Now, we consider the polynomials

$$p_n(t) := t^n(1-t)^n/n! ,$$

where  $n$  is a positive integer. It is well known (and easily seen) that  $p_n^{(n)}(t)$  is a polynomial of degree  $n$  with integer coefficients  $p_{n,0}, \dots, p_{n,n}$ . Thus the integral

$$J_n(z) := z^{n+1} \int_0^1 p_n^{(n)}(t) e^{zt} dt = \sum_{k=0}^n p_{n,k} z^{n-k} I_k(z) \quad (2)$$

is the sum of integer multiples of the integrals  $z^n I_0(z), z^{n-1} I_1(z), \dots, I_n(z)$ . Hence there are two polynomials  $Q_n(z), R_n(z) \in \mathbb{Z}[z]$  of degree  $\leq n$  with

$$J_n(z) = Q_n(z) + R_n(z)e^z. \quad (3)$$

On the other hand, by repeatedly using integration by parts we can transform the integral  $J_n(z)$  into

$$J_n(z) = (-1)^n \frac{z^{2n+1}}{n!} \int_0^1 t^n(1-t)^n e^{zt} dt. \quad (4)$$

Since  $0 \leq t(1-t) \leq 1/4$  for all  $t \in [0, 1]$  we get

$$|J_n(z)| \leq |z| e^{\operatorname{Re}(z)} \frac{|z/2|^{2n}}{n!}. \quad (5)$$

Now suppose that  $z$  and  $e^z$  are Gaussian rationals, that is,  $z \in \mathbb{Q} + \mathbb{Q}\mathbf{i}$  and  $e^z \in \mathbb{Q} + \mathbb{Q}\mathbf{i}$ . Then  $z = (a + a'\mathbf{i})/b$  and  $e^z = (c + c'\mathbf{i})/d$  with  $a, a', c, c' \in \mathbb{Z}$  and  $b, d \in \mathbb{N}$ . From (5) we have  $db^n J_n(z) \rightarrow 0$  as  $n \rightarrow \infty$  and from (3) we see that

$$db^n J_n(z) = db^n(Q_n(z) + R_n(z)e^z) \in \mathbb{Z} + \mathbb{Z}\mathbf{i},$$

i.e., a Gaussian integer. From this we obtain the following

**Proposition.** *If  $J_n(z)$  does not eventually vanish then not both of  $z$  and  $e^z$  can be Gaussian rationals.*

If  $x \neq 0$  is a real number, then obviously  $J_n(x)$  is nonzero, since the integral in (4) is positive. Thus, if  $x \neq 0$  is rational, the proposition implies the irrationality of  $e^x$  and if  $y \neq 1$  is a positive rational number, the proposition (applied to  $z = \ln y$ ) shows the irrationality of  $\ln(y)$ . For  $z = \mathbf{i}\pi$  we have

$$\operatorname{Im}\left(\int_0^1 t^n(1-t)^n e^{zt} dt\right) = \int_0^1 t^n(1-t)^n \sin(\pi t) dt > 0 \quad (6)$$

and in particular  $J_n(\mathbf{i}\pi) \neq 0$ . From  $e^{\mathbf{i}\pi} = -1$  and the proposition we obtain the irrationality of  $\pi$ . Moreover, since  $\cos(s) = -\cos(\pi - s)$ , the real part of the integral

in (6) vanishes and we obtain from (4) that  $J_n(\mathbf{i}\pi)$  is real. Denoting the coefficients of  $(Q_n - R_n)(z) \in \mathbb{Z}[z]$  by  $c_{n,0}, \dots, c_{n,n}$  we have

$$0 \neq J_n(\mathbf{i}\pi) = (Q_n - R_n)(\mathbf{i}\pi) = \operatorname{Re}(Q_n - R_n)(\mathbf{i}\pi) = \sum_{v=0}^{\lfloor n/2 \rfloor} (-1)^v c_{n,2v} \pi^{2v}.$$

If we suppose that  $\pi^2 = a/b$  with positive integers  $a$  and  $b$  then  $b^n J_n(\mathbf{i}\pi)$  is a nonzero integer. Similarly as above, this contradicts (5) for large  $n$ .

**Remark.** (3) and (4) imply that  $-Q_n/R_n$  is the  $(n, n)$ -Padé approximant of  $e^z$  (cf. [1, p. 318]). By explicitly computing the approximation polynomials used by Iwamoto, one then sees that these actually equal  $Q_n - R_n$ . The same can be observed regarding Niven's polynomials and  $\operatorname{Re}(Q_n + R_n \mathbf{i})$  which correspond to the case  $z = \mathbf{i}\pi/2$ . Also, in the note [5] Nesterenko used the explicit form of the Padé approximants to obtain an estimate for  $J_n$  similar to (5) and with that the irrationality of  $\pi$  and  $e^x$  for  $x \in \mathbb{Q} \setminus \{0\}$ .

## References

- [1] Borwein, J.M., Borwein, P.B., *Pi and the AGM*, Wiley, New York (1987).
- [2] Butlewski, Z., *A Proof that  $e^m$  is irrational*, Colloq. Math. 1 (1948), 197–198.
- [3] Fel'dman, N.I., Nesterenko, Yu.V., *Transcendental numbers. Number Theory IV*, Encyclopaedia of Mathematical Sciences. 44. Springer, Berlin (1998).
- [4] Iwamoto, Y., *A proof that  $\pi^2$  is irrational*, J. Osaka Institute of Science and Technology 1 (1949), 147–148
- [5] Nesterenko, Yu.V., *A simple proof of the irrationality of  $\pi$* , Russian Journal of Mathematical Physics 13 (2006), 473.
- [6] Niven, I., *A simple proof that  $\pi$  is irrational*, Bull. Amer. Math. Soc. 53 (1947), 509.

Jürgen Müller and Tom Müller  
 Universität Trier  
 FB IV, Mathematik  
 D-54286 Trier, Germany  
 e-mails: jmueller@uni-trier.de  
 s0tomuel@uni-trier.de