

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 73 (2018)
Heft: 4

Artikel: Walking on rational numbers and a self-referential formula
Autor: Fortman, Margaret / Kupiec, Kevin / Rawlings, Marina
DOI: <https://doi.org/10.5169/seals-787382>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Walking on rational numbers and a self-referential formula

Margaret Fortman, Kevin Kupiec,
Marina Rawlings and Enrique Treviño

Margaret Fortman is a student at Lake Forest College studying mathematics and physics. She completed three summers of undergraduate research in mathematics, applied mathematics, and materials science at Lake Forest College, San Diego State University, and Northwestern University, respectively.

Kevin Kupiec received a Bachelor of Arts Degree in Mathematics from Lake Forest College with specializations in statistics and computer science. He has contributed to research programs at Harvard University, as well. Kevin is pursuing a career in software development. He is from Chicago, IL.

Marina Rawlings graduated from Lake Forest College in 2017 with a B.A. in Mathematics and Economics. She is currently in a rotational program for developing leadership with a top U.S. insurance corporation. She also has various artistic pursuits in her spare time.

Enrique Treviño is an Assistant Professor at Lake Forest College. He got his PhD from Dartmouth College in 2011 working in number theory with Carl Pomerance. When not doing mathematics, he enjoys playing fútbol with his students and running around the house with his 3 year old daughter Katya.

Eine in Basis vier notierte reelle Zahl kann als zweidimensionaler Gitterweg interpretiert werden: Man startet im Ursprung und betrachtet die Ziffern 0, 1, 2 und 3 je als Schritt nach rechts, oben, links und unten. Umgekehrt kann man jeden Gitterweg, der nicht mit einer Bewegung nach rechts beginnt, in entsprechender Weise als reelle Zahl auffassen. Insbesondere lassen sich beliebige Schriftzeichen, die man sich als Gitterpfade gezeichnet denkt, in reelle Zahlen verwandeln. Die Autoren der vorliegenden Arbeit geben eine Formel an, die als Input einen Satz aus gegebenen Schriftzeichen hat, und als Output eine rationale Zahl liefert, die genau diesen Satz ins Gitter schreibt. Zudem wird eine Variante von Tuppers Formel diskutiert, die mit Pixeln statt mit Pfaden arbeitet: Tuppers Formel kann in Abhängigkeit einer natürlichen Zahl jede Graphik in einem 106×17 -Display darstellen, inklusive die Formel selber.

1 Walking on numbers

In [1], Aragón Artacho, et al. describe the process of a walk on the plane using the digits of a number base 4.¹ Consider a number x written in base 4, $x = d_n d_{n-1} \dots d_0 . d_{-1} d_{-2} \dots$. We start at the origin in the Cartesian plane. If $d_n = 0$ we move a unit to the right, if it is 1, we move a unit upwards, if it is 2, we move a unit to the left, and if it is 3, we move downwards. We continue this process with $d_{n-1}, d_{n-2}, \dots, d_1, d_0, d_{-1}, \dots$, and this process creates a “walk.” For example, the number 419636198 is rewritten in base 4 as 121000302033212_4 . The walk would look like Figure 1.




Figure 1 Walk for the number 419636198, which is 121000302033212_4 .

In [1], they show several other walks, including walks on π and e using 100 billion digits. The one that inspired this paper is the following: Consider the rational number

$$\begin{aligned}
 Q = & 10490122716774994374866192805654486016175673584915608761668483808431443 \\
 & 58447252875551629247027759555570453715679313058783247729772021770818187 \\
 & 96590637365767487981422801328592027861019258140957135748704712290267465 \\
 & 1513128059541953997504202061380373822338959713391954 \\
 & / \\
 & 16122269626942909129404900662735492142298807557254685123533957184651913 \\
 & 53017348814314017504539969445479353012064383327267097007933052629203035 \\
 & 09209736004509554561365966493250783914647728401623856513742952945308961 \\
 & 2268152748875615658076162410788075184599421938774835.
 \end{aligned}$$

The walk on this number is Figure 2.²



Figure 2 Walk on Q .

In Section 2 we describe an algorithm that, given any sentence σ , it can find a rational number whose walk creates an image that spells out σ . We do something similar in Section 4, but instead of using walks as our starting point we use Tupper’s self-referential formula, which is described in Section 3.

¹They describe the process in any base, but for the purposes of this paper, we’ll focus on base 4.

²This figure comes from [1]. As the authors explain, the color in this walk indicates the path followed by the walk. It is shifted up the spectrum (red-orange-yellow-green-cyan-blue-purple-red).

and

$$r = \sum_{i=1}^k \frac{r_{\alpha_i}}{4^{\left(\sum_{j=1}^{i-1} (n_{\alpha_j} + w)\right)}} + \frac{2 \cdot 4^w \left(1 - \frac{1}{4^{(k-1)w}}\right)}{3 \cdot 4^{\left(\sum_{j=1}^k (n_{\alpha_j} + w)\right)}}. \tag{4}$$

Then a walk on r of length n spells out σ . Furthermore, a walk on $\frac{4^n}{4^n - 1}r$ of any length $m \geq n$ spells out σ .

Proof. The idea is to focus on the digits first. We have n_{α_1} digits to represent α_1 , then we include w zeroes to give space for the next letter. We follow this with n_{α_2} digits of the second letter, followed by w zeroes, and so on. When we “write” the last letter, we have used $n_{\alpha_1} + n_{\alpha_2} + \dots + n_{\alpha_k} + (k - 1)w$ digits. But the walk is $(k - 1)w$ steps to the right. To get back to the origin, we need to take $(k - 1)w$ steps to the left, i.e., we need $(k - 1)w$ 2’s in the digit expansion. Therefore we’ve used n digits where n is the same as in (3).

Now we want to find the rational that has this digit expansion. To account for the letter α_i in the desired position, we need to multiply it by 4^{-x} where x is the number of digits used so far. For α_1 , we’ve used 0, for α_2 we’ve used $n_{\alpha_1} + w$, for α_3 we’ve used $(n_{\alpha_1} + w) + (n_{\alpha_2} + w)$, and in general for α_i , we’ve used $\sum_{j=1}^{i-1} (n_{\alpha_j} + w)$. Finally, we have to take into account the final $(k - 1)w$ 2’s. To do this, we can think of $r_{\alpha_{k+1}} = 0.22 \dots 2_4$ and place it after all the digits so far, which have been $\sum_{i=1}^{k-1} (n_{\alpha_i} + w) + n_{\alpha_k}$. Therefore

$$\begin{aligned} r &= \sum_{i=1}^k \frac{r_{\alpha_i}}{4^{\left(\sum_{j=1}^{i-1} (n_{\alpha_j} + w)\right)}} + \frac{r_{\alpha_{k+1}}}{4^{\left(\sum_{i=1}^k (n_{\alpha_i} + w) - w\right)}} \\ &= \sum_{i=1}^k \frac{r_{\alpha_i}}{4^{\left(\sum_{j=1}^{i-1} (n_{\alpha_j} + w)\right)}} + \frac{4^w}{4^{\left(\sum_{i=1}^k (n_{\alpha_i} + w)\right)}} \left(\frac{2}{4} + \frac{2}{4^2} + \frac{2}{4^3} + \dots + \frac{2}{4^{(k-1)w}}\right). \end{aligned}$$

By completing the geometric series we can verify this matches (4). By construction, the walk for r with n steps spells out σ . Furthermore, by the same process as that in (2), we find the rational whose infinite walk spells out σ . □

Theorem 1 suggests how to build a program to find a rational number for any sentence. As an example, a certain rational

$$r_\sigma = \frac{3.47783 \dots \times 10^{3195}}{5.42542 \dots \times 10^{3195}}, \tag{5}$$

creates Figure 4.



Figure 4 The walk for r_σ as in (5) after 10000 steps.

3 Tupper's self-referential formula

In [4], Tupper introduced the formula⁴

$$\frac{1}{2} < \left\lfloor \text{mod} \left(\left\lfloor \frac{y}{17} \right\rfloor 2^{-17\lfloor x \rfloor - \text{mod}(\lfloor y \rfloor, 17)}, 2 \right) \right\rfloor. \tag{6}$$

This formula has the amazing property that if you graph the equation⁵ for $0 \leq x < 106$ and $k \leq y < k + 17$ for k as in (7)⁶ you get Figure 5.




Figure 5 Graph of (6) in the range $0 \leq x < 106$ and $k \leq y < k + 17$.

$$\begin{aligned} k = &4858450636189713423582095962494202044581400587983244549483093085061934704708809 \\ &9284506447698655243648499972470249151191104116057391774078569197543265718554420 \\ &5721044573588368182982375413963433822519945219165128434833290513119319995350241 \\ &3758765239264874613394906870130562295813219481113685339535565290850023875092856 \\ &8926945559742815463865107300491067230589335860525440966643512653493636439571255 \\ &6569593681518433485760526694016125126695142155053955451915378545752575659074054 \\ &0157929001765967965480064427829131488548259914721248506352686630476300. \end{aligned} \tag{7}$$

It turns out that the formula doesn't only graph itself, by considering different values of k , we can graph anything that can be represented by pixels in a 106×17 table. For example, a certain value

$$k_0 = 1.4452 \dots \times 10^{536} \tag{8}$$

gives the interval in which the graph looks like Figure 6.



Figure 6 Graph of (6) in the range $0 \leq x < 106$ and $k_0 \leq y < k_0 + 17$.

The main reason why we can build anything in a 106×17 grid is the following lemma:

Lemma 1. *Let $k = 17k'$ for a nonnegative integer $k' < 2^{106 \times 17}$. Suppose we write k' in binary as follows:*

$$k' = \sum_{m=0}^{105} \sum_{n=0}^{16} a_{17m+n} 2^{17m+n}. \tag{9}$$

⁴The formula was given as an example of a formula that graphing software had difficulties with, but Tupper's graphing software can handle.

⁵By this we mean that the point (x, y) is painted black if it satisfies the inequality and not painted if it doesn't.

⁶In [2] and many other places, the value of k is given differently because of the convention in computer science that positive y go downwards.

Then

$$\left\lfloor \text{mod} \left(\left\lfloor \frac{y}{17} \right\rfloor 2^{-17\lfloor x \rfloor - \text{mod}(\lfloor y \rfloor, 17)}, 2 \right) \right\rfloor = a_b, \tag{10}$$

for $b = 17\lfloor x \rfloor + \text{mod}(\lfloor y \rfloor, 17)$.

Therefore, the point (x, y) is painted whenever $a_b = 1$ and not painted when $a_b = 0$, i.e., it depends only on the binary expansion of k' .

Proof. Let $\lfloor x \rfloor = i$ and $\lfloor y \rfloor = j = 17k' + j'$ for some $0 \leq j' \leq 16$. Then $\lfloor \frac{y}{17} \rfloor = k'$, $j' = \text{mod}(\lfloor y \rfloor, 17)$, and $b = 17i + j'$. Now

$$\left\lfloor \frac{y}{17} \right\rfloor 2^{-17\lfloor x \rfloor - \text{mod}(\lfloor y \rfloor, 17)} = k' 2^{-17i - j'} = \sum_{m=0}^{105} \sum_{j=0}^{16} a_{17m+n} 2^{17m+n-17i-j'}.$$

When we consider mod 2, we can eliminate any term where the exponent of 2 is at least 1, i.e., we're left with exponents satisfying $17m + n - 17i - j' \leq 0$. When we take the floor, we exclude any of the small exponents because $1/2 + 1/4 + \dots + 1/2^c < 1$ for any finite c . Therefore the only exponent of 2 we allow is 0. Hence $17m + n - 17i - j' = 0$. This implies $n \equiv j' \pmod{17}$, but both n and j' are between 0 and 16, so $n = j'$, and then $m = i$, which is what we wanted to prove. \square

4 Writing using Tupper's self-referential formula

From Lemma 1 we can extrapolate an algorithm to find a k to build any picture in a 106×17 grid. Indeed, write a 1 on any unit square that is painted black and a 0 otherwise. Now starting at the square with bottom-left corner $(0, 0)$, read the digits from bottom to top on each column. This binary number (read from right to left) will be k' and so $k = 17k'$.

Problem: For a given sentence, find the integer k such that the graph of Tupper's formula looks like that sentence for $0 \leq x < 106$ and $k \leq y < k + 17$.

As in the walk example, the key is figuring out how to do a letter first. Let's demonstrate how to do the letter a . Consider Figure 7. We read the number as 11101 10101 11111. To transform it into a number that fits in the 106×17 grid, we need to fill in the necessary 0's, which is equivalent to multiplying numbers in the ℓ -column by $2^{17(\ell-1)}$. Therefore, we associate the letter "a" with the number

$$17 \left((1 + 2 + 4 + 16) + (1 + 4 + 16)2^{17} + (1 + 2 + 4 + 8 + 16)2^{34} \right). \tag{11}$$



Figure 7 Breaking down the letter "a" in binary.

We can now move a letter around the grid by multiplying it by 2^{17m+n} to place it where the bottom-left corner of the letter is (m, n) . If we create all letters with a height of at most 5 squares and width of at most 5 squares (the letters "m" and "w" need 5 squares, and the

rest need 3), we can then fit up to three rows of letters to spell a short sentence. Given a letter α , let $f(\alpha)$ be the number we associate with α with bottom-left corner on $(0, 0)$. We'll let $f_{\text{blank}} = 0$ and for letters with width 3, we'll multiply their numbers by 2^{17} ,⁷ to create a buffer between letters.

Theorem 2. *Given a sentence $\sigma = \alpha_1\alpha_2 \cdots \alpha_k$, where α_i represents a single letter or a blank space and $k \leq 63$, we use the following formula to figure out the value of k for the range where the plot of Tupper's formula is σ :*

$$\sum_{i=1}^{\min(21,k)} 2^{85(i-1)+12} f(\alpha_i) + \sum_{i=22}^{\min(42,k)} 2^{85(i-22)+6} f(\alpha_i) + \sum_{i=43}^k 2^{85(i-43)} f(\alpha_i). \quad (12)$$

Proof. Each letter fits in a block of width 5 and height 5. To move from one letter to the next (to the right), we need to multiply by $2^{17 \times 5} = 2^{85}$. This is where the 85's in the exponents come from. The reason we add 12 and 6 (depending on how many letters we have) is because the first row consists of numbers in the top "strip" ($k + 12 \leq y < y + 17$), so we have to multiply by 2^{12} to move upwards. The numbers in the middle strip ($k + 6 \leq y \leq y + 11$) need a shift of 2^6 , and the bottom row needs no translation. The formula follows. \square

As an example of finding a k for a particular phrase, Figure 8 is the plot of Tupper's formula for $0 \leq x < 106$ and $k_1 \leq y < k_1 + 17$ for a certain

$$k_1 = 6.20234 \dots \times 10^{461}. \quad (13)$$



Figure 8 Graph of (6) in the range $0 \leq x < 106$ and $k_1 \leq y < k_1 + 17$.

Appendix: Full decimal digit expansion of constants in the paper.

The value of δ in (1) is

$$\begin{aligned} \delta = & 6384779382043951036217348661253680515005885357484535471589654514956414794662 \\ & 721006368542597248986985323127416704519810815261318970154183 \\ & / \\ & 2325883917745942049757836185241614509931652354199417792900768637378045721962 \\ & 8733546438113622840434097944400691400517693873107252115668992. \end{aligned} \quad (14)$$

The value of k_0 in (8) is

$$\begin{aligned} k_0 = & 1445202489708975828479425373371945674812777822151507024797188139685490 \\ & 8873568298734888825132090576643817888323197692344001666776474924212512 \end{aligned}$$

⁷The only letters not multiplied by 2^{17} are "m" and "n".

$$\begin{aligned}
& 8995265907053708020473915320841631792025549005418004768657201699730466 \\
& 3833949016013743197155209961811452497819450190683595005106578043256408 \\
& 0119786755686314228025969420625409608166564241736740394638417077453742 \\
& 7319606443899923010379398938675025786929455234476319291860957618345432 \\
& 2480049217280333494198162067498544720381939397385138489604767597826733 \\
& 1343769705199458068186981933044636774047268864, \tag{15}
\end{aligned}$$

The value of k_1 in (13) is

$$\begin{aligned}
k_1 = & 6202342045523518696372190728132145377913289497819263812843155643364944 \\
& 0461939615515996102292719598762206682015817244444562918664906697777197 \\
& 5007949955351598702405129648571930754026169504789347614953307622064658 \\
& 7662203381308047340029024837030531000814297140117523848644113896733785 \\
& 5616409282734138890208876466646383764272086299397454808405688789312744 \\
& 7832949435883695715278636348898143061593729742606126050532003884145813 \\
& 574480854000747397523613796592272870866944. \tag{16}
\end{aligned}$$

The value of r_σ in (5) is

$$\begin{aligned}
r_\sigma = & 3477838176632809766274027652998361070109305198798526822993346685629729543816284212324168906789690775547148059136594 \\
& 5650708415322762044997606235997855677049782095397960796362753436599841518280673260542696144332265810919906062866086 \\
& 7300310028185786493083481099698081973220156381116333517929438873090176653898133176194667067667046684812642590682921 \\
& 7676064926655968063069943083237148802645189802820278834066659917243976811957017195714300536678576755193478646225748 \\
& 0029718138397905599265918142427252199419439333830496840583774337111547232356603947260968981693338103053145854635253 \\
& 6556487569357115640975974485570693172272764417938367108726707881765559137578753568579282574517818911608998412741042 \\
& 0227212620869607503190020705525648927347045787545010799076183740241404923921318243481736923524124663788869995658064 \\
& 9792376238237930407962450320389159643655424259838617670381363425821742408854318599877016505639048938102659648790392 \\
& 9595397634307377756861919915276546513631911373910771937588542354894963935837136025758958018813262204515835086096680 \\
& 5708164505804035890264351831753362652746839651784749459769618297620283942247979473521463646695449617575428923630172 \\
& 04835915387116752765669677607707093216794342308880360116044571717992965087557491510480694730625476830869629084918 \\
& 0330152435295444522317092256763450131932664980244104376549124516719713135304140238008592647538068871086358474426810 \\
& 3228839275062729870736963264331850092569189868395203739083590687341840855847621655276649131461408882329507967011133 \\
& 9673513989596387755006734087403631374900979733345825183520000390763148149610355789986385442991553413321157523278681 \\
& 4570776631231717821839613245063461887765374214685496293414305359730563242820842921560948381487459668207389426718341 \\
& 0208201125760638630177927567438294560420792809301856500683312377648424972104939732771020518033666376045372522152359 \\
& 6365323093742860258648736509758715341472162499794997150383376599167041645495565423248191931830700401690739885894446 \\
& 5925138828780782529332465419011534533168166732703212020724420330540765914808719296089874268534962295074250221700319 \\
& 4354384010924295515836192696117735836407353432915120743106575043357828096478475426584678135447571530015321875975666 \\
& 0288499555262323407672275144428184247080572013967367202580538705898188934186489884901842932385285873056974563406043 \\
& 2972003424454777927419274342586200406369883986949725285622098211132577303257888674921256677207569164691929862742553 \\
& 6434760361740959640882424911067801423858038854691012806455532141473569316256449253261282819184164210728465538046128 \\
& 8689316433179357074775456515481274301511980843961573141008058309365269614398459837204683190833300361309048467141298 \\
& 7077183501524879707368429696020266759574384790866466734735759122149901064852517874055671305612116400142877895886450 \\
& 1717055895264820300249217342710966276085484922766003610831279480622647998891816267884673999729550256532945744911484 \\
& 6457331356310576117617522162604119512965408010784378997607700863451724450862932076743527823981076515112164422001921 \\
& 6801892221050493956203183395973148259212386859598795634512915298898905336948973956044452272241532431935859131543491 \\
& 8841290611039546092076535887617550365638779065367323042885225828369392365302807084634122922 \\
& / \\
& 5425427555547183235387483138677443269370516110125718034418458072780400020660097464368420331167768059210326095280700 \\
& 7960754517818980320913237680771539405202704824855426057414759118003163672774854142023984371664371147816596771028104 \\
& 6441014848078122087947108444254899832081659029137741969600347998581332358020366378254707477828257293605011341272917 \\
& 7555747527995632047517011038746128120784347825211963664998369471837665640062502456285346080810630834125801925184520 \\
& 2625907527220125563925851141313331030922107272915406124790182288452890603674783060337942569320897590240195578623903 \\
& 9566157200553376651298625999377015571387110468470489015343875859756592220376633541514211837269398010309207318883562 \\
& 6661594155820595537918980228922993377013997290067594729152329146585615037257856776126270118053463580693713572941709
\end{aligned}$$

0234335508300132571024621930514890212320934438015747210297144734254112822232392753999420252062509544044913507097587
 9350699695390168912767463858081679059061108453821615559680371903283785451415371400938993626727592450435669695632669
 7254222479018247794025921399900915178303441159892932368904241515358090051156613040271864216857817861614058082054879
 7992961270430349541616758800746625539427565309875638214927919257745396009984319529064787174938772005587710967554787
 3134637767111202690659149688125290400377854774888701800589697431906307040676654222644000854136099592624341159355680
 4025055285850722385200868824064066076467668623572504966498893887720846592451499855656864232341936775024254014988573
 6479995630540608861215015614298371308536980538926684840420124076706834851544084593920349240191074869574272723680521
 7127047049869920301330050108161034628412359529481374507528437783717961628384774969135064191568538488550802164466946
 2991844172716763186432074166709520943957216137007438680496257872185964930482173472798553657122383518454992612482965
 8983430312485656075937606211310154685225690149806271811259712611252549777641482169337757296336319039098051110001043
 7499227486471556829577571187480183096203689144604478969275431212534506308782256889985344653287072554683775069061486
 3014073442952271232361355689066831264378429515802977509102274184384658882941102930680858230829056588956526453972913
 517424823594276356517394477107632725305886992825822461928518132640292288853044020007731096127030815289259831325582
 5255354353810146217297245345948544438608118775507870315659930862311032719187735199457377847675540579010031134555663
 448900883567256390076131806135532284916027030170720880175003277202061389610811502156139259718144929520269648838872
 0466980806627099676908203772701028988981196334425767237037978129102064210173720253085972283392852129043684301723824
 2526468727429343973246435674499354269034727279971334516277822171422775378729827488841283446462926242824176152611082
 7525355700871671055020654492230456283388123851578103617271263271404921176203093075586296285974629333519569598503406
 2164519084107133103865309775307272676431232643151623368126225771314148471748923659994901815532321569291863200955776
 1945338825314444239627509164817263099215974450756080971015744259843605203901279813501743110877702438536565400720937
 3657055236584191717148472650721961987021467849755631867227974194334896125257314027790401535.

Acknowledgement

We used Mathematica to create the walks and to plot Tupper's self-referential formula in different ranges. We would like to thank Scott T. Chapman for his suggestions that improved the exposition of this paper, and to Pete Winkler for comments on the paper. We would also like to thank the anonymous referee for his thoughtful comments.

References

- [1] Francisco J. Aragón Artacho, David H. Bailey, Jonathan M. Borwein, and Peter B. Borwein, *Walking on real numbers*, Math. Intelligencer **35** (2013), no. 1, 42–60.
- [2] B. Haran and M. Parker, *The 'everything' formula – numberphile*, https://www.youtube.com/watch?v=_s5RFgd59ao, Accessed: 2017-05-23.
- [3] Wolfram Research, Inc., *Mathematica, Version 10.0*, Champaign, IL, 2017.
- [4] Jeff Tupper, *Reliable two-dimensional graphing methods for mathematical formulae with two free variables*, Proceedings of the 28th Annual Conference on Computer Graphics and Interactive Techniques (New York, NY, USA), SIGGRAPH '01, ACM, 2001, pp. 77–86.

Margaret Fortman
 Lake Forest College student
 e-mail:
 fortmanma@mx.lakeforest.edu

Marina Rawlings
 Lake Forest College 17'
 e-mail:
 rawlingsmn@mx.lakeforest.edu

Kevin Kupiec
 Lake Forest College 17'
 e-mail:
 kupieck@mx.lakeforest.edu

Enrique Treviño
 Department of Mathematics and
 Computer Science
 at Lake Forest College
 e-mail:
 trevino@lakeforest.edu