Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 75 (2020)
Heft: 3

Artikel: Mixtilinear radii and Finsler-Hadwiger inequality
Autor: Lukarevski, Martin / Wanner, Gerhard
DOI: https://doi.org/10.5169/seals-880889

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 17.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Elem. Math. 75 (2020) 121 – 124
0013-6018/20/030121-4
DOI 10.4171/EM/412

© Swiss Mathematical Society, 2020

Elemente der Mathematik

Short note

Mixtilinear radii and Finsler–Hadwiger inequality

Martin Lukarevski and Gerhard Wanner

Abstract. In this short note we derive few relations for the mixtilinear radii of a triangle and then as a curious application give an interesting proof of the celebrated Finsler–Hadwiger inequality.

The *mixtilinear incircle* is a circle tangent to two sides of a triangle and internally to the circumcircle of the triangle, [1]. Another, more suggestive name is *inarc circle*, [6]. We de-



Figure 1 Mixtilinear incircle and proof of $\rho_A = r\sec^2\frac{A}{2}$.

note by ρ_A, ρ_B, ρ_C the radii of the mixtilinear incircles of a triangle ABC with inradius r, circumradius R and semiperimeter s. It is known, [11], that the center of the mixtilinear incircle is the orthogonal projection on the bisector of the point on the side which is itself the orthogonal projection of the incenter. See Figure 1. Hence one readily gets that

$$\rho_A = r\sec^2\frac{A}{2}.$$

Theorem 1. *It holds for the radii of the mixtilinear incircles*

$$\rho_A + \rho_B + \rho_C = r\left(1 + \left(\frac{4R + r}{s}\right)^2\right), \tag{1}$$

$$\rho_A\rho_B + \rho_B\rho_C + \rho_C\rho_A = \frac{8Rr^2(4R+r)}{s^2}, \tag{2}$$

$$\rho_A\rho_B\rho_C = \frac{16R^2r^3}{s^2}, \tag{3}$$

$$\frac{1}{\rho_A} + \frac{1}{\rho_B} + \frac{1}{\rho_C} = \frac{4R+r}{2Rr}. \tag{4}$$

Proof. The cosines $\cos A$, $\cos B$, $\cos C$ are the roots of the polynomial [10, p. 56]

$$4R^2x^3 - 4R(R+r)x^2 + (s^2+r^2-4R^2)x + (2R+r)^2 - s^2 = 0.$$

Putting $x = 2y - 1$ and rearranging, we get that the polynomial

$$16R^2y^3 - 8R(4R+r)y^2 + (s^2 + (4R+r)^2)y - s^2 = 0 \tag{5}$$

has $\cos^2\frac{A}{2}$, $\cos^2\frac{B}{2}$, $\cos^2\frac{C}{2}$ as its roots, and since $\cos^2\frac{A}{2} = r/\rho_A$, we obtain that the mixtilinear radii ρ_A, ρ_B, ρ_C are roots of the polynomial

$$s^2t^3 - r(s^2 + (4R+r)^2)t^2 + 8Rr^2(4R+r)t - 16R^2r^3 = 0.$$

By Vieta's formulas the identities (1), (2) and (3) follow. The last identity (4) follows from

$$\frac{1}{\rho_A} + \frac{1}{\rho_B} + \frac{1}{\rho_C} = \frac{1}{r}\left(\cos^2\frac{A}{2} + \cos^2\frac{B}{2} + \cos^2\frac{C}{2}\right)$$

and the polynomial (5). □

We note that one can use the identity (1) to give a nice linear bound for the sum of the mixtilinear radii only in terms of R and r

$$4r \leqslant \rho_A + \rho_B + \rho_C \leqslant R + 2r, \tag{6}$$

which is sharper than $4r \leqslant \rho_A + \rho_B + \rho_C \leqslant (5R+6r)/4$, given in [12]. The LHS of (6) follows easily from the convexity of $f(x) = \sec^2\frac{x}{2}$ and Jensen's inequality. For the RHS we shall make use of the Gerretsen inequality $s^2 \geqslant r(16R-5r)$, [4, 5]. By (1), the RHS of (6)

$$r\left(1 + \left(\frac{4R+r}{s}\right)^2\right) \leqslant R + 2r$$

is equivalent to $s^2(R+r) \geqslant r(4R+r)^2$. By Gerretsen's inequality and Euler's inequality $R \geqslant 2r$,

$$\begin{aligned} s^2(R+r) &\geqslant r(16R-5r)(R+r) \\ &= r\left[(4R+r)^2 + 3(R-2r)\right] \\ &\geqslant r(4R+r)^2, \end{aligned}$$

which proves (6), with equality for the equilateral triangle.

For the end of this note, we show how the relations for the mixtilinear radii can be used to give an interesting proof of the celebrated Finsler–Hadwiger inequality from 1937 [3, 13]

$$a^2 + b^2 + c^2 \geqslant 4\sqrt{3}\mathcal{A} + (a-b)^2 + (b-c)^2 + (c-a)^2, \tag{7}$$

where $\mathcal{A}$ is the area of ABC. It can be rewritten as

$$2\sqrt{3}\mathcal{A} \leqslant a(s-a)+b(s-b)+c(s-c) = 2s^2-a^2-b^2-c^2.$$

By the identities $a^2+b^2+c^2 = 2(s^2-4Rr-r^2)$ and $\mathcal{A} = sr$, known since Euler's times (see, e.g., [2, §11, §19]), the Finsler–Hadwiger inequality is equivalent to

$$s \leqslant (4R+r)/\sqrt{3}. \tag{8}$$

We can apply the AM-HM inequality to (1) and (4) to obtain

$$(\rho_A+\rho_B+\rho_C)\left(\frac{1}{\rho_A}+\frac{1}{\rho_B}+\frac{1}{\rho_C}\right) \geqslant 9,$$

which after substituting and rearranging is

$$s^2 \leqslant \frac{(4R+r)^3}{14R-r}.$$

It holds $3(4R+r) \leqslant 14R-r$ by Euler's inequality. Hence

$$s^2 \leqslant \frac{(4R+r)^3}{14R-r} \leqslant \frac{(4R+r)^2}{3},$$

and that is the Finsler–Hadwiger inequality (8). Equality holds for $\rho_A = \rho_B = \rho_C$, that is $\sec^2\frac{A}{2} = \sec^2\frac{B}{2} = \sec^2\frac{C}{2}$, and $R = 2r$, which implies that the triangle is equilateral.

The Hadwiger–Finsler inequality (7) is actually equivalent [7] to the seemingly weaker Weitzenböck inequality

$$a^2+b^2+c^2 \geqslant 4\sqrt{3}\mathcal{A}.$$

One sharpening of (7) is

$$a^2+b^2+c^2 \geqslant 4\mathcal{A}\sqrt{3+\frac{R-2r}{R}}+(a-b)^2+(b-c)^2+(c-a)^2,$$

which is equivalent to Kooi's inequality [8, 9].

References

[1] L. Bankoff, A Mixtilinear Adventure, *Crux Math.* **9** (1983), 2–7.

[2] L. Euler, Solutio facilis problematum quorundam geometricorum difficillimorum [E325], Novi Commentarii Academiae Scientiarum Petropolitanae 11, 1765 (1767) 103–123; reprinted in Opera Omnia, series prima, vol. 26, 139–157, Orell Füssli, Zürich 1953.

[3] P. Finsler, H. Hadwiger, Einige Relationen im Dreieck, *Commentarii Mathematici Helvetici*, **10** (1937), no.1, 316–326.

[4] J.C.H. Gerretsen, *Ongelijkheden in de driehoek.* Nieuw Tijdschr **41**, 1–7 (1953).

[5] M. Lukarevski, *An alternate proof of Gerretsen's inequalities*, Elem. Math. **72**, (2017), 2–8.

[6] M. Lukarevski, An inequality arising from the inarc centres of a triangle, *Math. Gaz.* **103** (November 2019) pp. 538–541.

[7] M. Lukarevski, The excentral triangle and a curious application to inequalities, *Math. Gaz.* **102** (November 2018) pp. 531–533.

[8] M. Lukarevski, D.S. Marinescu, A refinement of the Kooi's inequality, Mittenpunkt and applications, *J. Inequal. Appl.* **13** (3), (2019) pp. 827–832.

[9] M. Lukarevski, A Simple Proof of Kooi's Inequality, *Math. Mag.* **93** (3), (2020), p. 225.

[10] D.S. Mitrinović, J. Pečarić, V. Volenec, *Recent Advances in Geometric Inequalities*. Kluwer Academic Publishers. Dordrecht (1989).

[11] A. Ostermann, G. Wanner, *A dynamic proof of Thebault's theorem*, Elem. Math. **65**, (2010), 12–16.

[12] Problem 11912, (proposed by Pal Peter Delyay, solution by Tamas Wiandt) *Amer. Math. Monthly* **123** (2016), 505 and **125** (2018), 281–283.

[13] G. Wanner, *Gerretsen and the Finsler–Hadwiger inequality*, Elem. Math. **72**, (2017), 173–174.

Martin Lukarevski
Department of Mathematics and Statistics
University "Goce Delcev"
Stip, Macedonia
email: `martin.lukarevski@ugd.edu.mk`

Gerhard Wanner
Univ. Geneva
email: `Gerhard.Wanner@unige.ch`