

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 75 (2020)
Heft: 3

Artikel: A proof of the Schwarz theorem on mixed partial derivatives via elementary approximation theory
Autor: Camargo, André Pierro de
DOI: <https://doi.org/10.5169/seals-880890>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 04.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Short note **A proof of the Schwarz theorem on
mixed partial derivatives via
elementary approximation theory**

André Pierro de Camargo

Abstract. We show that the equality of the mixed second-order partial derivatives holds for functions that can be well approximated by linear combinations of functions whose variables separate and that this is the case for functions satisfying the hypothesis of Schwarz' theorem.

1 Introduction

A fundamental theorem of multivariate Calculus states that if a function $f(x, y)$ has partial derivatives f_x and f_y , and mixed derivatives f_{xy} and f_{yx} in a neighborhood Ω of a point (x_0, y_0) and if f_{xy} and f_{yx} are continuous at (x_0, y_0) , then¹

$$f_{xy}(x_0, y_0) = f_{yx}(x_0, y_0). \quad (1)$$

According to [5], many outstanding mathematicians, among them Euler, Lagrange and Cauchy, offered proofs of (1), but none of them without a fault. In fact, in 1867, Lindelöf published an article criticizing all those proofs in detail. The first satisfactory proof of (1) appeared only six years later in an article by Schwarz.

All proofs of (1) we found in a collection of Calculus textbooks [1, 2, 7, 8, 9, 10, 11] mainly reproduce Schwarz' original proof [12], which proceeds by showing that the function

$$\phi(h_x, h_y) = \frac{\frac{f(x_0 + h_x, y_0 + h_y) - f(x_0, y_0 + h_y)}{h_x} - \frac{f(x_0 + h_x, y_0) - f(x_0, y_0)}{h_x}}{h_y} \quad (2)$$

has a limit as $(h_x, h_y) \rightarrow (0, 0)$ and that this limit is equal to both $f_{xy}(0, 0)$ and $f_{yx}(0, 0)$. The reader can also find lightly different proofs in [3] and [4]; and [5] also points a proof of (1) based on Lebesgue integration.

What we found curious about this theorem is that none of the proofs presented in the references explores the obvious fact that (1) holds whenever f is a polynomial in the

¹A few refinements can be found in the literature [6], but we shall not discuss it here.

variables x and y or, more in general, if f is of the form

$$\sum_{i=1}^n u_i(x)v_i(y), \quad (3)$$

for some differentiable functions $u_1, u_2, \dots, u_n, v_1, v_2, \dots, v_n$.

Actually this hypothesis can be somewhat relaxed and (1) also holds for functions f that can be well approximated by functions g of the form (3) in the sense that

$$\lim_{(x,y) \rightarrow (x_0,y_0)} \frac{f(x,y) - g(x,y)}{(x-x_0)(y-y_0)} = 0. \quad (4)$$

This is true because

Lemma 1. *If f and g satisfy (4) and have partial derivatives and mixed partial derivatives of second order in Ω , then both first partial derivatives and mixed partial derivatives of second order of f and g coincide at the point (x_0, y_0) .*

In this note we show that, for each function f satisfying the hypothesis of Schwarz' theorem, there is a function g of the form (3) such that Lemma 1 holds for f and g and this proves (1).

Remark 1. We emphasize that, although a little bit longer than the usual argument for proving Schwarz' theorem, our strategy is based on the largely employed technique of *reducing the problem to a simpler one* (just as in the case of the analysis of local extrema by the analysis of quadratic approximations). Therefore, our reasoning is totally coherent and it is somewhat surprising that no one explored it before (at least according to our records).

Proof of (1)

To simplify the notation, let us assume that $(x_0, y_0) = (0, 0)$. We start with the initial approximation $g_1(x, y) := f(0, y) + f_x(0, y)x$ obtained by means of the partial derivative f_x :

$$f(x, y) = f(0, y) + f_x(0, y)x + R(x, y), \quad (5)$$

with $\lim_{x \rightarrow 0} \frac{R(x, y)}{x} = 0$ for every fixed y . We want to check whether f and g_1 satisfy the hypothesis of Lemma 1. In this vein, we note that (5) is not sufficient for our purpose and we must also analyze how

$$R(x, y) = f(x, y) - f(0, y) - f_x(0, y)x$$

varies with respect to y .

This expression shows that R is differentiable with respect to y and we have

$$R_y(x, y) = f_y(x, y) - f_y(0, y) - f_{xy}(0, y)x,$$

or, by the mean value theorem,

$$R_y(x, y) = [f_{yx}(\tau(x, y), y) - f_{xy}(0, y)] x,$$

with $|\tau(x, y)| < |x|$. Once again by the mean value theorem, we obtain

$$\begin{aligned} R(x, y) &= R(x, 0) + R_y(x, \zeta(x, y)) y \\ &= R(x, 0) + [f_{yx}(\tau(x, \zeta(x, y)), \zeta(x, y)) - f_{xy}(0, \zeta(x, y))] x y \\ &= R(x, 0) + [f_{yx}(0, 0) - f_{xy}(0, 0)] x y \\ &\quad + [f_{yx}(\tau(x, \zeta(x, y)), \zeta(x, y)) - f_{yx}(0, 0)] x y \\ &\quad + [f_{xy}(0, 0) - f_{xy}(0, \zeta(x, y))] x y, \end{aligned} \tag{6}$$

with $|\zeta(x, y)| < |y|$.

The expression above shows that f and g_1 do not satisfy the hypothesis of Lemma 1 in general, but (5) and (6) and the continuity of f_{xy} and f_{yx} at $(0, 0)$ shows that Lemma 1 holds for f and

$$\begin{aligned} g(x, y) &= f(0, y) + f_x(0, y) x + R(x, 0) + [f_{yx}(0, 0) - f_{xy}(0, 0)] x y \\ &= f(0, y) + f_x(0, y) x + f(x, 0) - f(0, 0) - f_x(0, 0) x \\ &\quad + [f_{yx}(0, 0) - f_{xy}(0, 0)] x y. \end{aligned}$$

In fact, we have

$$\begin{aligned} \lim_{(x,y) \rightarrow (0,0)} \frac{f(x,y) - g(x,y)}{xy} &= \lim_{(x,y) \rightarrow (0,0)} [f_{yx}(\tau(x, \zeta(x, y)), \zeta(x, y)) - f_{yx}(0, 0)] \\ &\quad + \lim_{(x,y) \rightarrow (0,0)} [f_{xy}(0, 0) - f_{xy}(0, \zeta(x, y))] \\ &= 0. \end{aligned}$$

In addition, note that g is of the form (3) for $n = 4$ and

$$\begin{aligned} v_1(y) &= f(0, y); & u_1(x) &= 1; \\ v_2(y) &= f_x(0, y); & u_2(x) &= x; \\ v_3(y) &= 1; & u_3(x) &= f(x, 0) - f(0, 0) - f_x(0, 0) x; \\ v_4(y) &= y; & u_4(x) &= [f_{yx}(0, 0) - f_{xy}(0, 0)] x. \end{aligned}$$

Because all these functions are differentiable (some of them by the assumptions on f and other simply because they are polynomials), the proof is complete.

Remark 2. We emphasize that, while $[f_{yx}(0, 0) - f_{xy}(0, 0)]$ appears explicitly in the definition of g , our argument does not depend on any previous knowledge about its value.

Proof of Lemma 1

Again, assume that $(x_0, y_0) = (0, 0)$. Let $h(x, y) = f(x, y) - g(x, y)$. By assumption, given any positive number ϵ there is a number $\delta > 0$ such that

$$|h(x, y)| \leq \epsilon |x| |y|, \tag{7}$$

for $0 < |x| < \delta$ and $0 < |y| < \delta$. Hence, we obtain

$$h(x, 0) = 0 \text{ for } |x| < \delta \text{ and } h(0, y) = 0 \text{ for } |y| < \delta.$$

This is true because $h(x, \cdot)$ and $h(\cdot, y)$ are differentiable (and therefore continuous) for every x and y . In particular,

$$h_x(0, 0) = 0. \quad (8)$$

Moreover, for $0 < |x| < \delta$ and $0 < |y| < \delta$ we have

$$\left| \frac{h(x, y) - h(0, y)}{x} \right| = \left| \frac{h(x, y)}{x} \right| \leq \epsilon |y|$$

and this shows that $|h_x(0, y)| \leq \epsilon |y|$ for $0 < |y| < \delta$. Combining this inequality with (8), we obtain

$$\left| \frac{h_x(0, y) - h_x(0, 0)}{y} \right| = \left| \frac{h_x(0, y)}{y} \right| \leq \epsilon$$

for $0 < |y| < \delta$ and, therefore,

$$|h_{xy}(0, 0)| \leq \epsilon.$$

Because ϵ can be chosen arbitrarily small, we have that $h_{xy}(0, 0) = 0$.

By symmetry, we also have that $h_{yx}(0, 0) = 0$.

References

- [1] R. Bartle, *The Elements of Real Analysis*, 2nd ed., John Wiley & Sons, New York, USA, 1964.
- [2] R. Courant, *Differential and Integral Calculus*, Black & Son, Glasgow, UK, 1961.
- [3] H.Y. Chen, *A theorem on the difference quotient*, Tohoku Math. Journal, 42 1936, pp. 86–89.
- [4] T. Drammatico, *A new proof of the equality of mixed second partial derivatives*, Amer. Math. Monthly, 122 (6) 2015, pp. 602–603.
- [5] T.J. Higgins, *A note on the history of mixed partial derivatives*, Scripta. Math., 7, 1940, pp. 59–62.
- [6] E. Minguzzi, *The equality of mixed partial derivatives under weak differentiability conditions*, Real. Anal. Exchange, 41 (1) 2015, pp. 81–98.
- [7] N. Piskounov, *Differential and Integral Calculus*, Mir, Moscow, RUS 1996.
- [8] W. Rudin, *Principles of Mathematical Analysis*, MacGraw-Hill 3th ed., New York, USA, 1976.
- [9] J. Stewart, *Calculus*, Brooks/Cole 4th ed., Belmont, CA, 2012.
- [10] E. Swokowski, *Calculus With Analytic Geometry*, 2nd ed., Prindle, Weber & Schmidt, Boston, USA, 1979.
- [11] G. B. Thomas, *Calculus*, 13th ed. Pearson, Boston, USA, 2014.
- [12] D. H. Trahan, *The mixed partial derivatives and the double derivative*, Amer. Math. Monthly, 76 (1) 1969, pp. 76–77.

André Pierro de Camargo

Centro de Matemática, Computação e Cognição

Universidade Federal do ABC – UFABC, BRA

e-mail: andrecamargo.math@gmail.com