

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 75 (2020)
Heft: 4

Artikel: The irrationality measure of π as seen through the eyes of $\cos(n)$
Autor: Chen, Sully F. / Pearse, Erin P. J.
DOI: <https://doi.org/10.5169/seals-880892>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 02.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The irrationality measure of π as seen through the eyes of $\cos(n)$

Sully F. Chen and Erin P.J. Pearse

Sully Chen is an undergraduate student who previously attended Cal Poly San Luis Obispo and is currently attending the University of Southern California studying biochemistry. Though he hopes to pursue a career in medicine, he has a strong passion for mathematics and hopes to continue his mathematical studies alongside his current endeavors.

Erin Pearse is an Associate Professor of Mathematics at Cal Poly San Luis Obispo. He has held postdoctoral positions at Cornell University, University of Iowa, and University of Oklahoma, after graduating from UC Riverside. This article began as a student project.

1 Introduction

1.1 Irrationality measure

An irrational number α is considered to be well approximated by rational numbers iff for any $\nu \in \mathbb{N}$, there are infinitely many choices of rational numbers p/q that satisfy

$$|\alpha - (p/q)| < 1/q^\nu, \quad \text{for some } \nu \in \mathbb{N}. \quad (1)$$

Für natürliche Zahlen $n > 0$ liegen die Werte $\cos n$ im offenen Intervall $(-1, 1)$. Man könnte daher vermuten, dass die Zahlen der Folge $(\cos n)^n$ mit wachsendem n gegen 0 streben. Andererseits liegen die Zahlen $\cos n$ dicht in $[-1, 1]$, und es könnte daher sein, dass gewisse Werte $\cos n$ so nahe bei 1 oder -1 liegen, dass dies auch noch für die Potenz $(\cos n)^n$ zutrifft. Tatsächlich ist letzteres der Fall. Mehr noch: Die Autoren der Arbeit zeigen, dass es ein ganzes Spektrum von Teilfolgen gibt, die sich aus der Umgebung von 0 lösen. Diese Teilfolgen hängen zusammen mit den Nennern, die in der diophantischen Approximation von π erscheinen und geben Einblick in das Irrationalitätsmass von π . Das Irrationalitätsmass einer Zahl beschreibt, wie effizient sie durch rationale Zahlen approximiert werden kann. Man weiss, dass das Irrationalitätsmass von π (und jeder anderen transzendenten Zahl) grösser oder gleich 2 ist, der genaue Wert ist jedoch unbekannt. Die numerischen Indizien in dieser Arbeit deuten nun darauf hin, dass das Irrationalitätsmass von π genau 2 ist.

The idea is that if the set

$$U(\alpha, \nu) = \left\{ \frac{p}{q} \in \mathbb{Q} : 0 < \left| \alpha - \frac{p}{q} \right| < \frac{1}{q^\nu} \right\} \quad (2)$$

is infinite for a fixed ν , then $U(\alpha, \nu)$ contains rationals with arbitrarily large denominators and so one can find a sequence of rationals $\frac{p_n}{q_n} \in U(\alpha, \nu)$ which converge to α . One could say that this sequence converges to α “at rate ν ”. When this can be accomplished for any $\nu \in \mathbb{N}$, the number α can be approximated by sequences with arbitrarily high rates of convergence. Such numbers are called *Liouville numbers* after Joseph Liouville who proved that all such numbers are transcendental, thereby giving the first proof of the existence of transcendental numbers [5]. See Appendix A for a proof of the transcendentality of Liouville numbers.

When the number α is not well approximated by a sequence of rationals, one can grade “how approximable” α is in terms of the optimal exponent for which an approximating sequence of rationals may be found, and this leads to the notion of irrationality measure (also variously called the Liouville–Roth irrationality measure, irrationality exponent, approximation exponent, or Liouville–Roth constant); cf. [3, 9].

Definition 1.1 (Irrationality Measure). The *irrationality measure* of $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ is

$$\mu(\alpha) = \inf\{\nu : |U(\alpha, \nu)| < \infty\}.$$

This means, for example, that except possibly for at most finitely many “lucky” choices of $\frac{p}{q}$, every rational approximation $\frac{p}{q}$ that gives $n + 1$ correct decimal digits of α must satisfy

$$\frac{1}{10^n} \geq \left| \alpha - \frac{p}{q} \right| \geq \frac{1}{q^{\mu(\alpha) + \varepsilon}}$$

for any $\varepsilon > 0$.

Example 1.2. Liouville numbers are precisely those numbers α with $\mu(\alpha) = \infty$; cf. [3]. The most well-known example is Liouville’s Constant [10], defined as:

$$\alpha = \sum_{k=1}^{\infty} 10^{-k!} = 0.110001 \dots$$

One can see by inspection that the partial sums in the 10-ary expansion of α (more commonly called the *decimal expansion*) of α approach the true value extremely quickly. If s_n denotes the n th partial sum, we have the first few rational approximations and their corresponding error bounds

n	1	2	3	4
s_n	0.1	0.11	0.110001	0.11000100...001
$ \alpha - s_n \leq$	2×10^{-2}	2×10^{-6}	2×10^{-24}	2×10^{-120}

Observe that the number of decimal places for which s_n agrees with α increases much faster than n . In terms of Definition 1.1, we have found that $U(\alpha, 2)$ contains infinitely many points.

In contrast, for $\sqrt{2}$ (whose irrationality measure is equal to 2), no such patterns exist in the 10-ary expansion, or indeed, in the q -ary expansion of $\sqrt{2}$ for any $q \in \mathbb{Q}^+$.

It is clear that there exists a deep underlying meaning and complexity to the irrationality measure of a number, and thus any efforts towards determining the exact values of the irrationality measures of different numbers would yield considerable insight.

It can be shown that $\mu(\alpha) = 1$ for any rational number $\alpha \in \mathbb{Q}$, and that $\mu(\alpha) = 2$, for every algebraic number α of order greater than 1, from Roth's theorem on arithmetic progressions; cf. [4].¹ However, transcendental numbers may have any irrationality measure greater than or equal to 2. There is no known method to compute or determine the exact value of the irrationality measure of any transcendental number, or even estimate the value. Upper bounds have been proven for many interesting transcendental constants, for example, $\mu(\pi) \leq 7.6063$ [7] and $\mu(\ln 3) \leq 5.125$ [6]. Proofs of exact values are rare, and one such gem is e , for which it is known that $\mu(e) = 2$ [3]. This is perhaps not surprising in light of the fact that almost all numbers have irrationality measure 2.

An exact value for the irrationality measure of the most famous of mathematical constants, π , has evaded us thus far, though it is widely believed that $\mu(\pi) = 2$. Using Corollary 1.4, we provide further numerical evidence that $\mu(\pi) = 2$ in §4.

Our main result concerns the following set:

$$\Phi = \{\gamma \geq 0 : \limsup\{|\cos k|^{k^\gamma}\}_{k \in \mathbb{N}} = 1\}. \quad (3)$$

Theorem 1.3. *For Φ as defined in (3), we have $\sup \Phi = 2\mu(\pi) - 2$.*

Corollary 1.4. *If $\sup \Phi = 2$, then $\mu(\pi) = 2$.*

Remark 1.5 (The relation of $\mu(\pi)$ to the end behavior of special sequences). The value of $\mu(\pi)$ is closely tied to the behavior and convergence of special sequences of transcendental numbers. For example, Alekseyev proved that the convergence of the Flint Hills series directly implies $\mu(\pi) < 2.5$, in [1].

The investigation in the present paper began with a question raised by a student regarding the behavior of the sequence $a_n = \cos(n)^n$. Since $|\cos n| < 1$, one might expect that raising this number to a large power would cause $a_n \rightarrow 0$. However, preliminary numerical investigations revealed that this is not the case. Indeed, while most values of this sequence are extremely close to 0, certain subsequences form oscillations with amplitude 1. The top of Figure 1 shows the first million terms of the sequence, and the bottom shows a magnification of just the first 100,000 terms, which reveals several rapidly decaying subsequences. In this note, we examine the sequence $a_n = \cos(n)^{n^\gamma}$, for $\gamma \geq 0$, and show that it is closely related to the value of $\mu(\pi)$.

¹This is a deep result, and Roth's work in this area earned him the 1958 Fields Medal.




Figure 1 The top plot shows the first 20,000 terms of $a_n = \cos(n)^n$; the bottom figure shows the first 1,000,000. Two subsequences, even and odd k are plotted. Note: each peak actually corresponds to a different subsequence, an arithmetic progression of the form $n_k = 355k + 22j$; see §5 for details.

2 Proof of the main result

Definition 2.1. We define a function $Q(n) : \mathbb{Z} \mapsto \mathbb{Z}$ such that for any integer n , $Q(n)$ maps to the integer minimizing $|n - Q(n)\pi|$. In other words, if $[x]$ denotes the nearest integer to x ,² then

$$Q(n) = \left[\frac{n}{\pi} \right] = \operatorname{argmin}\{|n - m\pi| : m \in \mathbb{Z}\}. \quad (4)$$

Outline of the proof of Theorem 1.3

We prove Theorem 1.3 by establishing

$$\limsup\{|\cos k|^{k^\gamma}\}_{k \in \mathbb{N}} = \begin{cases} 1, & \gamma < 2\mu(\pi) - 2, \\ 0, & \gamma > 2\mu(\pi) - 2. \end{cases} \quad (5a)$$

$$(5b)$$

The proof will require a couple of technical lemmas, which we state and prove before proceeding. From Lemma 2.2, it follows that $Q(n) = \mathcal{O}(n)$, $n \rightarrow \infty$, and this is the observation used to show (5a).

For (5b), we establish a bound of the form

$$|\cos p|^{p^\gamma} \leq \left(1 - \frac{1}{\phi_0(p)}\right)^{\phi_1(p)}$$

for

$$\phi_0(p) = \frac{24Q(p)^{4(\mu(\pi)-1+\lambda)}}{12Q(p)^{2(\mu(\pi)-1+\lambda)} - 1} \quad \text{and} \quad \phi_1(p) = p^\gamma,$$

and then apply Lemma 2.3.

²Note that $[x]$ is *not* the same as the floor or ceiling function.

Lemma 2.2. $\lim_{n \rightarrow \infty} \frac{n}{Q(n)} = \pi$.

Proof. This is immediate from Definition 2.1 by Diophantine approximation. \square

Lemma 2.3. Let $\phi_0(x)$ and $\phi_1(x)$ be \mathbb{R} -valued functions satisfying

$$\lim_{x \rightarrow \infty} \phi_0(x) = \infty \quad \text{and} \quad \lim_{x \rightarrow \infty} \phi_1(x) = \infty.$$

If $\phi_0(x) \neq 0$ for all $x \in \mathbb{R}$ and $\phi_0 = o(\phi_1)$, as $x \rightarrow \infty$, then

$$\lim_{x \rightarrow \infty} \left(1 - \frac{1}{\phi_0(x)}\right)^{\phi_1(x)} = 0. \quad (6)$$

Proof. Observe that $r(r-1)(r-2)\dots(r-n+1) \leq r^n$, so comparing the Taylor series for $(1+x)^r$ and e^{rx} gives

$$(1+x)^r \leq e^{rx}.$$

Now making the substitutions $x = -\frac{1}{\phi_0(x)}$ and $r = \phi_1(x)$, we obtain

$$\left(1 - \frac{1}{\phi_0(x)}\right)^{\phi_1(x)} \leq \exp\left(-\frac{\phi_1(x)}{\phi_0(x)}\right),$$

and the right side tends to 0 as $x \rightarrow \infty$ because $\phi_0 = o(\phi_1)$. \square

Proof of Theorem 1.3. Step 1: we prove (5a). By Definition 1.1, there are infinitely many integers $p, q > 0$ such that:

$$0 < \left| \pi - \frac{p}{q} \right| < \frac{1}{q^{\mu(\pi)-\varepsilon}}, \quad \varepsilon > 0.$$

We will make repeated use of the following Taylor approximation:

$$|\cos x| \geq 1 - \frac{x^2}{2}, \quad \text{for } x \in \mathbb{R}. \quad (7)$$

Choosing arbitrary integers p and $Q(p)$ satisfying the above inequality and combining the above inequalities, we have:

$$|\cos p| = |\cos(p - \pi Q(p))| \geq 1 - \frac{(p - \pi Q(p))^2}{2} > 1 - \frac{1}{2Q(p)^{2(\mu(\pi)-1-\varepsilon)}}. \quad (8)$$

Raising (8) to p^γ and applying the Bernoulli inequality we have:

$$|\cos p|^{p^\gamma} \geq \left(1 - \frac{1}{2Q(p)^{2(\mu(\pi)-1-\varepsilon)}}\right)^{p^\gamma} > 1 - \frac{p^\gamma}{2Q(p)^{2(\mu(\pi)-1-\varepsilon)}}. \quad (9)$$

Observe $\limsup |\cos k|^{k^\gamma} \leq 1$.

Let $\gamma < 2(\mu(\pi) - 1 - \varepsilon)$. We have $p^\gamma = o(Q(p)^{2(\mu(\pi)-1-\varepsilon)})$ because $Q(p) = \mathcal{O}(p)$ by Lemma 2.2. Since we can choose arbitrarily large p and $Q(p)$ satisfying (1), we can bring $|\cos p|^{p^\gamma}$ arbitrarily close to 1 by (9). This establishes (5a).

Step 2: we show (5b). First, observe that $|\cos x| \leq 1 - \frac{x^2}{2} + \frac{x^4}{24}$ for $x \in [-\frac{\pi}{2}, \frac{\pi}{2}]$. Here, the interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$ is chosen arbitrarily; any interval which is small enough to ensure the inequality would suffice (the inequality does not hold for all x). Fix $\varepsilon > 0$ and let $\gamma = 2(\mu(\pi) - 1 + \varepsilon)$. Then for arbitrary integers p and $Q(p)$, $|p - \pi Q(p)| \leq \frac{\pi}{2}$, we have:

$$|\cos p|^{p^\gamma} \leq \left(1 - \frac{(p - \pi Q(p))^2}{2} + \frac{(p - \pi Q(p))^4}{24}\right)^{p^\gamma}. \quad (10)$$

By Definition 1.1, for any real number $\lambda > 0$, there exists an integer N such that

$$|p - q\pi| > \frac{1}{q^{\mu(\pi)-1+\lambda}}, \quad \text{for all } p, q > N. \quad (11)$$

Since $1 - \frac{x^2}{2} + \frac{x^4}{24}$ is monotonically increasing on $[-\frac{\pi}{2}, 0)$ and monotonically decreasing on $(0, \frac{\pi}{2}]$, for $p, Q(p) > N$, we have:

$$\begin{aligned} |\cos p|^{p^\gamma} &\leq \left(1 - \frac{(p - \pi Q(p))^2}{2} + \frac{(p - \pi Q(p))^4}{24}\right)^{p^\gamma} \\ &\leq \left(1 - \frac{1}{2Q(p)^{2(\mu(\pi)-1+\lambda)}} + \frac{1}{24Q(p)^{4(\mu(\pi)-1+\lambda)}}\right)^{p^\gamma} \\ &= \left(1 - \frac{12Q(p)^{2(\mu(\pi)-1+\lambda)} - 1}{24Q(p)^{4(\mu(\pi)-1+\lambda)}}\right)^{p^\gamma}. \end{aligned} \quad (12)$$

Observe that if we define a function $\phi(p)$ such that:

$$\frac{1}{\phi(p)} = \frac{12Q(p)^{2(\mu(\pi)-1+\lambda)} - 1}{24Q(p)^{4(\mu(\pi)-1+\lambda)}},$$

then $\phi(p) = \mathcal{O}(Q(p)^{2(\mu(\pi)-1+\lambda)})$. If $\gamma > 2(\mu(\pi) - 1 + \lambda)$, then we have $\phi(Q(p)) = o(p^\gamma)$. Since we can choose arbitrarily large p and $Q(p)$ satisfying (11), by Lemma 2.3 we can bring (10) arbitrarily close to 0.

Thus $\limsup\{|\cos k|^{k^\gamma}\}_{k \in \mathbb{N}} = 0$ for all $\gamma > 2(\mu(\pi) - 1 + \lambda)$. Since our choice of λ is arbitrary but greater than zero, this establishes (5b). By (5a)–(5b) the proof is complete. \square

3 Further investigations

Theorem 3.1. $\limsup |\cos n|^{n^2} \geq \exp(-\pi^2/2) \approx 0.007192$.

Proof. By Diophantine approximation and (7), (8) becomes:

$$|\cos p| = |\cos(p - \pi Q(p))| \geq 1 - \frac{(p - \pi Q(p))^2}{2} > 1 - \frac{1}{2Q(p)^2} \quad (13)$$

Raising (13) to the p^2 , we obtain

$$|\cos p|^{p^2} \geq \left(1 - \frac{1}{2Q(p)^2}\right)^{p^2}. \quad (14)$$

Fix $\varepsilon > 0$. Since $\frac{p}{Q(p)} \rightarrow \pi$ by Lemma 2.2, we have $\pi^2 - \varepsilon < \frac{p^2}{Q(p)^2} < \pi^2 + \varepsilon$ for all sufficiently large p , whence

$$\left(1 - \frac{\frac{\pi^2 - \varepsilon}{2}}{p^2}\right)^{p^2} > \left(1 - \frac{p^2}{2p^2 Q(p)^2}\right)^{p^2} > \left(1 - \frac{\frac{\pi^2 + \varepsilon}{2}}{p^2}\right)^{p^2}.$$

It follows that the limit of the right side of (14) is $\exp(-\pi^2/2)$. \square

3.1 Cosine identities

We manipulate the following identity to obtain insight on the sequence:

$$\prod_{k=1}^n \cos \theta_k = \frac{1}{2^n} \sum_{e \in S} \cos(e_1 \theta_1 + \cdots + e_n \theta_n), \quad (15)$$

where $S = \{1, -1\}^n$.

Theorem 3.2. *We have the estimate*

$$\cos(n)^{n^2} \geq 1 - \frac{n^{4-2\mu(\pi)}}{2}. \quad (16)$$

Proof. Choosing $\theta_k = n$ and $S = \{1, -1\}^{n^2}$ in (15), the identity becomes:

$$\cos(n)^n = \frac{1}{2^n} \sum_{e \in S} \cos((e_1 + \cdots + e_n)n) = \frac{1}{2^n} \sum_{i=0}^n \binom{n}{i} \cos((2i - n)n).$$

Observe that for $\cos(n)^{n^2}$, we have:

$$\cos(n)^{n^2} = \frac{1}{2^{n^2}} \sum_{i=0}^{n^2} \binom{n^2}{i} \cos((2i - n^2)n). \quad (17)$$

Applying the cosine approximation (7) and writing $Q(n) = \left[\frac{n}{\pi}\right]$, for large choices of n which satisfy (1), we have:

$$\cos(n)^{n^2} \geq \frac{1}{2^{n^2}} \sum_{i=0}^{n^2} \binom{n^2}{i} \left(1 - \frac{(2i - n^2)^2 (n - \pi \left[\frac{n}{\pi}\right])^2}{2}\right)$$

$$\begin{aligned}
&= \frac{1}{2^{n^2}} \sum_{i=0}^{n^2} \left(\binom{n^2}{i} - \binom{n^2}{i} \frac{(2i - n^2)^2}{2} (n - \pi \lfloor \frac{n}{\pi} \rfloor)^2 \right) \\
&= 1 - \frac{(n - \pi Q(n))^2}{2 \cdot 2^{n^2}} \sum_{i=0}^{n^2} \binom{n^2}{i} (2i - n^2)^2 \\
&= 1 - \frac{n^2 (n - \pi Q(n))^2}{2} \tag{18}
\end{aligned}$$

where we have used the identity $\sum_{i=0}^{n^2} \binom{n^2}{i} (2i - n^2)^2 = n^2 2^{n^2}$ in the last step. Definition 1.1 then yields

$$\cos(n)^{n^2} \geq 1 - \frac{n^2}{2n^{2\mu(\pi)-2}}$$

and (16) follows. \square

From (16), we can see that the existence of a lower bound (as we pick larger n which better approximate multiples of π) is determined by the growth rate of the second term. If the rate is $\mathcal{O}(1)$, then there is a positive lower bound. If the growth rate is $o(1)$, then the lower bound is 1, and the limsup is also 1.

Additionally, we can get a more precise lower bound of (16) by adding more terms to the Taylor expansion (note that we continue choose large values of n which satisfy (1)):

$$\begin{aligned}
\cos(n)^{n^2} &= 1 - \frac{n^2 (n - \pi Q(n))^2}{2!} + \frac{n^2 (3n^2 - 2) (n - \pi Q(n))^4}{4!} \\
&\quad - \frac{n^2 (16 + 15n^2 (n^2 - 2)) (n - \pi Q(n))^6}{6!} \dots \tag{19}
\end{aligned}$$

where all of the identities of the form $\sum_{i=0}^{n^2} \binom{n^2}{i} (2i - n^2)^a$ were computed with Mathematica. Thus, we have shown the following.

Corollary 3.3. *The bound in Theorem 3.2 can be improved to*

$$\cos(n)^{n^2} \geq 1 - \frac{n^2}{2!n^{2\mu(\pi)-2}} + \frac{n^2 (3n^2 - 2)}{4!n^{4\mu(\pi)-4}} - \frac{n^2 (16 + 15n^2 (n^2 - 2))}{6!n^{6\mu(\pi)-6}} \dots \tag{20}$$

Now it is clear from (20) that if we assume $\mu(\pi) = 2$, then only the highest coefficient of the numerator will remain when we take the lim sup:

$$\limsup |\cos(n)^{n^2}| \geq 1 - \frac{1}{2!} + \frac{3}{4!} - \frac{15}{6!} + \frac{105}{8!} - \frac{945}{10!} \dots \approx 0.6065 \tag{21}$$

Note that the numerator coefficients we use in (21) were from identities found using Mathematica. A full table of these coefficients up to denominator $16!$ is shown below:

Denominator	Numerator
2!	-1
4!	3
6!	-15
8!	105
10!	-945
12!	10395
14!	-135135
16!	2027025

Since we are determining these coefficients through Mathematica, we do not know the growth rate of the coefficients. It is possible that the series could diverge, but as long as we truncate the series at a negative term, the lower bound holds.

Corollary 3.4. $\limsup |\cos n|^{n^2} \geq 0.6065 \dots$

Proof. $\mu(\pi) \geq 2$, by transcendentality of π . If $\mu(\pi) = 2$, then $\limsup |\cos n|^{n^2} \geq 0.6065 \dots$ by (21). If $\mu(\pi) > 2$, then $\limsup |\cos n|^{n^2} = 1$ by Theorem 1.3. This completes the proof. \square

4 Numerical results

Computationally, we find values of $|\cos k|^{k^\gamma}$ very close to one for values of k on the order of 10^8 , with values of γ very close to 2. When we set $\gamma = 2$, we do not find values of k close to one. This is evidence that $\mu(\pi) = 2$, but a proof has evaded us so far. Various values of γ are plotted in Figure 2 and Figure 3.

We see that for larger values of γ , points near one get more sparse but still reach values very close to 1. In the case of $\gamma = 1$, there are interesting subsequences, some of which persist quasiperiodically or die out entirely. Investigations of these subsequences are an interesting subtopic which is open to exploration. Figure 4 shows plots of γ near and above 2.




Figure 2 Left: Varying values of γ , plotted from 0 to 170000. Right: $\gamma = 1$ plotted from 0 to 10^6 .




Figure 3 Left: $\gamma = 1$ plotted from 0 to 200000. Right: $\gamma = 1.5$ plotted from 0 to 10^6 .




Figure 4 Plots of $(\cos n)^{\gamma}$, for $\gamma \approx 2$.

When γ is just below two, we still see subsequences that are close to 1, but they become more sparse as γ is closer to 2. In the case where $\gamma = 2.1$, we see that the values approach zero, and do not appear to take on values close to one in the range sampled, which is expected if $\mu(\pi) = 2$.

5 Subsequences of $(\cos n)^{n^\gamma}$

Various interesting subsequences appear in $(\cos n)^{n^\gamma}$, and the following theorems shed some light on the topic.

Theorem 5.1. *For all $0 \leq \gamma < 2(\mu(\pi) - 1)$, for any real number $\alpha \in (0, 1)$, there exist infinitely many arbitrarily long arithmetic progressions \mathbf{S} such that for all $a_n \in \mathbf{S}$, $|\cos a_n|^{a_n^\gamma} > \alpha$.*

Proof. The proof relates closely to the proof of (5a). We modify (9) to obtain:

$$|\cos(np)|^{(np)^\gamma} \geq \left(1 - \frac{(n(p - \pi Q(p))^2)^{(np)^\gamma}}{2}\right) > 1 - \frac{n^{\gamma+2} p^\gamma}{2Q(p)^{2(\mu(\pi)-1-\epsilon)}}$$

By (1), for any $n \in \mathbb{N}$, we can bring $|\cos(np)|^{(np)^\gamma}$ arbitrarily close to 1, which is greater than α . Also observe that for any $a \in \mathbb{N}$, $a < n$ we have:

$$\frac{n^{\gamma+2} p^\gamma}{2Q(p)^{2(\mu(\pi)-1-\epsilon)}} > \frac{a^{\gamma+2} p^\gamma}{2Q(p)^{2(\mu(\pi)-1-\epsilon)}}$$

Thus for any series of length $n \in \mathbb{N}$, there exist infinitely many integers p such that the sequence $\{|\cos(kp)|^{(kp)^\gamma}, k = 0, 1, 2, \dots, n\}$ is bounded below by $\alpha \in (0, 1)$. This completes the proof. \square

Definition 5.2. We refer to the subsequences which Theorem 5.1 guarantees to exist as *persistent subsequences*.

The terminology of Definition 5.2 stems from the idea that these values of $\cos(n_k)$ are sufficiently close to 1 that not even raising them to the (typically extremely large) number n_k will result in a number near 0.

Numerical experimentation reveals that **each peak** in Figure 1 corresponds to a separate persistent subsequence

$$\cos(n_k)^{n_k}, \quad \text{where } n_k = 355k + (3 + 22j), \quad k \in \mathbb{N}, \quad (22)$$

for some fixed $j \in \mathbb{N}$. The appearance of $p = 355$ is due to its role in the continued fraction expansion of π and the fact that $\frac{355}{113}$ is the best rational approximation of π with denominator less than 16604, and the appearance of 22 is also likely due to $\pi \approx \frac{22}{7}$. This suggests that other persistent subsequences may be found by looking at other exceptionally accurate fractional approximations to π , for example: $\frac{52163}{16604}$, $\frac{833719}{265381}$ and $\frac{42208400}{13435351}$. Indeed, we find persistent subsequences for $n_k = 833719k$ and $n_k = 833719k + 42208400$, as in Figure 5.

Each persistent subsequence appears to have a single peak, of a shape very similar to a Gaussian distribution (for those whose peak is not too close to 0). Curve matching in MATLAB reveals that these peaks match well. For example, the persistent subsequence $\cos(n_k)^{n_k}$ with $n_k = 644 + 355k$ has a coefficient of determination $R^2 = 0.9983$ with the




Figure 5 Left: the persistent subsequence for $n_k = 833719k$. Right: the persistent subsequence for $n_k = 833719k + 42208400$. Notice: the scale on the horizontal axis is $\times 10^8$ (right) and $\times 10^9$ (left).




Figure 6 $|\cos(8.49137 \cdot 10^{-8}x - 0.0353982)|^{x^{1.2}}$ plotted against the sequence $|\cos(a_n)|^{a_n^{1.2}}$.

Gaussian $0.9978 e^{-\left(\frac{x-260.8}{84.57}\right)^2}$. However, these curves are not truly Gaussian; the following theorem derives the general peak shape we observe in persistent subsequences and is matched against a sample subsequence in Figure 6.

Theorem 5.3. *For any arithmetic progression $a_n = pn + d$, the sequence $\{|\cos a_n|^{a_n^\gamma}\}$ lies on the curve*

$$f(x) = |\cos((p - \pi Q(p))\alpha(x) + d - \pi Q(d))|^{x^\gamma}. \quad (23)$$

Proof. The proof begins similarly to the proof of Theorem 5.1. Let \mathbf{S} be an arithmetic progression of the form $S_n = pn + d$, where $p, d \in \mathbb{Z}$. Then we have:

$$|\cos(pn + d)|^{(pn+d)^\gamma} = |\cos(n(p - \pi Q(p)) + (d - \pi Q(d)))|^{(pn+d)^\gamma}.$$

The domain can be extended into the reals via the transformation $\alpha(x) = \frac{x-d}{p}$, and this yields (23). \square

Appendix: Irrationality measure of rational and algebraic numbers

Theorem A.1 ([3, Thm. E.2]). *If $\alpha \in \mathbb{Q}$, then $\mu(\alpha) = 1$.*

Proof. Since $|q\alpha - [q\alpha]| < 1$ for every $q \in \mathbb{N}$, we have $\mu(\alpha) \geq 1$. If $\alpha = \frac{a}{b}$ in lowest terms, and $\alpha \neq \frac{p}{q}$, then $|\alpha - \frac{p}{q}| \geq \frac{1}{|bq|}$, whence $\mu(\alpha) = 1$. \square

Using continued fraction expansions, one can prove that $\mu(\alpha) \geq 2$ whenever $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, but that is beyond the scope of this paper. However, we can show that all Liouville numbers are transcendental. It suffices to prove the following result; our treatment is adapted from [8].

Theorem A.2. *For all irrational algebraic numbers α of degree n , there exists a constant $c > 0$ such that: $|\alpha - \frac{p}{q}| \geq \frac{c}{q^n}$ for any $p, q \in \mathbb{Z}$ with $q > 0$.*

Proof. Let $P(x) = \sum_{k=0}^n a_k x^k$ be a polynomial of degree n such that $P(\alpha) = 0$, and denote the set of rational roots of P by $R = \{\zeta \in \mathbb{Q} : P(\zeta) = 0\}$. For a given rational $r = \frac{p}{q}$, we consider the following three cases: (i) $|\alpha - r| \geq 1$, (ii) $r \in R$, and (iii) $r \notin R$ but $|\alpha - r| \leq 1$.

Case (i). For any $r = \frac{p}{q}$ satisfying $|\alpha - r| \geq 1$, we have $|\alpha - r| \geq \frac{1}{q^n}$, and $c = 1$ would work.

Case (ii). For $r \in R$, we can define $\xi > 0$ by

$$\xi = \begin{cases} \min\{|\alpha - r| : r \in R\}, & R \neq \emptyset \\ 1, & R = \emptyset, \end{cases}$$

and we immediately have $|\alpha - r| \geq \xi \geq \frac{\xi}{q^n}$.

Case (iii). For $r \notin R$, we know that $P(r)$ is some multiple of $\frac{1}{q^n}$. Since $r \notin R$, this implies

$$|P(\alpha) - P(r)| = |P(r)| \geq \frac{1}{q^n}. \quad (24)$$

Observe that

$$\alpha^k - r^k = (\alpha - r) \sum_{i=0}^{k-1} r^i \alpha^{k-1-i}.$$

Solving this identity for r^k and substituting it into the definition of P leads to

$$P(\alpha) - P(r) = (\alpha - r) \sum_{k=1}^n a_k \sum_{i=0}^{k-1} r^i \alpha^{k-1-i}.$$

Recall that for this case we have also assumed $|\alpha - r| \leq 1$. For such r , we have $|r| \leq |\alpha| + 1$, and so for c_α defined by

$$c_\alpha = \sum_{k=1}^n |a_k| k (|\alpha| + 1)^{k-1},$$

we get $|P(\alpha) - P(r)| \leq |\alpha - r| c_\alpha$ by the triangle inequality.

In combination with (24), this gives

$$|\alpha - r| \geq \frac{|P(\alpha) - P(r)|}{c_\alpha} \geq \frac{1}{c_\alpha q^n}.$$

For $c = \min\left(1, \zeta, \frac{1}{c_\alpha}\right)$, all cases are covered. \square

Acknowledgement

The first author would like to thank the second author for suggesting this problem and assisting greatly with the development of the paper and for the second author's continued mentoring and guidance. The first author would also like to thank Math StackExchange user "T. Bongers" [2] for his/her/their insight on a more specific version of the problem. The second author acknowledges that the first author did all the hard work, and is grateful to Jordan Rowley for raising the question about $\sin(n)^n$ that led to the current work ($\cos(n)^n$ turned out to be more convenient to study). Both authors would like to thank the referee for helpful comments which we feel improved the paper significantly.

References

- [1] Max A. Alekseyev. *On convergence of the Flint Hills series*. arXiv:1104.5100 [math.CA]
- [2] T. Bongers. *End behavior of $(\sin k)^k$ where $k \in \mathbb{N}$* . URL (version: 2018-04-24): <https://math.stackexchange.com/q/2752127>
- [3] Yann Bugeaud. *Distribution modulo one and Diophantine approximation* (2012). Cambridge University Press, 193.
- [4] Ernie Croot, & Olof Sisask. *A new proof of Roth's theorem on arithmetic progressions* (2009). Proceedings of the American Mathematical Society, 137(3), 805–809.
- [5] J. Liouville. *Remarques relatives à des classes très-étendues de quantités dont la valeur n'est ni algébrique, ni même réductible à des irrationnelles algébriques* (1844). C. R. Acad. Sci. Paris 18, 883–885.
- [6] V.Kh. Salikhov. *On the irrationality measure of $\ln 3$* . (2007) Dokl. Math. 76: 955.
- [7] V.Kh. Salikhov. *On the irrationality measure of π* . Uspekhi Mat. Nauk, 63:3(381) (2008), 163–164; Russian Math. Surveys, 63:3, 570–572
- [8] Simmons, George F. *Calculus gems: brief lives and memorable mathematics*. (2007) MAA 55.
- [9] Weisstein, Eric W. *Irrationality Measure*. From MathWorld—A Wolfram Web Resource. <http://mathworld.wolfram.com/IrrationalityMeasure.html>
- [10] Weisstein, Eric W. *Liouville's Constant*. From MathWorld—A Wolfram Web Resource. <http://mathworld.wolfram.com/LiouvillesConstant.html>
- [11] Weisstein, Eric W. *Liouville Number*. From MathWorld—A Wolfram Web Resource. <http://mathworld.wolfram.com/LiouvilleNumber.html>

Sully F. Chen
University of Southern California
Department of Chemistry
Los Angeles, CA 90089-1062, USA
e-mail: sullyche@usc.edu

Erin P.J. Pearse
Mathematics Department
California Polytechnic University
San Luis Obispo, CA 93407, USA
e-mail: epearse@calpoly.edu