

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 75 (2020)
Heft: 4

Artikel: An inequality for cevians and applications
Autor: Lukarevski, Martin / Marinescu, Dan Stefan
DOI: <https://doi.org/10.5169/seals-880893>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 02.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

An inequality for cevians and applications

Martin Lukarevski and Dan Stefan Marinescu

Martin Lukarevski obtained his Ph.D. from Leibniz University Hannover in 2012. Currently he works as an associate professor in the Faculty of Informatics at the University “Goce Delcev” (North Macedonia). His interests include classical geometry, analysis and history of mathematics. As a passionate cinephile, he is a frequent moviegoer.

Dan Stefan Marinescu is a teacher at the National College Iancu de Hunedoara, Hunedoara, Romania. He has a Ph.D. in mathematics, obtained from the University Babes-Bolyai of Cluj Napoca.

1 The inequality for cevians

In a triangle ABC with sides a, b, c , semiperimeter s , circumradius R and inradius r , let AD be a cevian. We give a lower bound for its length, in terms of its adjacent sides and corresponding angle.

Theorem 1. *For the cevian AD with $BD/DC = \lambda$ and angle $\angle BAC = \alpha$, we have the inequality*

$$AD \geq \left(\frac{\lambda}{\lambda + 1} b + \frac{1}{\lambda + 1} c \right) \cos \frac{\alpha}{2}. \quad (1)$$

Equivalently, if the ratio $BD/BC = k$ is given, with $k' = 1 - k$, the inequality is

$$AD \geq (kb + k'c) \cos \frac{\alpha}{2}. \quad (2)$$

Cevane oder Ecktransversalen sind Geraden, welche durch die Ecke eines Dreiecks verlaufen. Sie spielen eine zentrale Rolle in der Dreiecksgeometrie. Ein Beispiel ist der bekannte Satz von Ceva, bei dem es darum geht, in welchen Verhältnissen drei kopunktuale Ecktransversalen die gegenüberliegenden Dreiecksseiten teilen. Es ist aber auch nützlich, Aussagen über die Längen von Ecktransversalen zu finden. Die Autoren der vorliegenden Arbeit geben hier eine allgemeine untere Schranke für diese Längen an. Daraus ergeben sich interessante Folgerungen für spezielle Ecktransversalen sowie für das Produkt der Längen von drei kopunktalen Ecktransversalen.

Proof. The proof is given for the inequality in form (2). Let M be a point on the side AB , such that DM is parallel to CA . We construct two similar right triangles, AMF and DME , with right angles at vertices F and E , and angles equal to $\alpha/2$ at vertices A and D , see Figure 1.




Figure 1

Then the lengths of AF and DE are $k'c \cos \frac{\alpha}{2}$ and $kb \cos \frac{\alpha}{2}$ respectively. It is evident that the length of the cevian AD is greater than or equal to their sum. \square

The inequality is sharp, with equality when the cevian AD is the bisector w_a of α . In that case $\lambda = c/b$, Euclid, Book VI. 3. On the other hand by Pappus $w_a = \frac{2bc}{b+c} \cos \frac{\alpha}{2}$, hence equality holds in (1).

We could have proceeded applying the elementary inequality for a triangle ABC

$$\sin \frac{\alpha}{2} \leq \frac{a}{b+c} \quad (3)$$

to the triangle AMD , where $\angle M = \angle DMA = \pi - \alpha$, to obtain

$$AD \geq (MD + MA) \sin \frac{\alpha}{2} = (kb + k'c) \cos \frac{\alpha}{2}.$$

The inequality (3) is a direct byproduct of Mollweide's formula [5], [7]

$$\sin \frac{\alpha}{2} / \cos \frac{\beta - \gamma}{2} = \frac{a}{b+c}.$$

It can also be derived geometrically by drawing perpendiculars from vertices B and C to the bisector of α .

2 Applications

Next we apply the inequality (1) to some special cevians. For the median m_a , we have $\lambda = 1$, so (1) gives

$$m_a \geq \frac{b+c}{2} \cos \frac{\alpha}{2}. \quad (4)$$

By the AM-HM and AM-GM inequalities $\frac{b+c}{2} \geq \frac{2bc}{b+c}$, $\frac{b+c}{2} \geq \sqrt{bc}$ and $\cos(\alpha/2) = \sqrt{s(s-a)/(bc)}$, $w_a = \frac{2bc}{b+c} \cos \frac{\alpha}{2}$, (4) implies the known useful inequalities

$$m_a \geq \sqrt{s(s-a)} \geq w_a.$$

From (4) and $m_a = \sqrt{2(b^2 + c^2) - a^2}/2$, follows an inequality for cosines similar to inequality (3)

$$\cos \frac{\alpha}{2} \leq \frac{\sqrt{2(b^2 + c^2) - a^2}}{b+c}. \quad (5)$$

The next inequalities for symmedians, Gergonne and Nagel Cevians are possibly new.

The *symmedian* is a cevian which is a reflection of the median in the corresponding angle bisector. The three symmedians are concurrent in the *symmedian point*, sometimes referred to as the Lemoine point or Grebe point.

Corollary 1. *For the symmedian s_a , the following inequality holds*

$$s_a \geq \frac{bc(b+c)}{b^2 + c^2} \cos \frac{\alpha}{2}. \quad (6)$$

Proof. It is well known [3] that the symmedian $s_a = AD$ divides the side BC in the ratio of the squares of the adjacent sides, that is $\lambda = c^2/b^2$. Substituting in (1), gives (6). \square

From the equation of the symmedian $s_a = bc\sqrt{2(b^2 + c^2) - a^2}/(b^2 + c^2)$, which can be derived from Stewart's theorem, and (6), follows again the inequality (5).

The three lines connecting each vertex of a triangle to the point of contact of the incircle and the opposite side are concurrent at the *Gergonne point* and are called *Gergonne cevians*, named after the French geometer Joseph Diaz Gergonne (1771–1859). *Nagel cevians*, named after the German geometer Christian Heinrich von Nagel (1803–1882), are the three lines concurrent at the *Nagel point*, connecting each vertex to the point of contact of the corresponding excircle and the opposite side. We denote the length of the Gergonne cevian AG_a by g_a and the length of the Nagel cevian AN_a by n_a . All the notation appears in Figure 2.

For Gergonne cevians

$$\lambda = \frac{BG_a}{G_aC} = \frac{s-b}{s-c},$$

hence by (1)

$$g_a = AG_a \geq \frac{b(s-b) + c(s-c)}{a} \cos \frac{\alpha}{2}.$$




Figure 2

For Nagel cevians

$$\lambda = \frac{BN_a}{N_aC} = \frac{s-c}{s-b},$$

hence by (1)

$$n_a = AN_a \geq \frac{b(s-c) + c(s-b)}{a} \cos \frac{\alpha}{2}.$$

We derive as a consequence of (1) an inequality for the product of cevians. Let AD , BE , CF be three cevians such that

$$\frac{BD}{DC} = \lambda, \quad \frac{CE}{EA} = \mu, \quad \frac{AF}{FB} = \nu.$$

Corollary 2. For the product of cevians AD , BE , CF , we have the inequality

$$AD \cdot BE \cdot CF \geq \frac{8rs^2\sqrt{\lambda\mu\nu}}{(\lambda+1)(\mu+1)(\nu+1)} \quad (7)$$

Proof. By (1) and the AM-GM inequality $\frac{\lambda}{\lambda+1}b + \frac{1}{\lambda+1}c \geq 2\sqrt{bc\lambda}/(\lambda+1)$, we get

$$AD \geq \frac{2\sqrt{bc\lambda}}{\lambda+1} \cos \frac{\alpha}{2}. \quad (8)$$

Multiplying (8) by the analogous inequalities for the cevians BE, CF

$$BE \geq \frac{2\sqrt{ca\mu}}{\mu+1} \cos \frac{\beta}{2}, \quad CF \geq \frac{2\sqrt{abv}}{v+1} \cos \frac{\gamma}{2},$$

and using the well-known identities $abc = 4Rrs$ and

$$\cos \frac{\alpha}{2} \cos \frac{\beta}{2} \cos \frac{\gamma}{2} = \frac{s}{4R},$$

we obtain (7). □

For concurrent cevians, Ceva's theorem [1], [3], [8] says that $\lambda\mu\nu = 1$, so the inequality (7) in that case simplifies to

Corollary 3. *Let the cevians AD, BE, CF be concurrent. Then the following inequality holds*

$$AD \cdot BE \cdot CF \geq \frac{8rs^2}{(\lambda+1)(\mu+1)(\nu+1)}. \quad (9)$$

Let us consider now the product of cevians of some special triangle centers.

For the centroid G and the corresponding medians holds $\lambda = \mu = \nu = 1$. Hence by (9) we have

$$m_a m_b m_c \geq rs^2.$$

Since for the exradii of the triangle, the identity $r_a r_b r_c = rs^2$ holds, this last inequality is actually the known inequality $m_a m_b m_c \geq r_a r_b r_c$ [2, 8.21].

The bisectors are the cevians of the incenter I and they divide the sides in the ratio of the corresponding sides: $\lambda = c/b$, $\mu = a/c$, $\nu = b/a$. Thus by (9), we have the elegant inequality for the product of bisectors

$$w_a w_b w_c \geq \frac{8rs^2 abc}{(a+b)(b+c)(c+a)}, \quad (10)$$

which complements $rs^2 \geq w_a w_b w_c$ [2, 8.14]. We remark that the inequality (10) is a refinement of $w_a w_b w_c \geq 8Rr^2 s^2 / (2R^2 + 3Rr + 2r^2)$ given in [6, p. 217]. Indeed, by $abc = 4Rrs$ and the identity [6, p. 53]

$$\prod (b+c) = 2s(s^2 + 2Rr + r^2),$$

the inequality

$$\frac{8rs^2 abc}{(a+b)(b+c)(c+a)} \geq \frac{8Rr^2 s^2}{2R^2 + 3Rr + 2r^2}$$

is equivalent to Gerretsen's inequality [4] $s^2 \leq 4R^2 + 4Rr + 3r^2$.

Next we give an inequality for the product of symmedians. Since for symmedians $\lambda = c^2/b^2$, $\mu = a^2/c^2$, $\nu = b^2/a^2$, by (9), we have

$$s_a s_b s_c \geq \frac{8rs^2 a^2 b^2 c^2}{(a^2 + b^2)(b^2 + c^2)(c^2 + a^2)}.$$

We end this article by asking for a geometric proof of inequality (5), similar to the proof of inequality (3) or the one given in Theorem 1.

Acknowledgement

We would like to express our gratitude to the anonymous referee above all for his delightful proof of the inequality in Theorem 1. We are also grateful for his valuable suggestions which led to an improved version of the paper.

References

- [1] N. Altshiller-Court, *College Geometry*, Barnes & Noble, 1952
- [2] O. Bottema, R.Z. Djordjevic, R.R. Janic, D.S. Mitrinovic, P.M. Vasic, *Geometric inequalities.*, Groningen, Wolters-Noordhoff, 1969
- [3] R. Johnson, *Advanced Euclidean Geometry*, Dover, 1960
- [4] M. Lukarevski, An alternate proof of Gerretsen's inequalities, *Elem. Math.* **72**, No. 1, (2017), 2–8
- [5] M. Lukarevski, Mollweide's formula and circumcevians of the incenter, *Elem. Math.* to appear
- [6] D.S. Mitrinovic, J. Pecaric, V. Volenec, *Recent Advances in Geometric Inequalities*, Kluwer Academic Publishers. Dordrecht (1989)
- [7] C.B. Mollweide, *Zusätze zur ebenen und sphärischen Trigonometrie*, Monatliche Correspondenz zur Beförderung der Erd- und Himmelskunde, (1808), 394–400
- [8] A. Ostermann, G. Wanner, *Geometry by Its History*, Springer, 2012

Martin Lukarevski

Department of Mathematics and Statistics

University "Goce Delcev"

Stip, North Macedonia

e-mail: martin.lukarevski@ugd.edu.mk

Dan Stefan Marinescu

Colegiul National "Iancu De Hunedoara"

Hunedoara, Romania

e-mail: marinescuds@gmail.com