**Zeitschrift:** Elemente der Mathematik

**Herausgeber:** Schweizerische Mathematische Gesellschaft

**Band:** 75 (2020)

**Heft:** 4

**Artikel:** On the inequalities of Grüss-ebyshev and Kantorovich : a probabilistic approach

**Autor:** Heinrich, Lothar

**DOI:** https://doi.org/10.5169/seals-880894

**Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

**Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

**Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 17.02.2026

**ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch**

Elem. Math. 75 (2020) 172 – 175
0013-6018/20/040172-4
DOI 10.4171/EM/419

© Swiss Mathematical Society, 2020

Elemente der Mathematik

*Short note*

# On the Inequalities of Grüss–Čebyshev and Kantorovich: A Probabilistic Approach

Lothar Heinrich

**Abstract.** First we recall the original form of inequalities found by P.L. Čebyshev in 1882, G. Grüss in 1935 and V.L. Kantorovich in 1948. Then we formulate generalized versions of these inequalities in the language of probability theory which allows to prove them by simple probabilistic arguments. A further moment inequality of this type rounds off this note.

## 1 Introduction and Results

V. Pták, see [6], provided a very short proof of the following inequality which has been first proved by V.L. Kantorovich, see [3] or [8], p. 85.

**Kantorovich Inequality:** For any real numbers $x_1, \ldots, x_n$ satisfying $0 < x_1 < \cdots < x_n < \infty$ and $\lambda_1, \ldots, \lambda_n \geq 0$ such that $\sum_{i=1}^n \lambda_i = 1$

$$\left(\sum_{i=1}^n \lambda_i x_i\right)\left(\sum_{i=1}^n \frac{\lambda_i}{x_i}\right) \leq \frac{(x_1 + x_n)^2}{4\, x_1\, x_n}\,. \tag{1}$$

In the particular case $\lambda_1 = \cdots = \lambda_n = \frac{1}{n}$ the inequality (1) has been shown by P. Schweitzer, see [7], by applying the fact that the function $\phi(x_1, \ldots, x_n) = \frac{1}{x_1} + \cdots + \frac{1}{x_n}$ is strictly Schur-convex and decreasing in each coordinate, see [5], p. 71. In Theorem 1 we formulate a slightly more general version of (1) emphasizing its probabilistic nature, where its proof is similarly short as that of (1) given in [6].

To state the next two inequalities we define for any two bounded functions $f, g : [a, b] \mapsto \mathbb{R}$ with $-\infty < a < b < \infty$

$$\Delta_{a,b}(f, g) := \frac{1}{b-a}\int_a^b f(x)g(x)\mathrm{d}x - \frac{1}{(b-a)^2}\int_a^b f(x)\mathrm{d}x \int_a^b g(x)\mathrm{d}x\,. \tag{2}$$

**Čebyshev Inequality:** If $f$ and $g$ are Lipschitz continuous on $[a, b]$ with Lipschitz constants $L_f$ and $L_g$, respectively, then

$$|\,\Delta_{a,b}(f, g)\,| \leq \frac{1}{12}\,(b-a)^2\, L_f\, L_g\,, \qquad \text{see [1].} \tag{3}$$

**Grüss Inequality:** If $f$ and $g$ are Riemann integrable on $[a, b]$ with bounded oscillations $\operatorname{osc}_a^b(f) := \sup\{f(x) : x \in [a, b]\} - \inf\{f(x) : x \in [a, b]\}$ and $\operatorname{osc}_a^b(g)$, respectively, then

$$|\,\Delta_{a,b}(f, g)\,| \le \frac{1}{4}\operatorname{osc}_a^b(f)\operatorname{osc}_a^b(g)\,, \qquad \text{see [2]}. \tag{4}$$

**Remarks:** The factors 1/12 in (3) and 1/4 in (4) cannot be replaced by smaller constants. For $a = 0, b = 1$ this is easily seen by inserting $f(x) = g(x) = 2x - 1$ in (3) and $f(x) = g(x) = \operatorname{sign}(2x - 1)$ in (4). E. Landau, see [4], pointed out that (4) holds in case of continuous functions $f, g$ with the best possible factor 4/45 instead of 1/4. More about Čebyshev-type and Grüss-type inequalities including historical notes the reader can find in [1], Chapter 3 and references therein. Among others there are provided discrete and operator versions of (3) and (4) in different settings.

In what follows, let $X$ and $Y$ denote real-valued random variables (short: rv's) on some probability space $[\Omega, \mathcal{F}, \mathsf{P}]$, where $\mathsf{E}$, $\mathsf{Var}$ and $\mathsf{Cov}$ designate expectation, variance and covariance, respectively. We remind the reader that $\mathsf{Cov}(X, Y) := \mathsf{E}(XY) - \mathsf{E}X\,\mathsf{E}Y$ and $\mathsf{Var}(X) := \mathsf{Cov}(X, X)$.

**Theorem 1 (Probabilistic Kantorovich-Type Inequality).** *Let $X$ be an rv taking values only in $[a, b]$ for $a, b \in \mathbb{R}$ such that $0 < a < b < \infty$. Then*

$$\mathsf{E}(X)\,\mathsf{E}\Big(\frac{1}{X}\Big) \le \frac{(a+b)^2}{4\,a\,b}\,. \tag{5}$$

Equality holds in (5) iff $\mathsf{P}(X = a) = \mathsf{P}(X = b) = 1/2$. Obviously, (1) follows from (5) if $\lambda_i = \mathsf{P}(X = x_i)$ for $i = 1, \ldots, n$ and $a = x_1 < \cdots < x_n = b$.

The key to formulate probabilistic versions of (3) and (4) consists in the simple fact that $\Delta_{a,b}(f, g)$ coincides with $\mathsf{Cov}(f(X), g(X))$ when $X$ is uniformly distributed on $[a, b]$.

**Theorem 2 (Probabilistic Čebyshev- and Grüss-Type Inequality).** *Let $X$ be an rv taking values in a finite or infinite interval $[a, b]$ such that $\mathsf{E}X^2 < \infty$. If $f, g : [a, b] \mapsto \mathbb{R}$ are Lipschitz continuous functions on $[a, b]$ with Lipschitz constants $L_f$, $L_g$, then*

$$|\,\mathsf{Cov}(f(X), g(X))\,| \le L_f\,L_g\,\mathsf{Var}(X) \le L_f\,L_g\,(\mathsf{E}X - a)(b - \mathsf{E}X)\,. \tag{6}$$

*If $f, g : [a, b] \mapsto \mathbb{R}$ are Borel-measurable functions on $[a, b]$ with bounded oscillations $\operatorname{osc}_a^b(f)$ and $\operatorname{osc}_a^b(g)$, then*

$$|\,\mathsf{Cov}(f(X), g(X))\,| \le \frac{1}{4}\operatorname{osc}_a^b(f)\operatorname{osc}_a^b(g)\,. \tag{7}$$

Finally, we state an estimate of the absolute central moment $\mathsf{E}|X - \mathsf{E}X|^p$ of order $p \ge 2$.

**Theorem 3.** *Let the rv $X$ take values in a finite interval $[a, b]$ with $\mu := \mathsf{E}X$. Then*

$$\mathsf{E}|\,X - \mu\,|^p \le (\max\{\mu - a, b - \mu\})^{p-2}\,(\mu - a)(b - \mu) \tag{8}$$

*with equality for $p = 2$ iff $\mathsf{P}(X \in \{a, b\}) = 1$ and for $p > 2$ iff $\mathsf{P}(X = a) = \mathsf{P}(X = b) = 1/2$.*

## 2 Proofs of the Theorems 1, 2, and 3

*Proof of Theorem* 1. Putting $Y = X/\sqrt{a\,b}$ implies that $\mathsf{P}(\sqrt{a/b} \le Y \le \sqrt{b/a}) = 1$ and $\mathsf{E}(X)\ \mathsf{E}(1/X) = \mathsf{E}(Y)\ \mathsf{E}(1/Y)$. Making use of the well-known inequality $\sqrt{u\,v} \le (u+v)/2$ for any real two numbers $u, v > 0$ (with equality iff $u = v$) we obtain the relation

$$\mathsf{E}(Y)\,\mathsf{E}\Big(\frac{1}{Y}\Big) \le \frac{1}{4}\Big(\mathsf{E}(Y) + \mathsf{E}\Big(\frac{1}{Y}\Big)\Big)^2 = \frac{1}{4}\Big[\sqrt{\frac{b}{a}} + \sqrt{\frac{a}{b}} - \mathsf{E}\Big(\frac{1}{Y}\Big(Y - \sqrt{\frac{a}{b}}\Big)\Big(\sqrt{\frac{b}{a}} - Y\Big)\Big)\Big]^2 .$$

Since the expectation term and also the whole term within the brackets are non-negative it is clear that the right-hand side of the latter relation is less than or equal to $(\sqrt{b/a} + \sqrt{a/b})^2/4 = (a+b)^2/4ab$ which proves (5). Equality in this estimate holds iff the expectation term disappears and this takes place iff the rv $Y$ is concentrated on the numbers $\sqrt{a/b}$ and $\sqrt{b/a}$, say $p := \mathsf{P}(Y = \sqrt{a/b}) = 1 - \mathsf{P}(Y = \sqrt{b/a})$. Then equality in the first estimate, that is $\mathsf{E}(Y) = \mathsf{E}(1/Y)$, is equivalent with

$$p\sqrt{\frac{a}{b}} + (1-p)\sqrt{\frac{b}{a}} = p\sqrt{\frac{b}{a}} + (1-p)\sqrt{\frac{a}{b}} .$$

This equation is satisfied just for $p = 1/2$ completing the proof of Theorem 1. □

*Proof of Theorem* 2. The inequality of Cauchy–Schwarz–Bunyakovsky, see [8], yields $|\,\mathsf{Cov}(f(X), g(X))\,| \le \sqrt{\mathsf{Var}(f(X))}\,\sqrt{\mathsf{Var}(g(X))}$. We estimate the variance $\mathsf{Var}(f(X))$ in two ways: First by using $|f(y) - f(x)| \le L_f\,|y - x|$ and the probability distribution $P_X(\cdot) = \mathsf{P}(\{\omega \in \Omega : X(\omega) \in (\cdot)\})$ on the Borel sets of $\mathbb{R}$ we get

$$\begin{aligned}\mathsf{Var}(f(X)) &= \mathsf{E}\big(f(X) - \mathsf{E}f(X)\big)^2 = \frac{1}{2}\int_{\mathbb{R}}\int_{\mathbb{R}} \big(f(y) - f(x)\big)^2 P_X(\mathrm{d}y)\,P_X(\mathrm{d}x) \\ &\le \frac{L_f^2}{2}\int_{\mathbb{R}}\int_{\mathbb{R}} (y-x)^2 P_X(\mathrm{d}y)\,P_X(\mathrm{d}x) = L_f^2\,\mathsf{Var}(X)\end{aligned}$$

implying $|\,\mathsf{Cov}(f(X), g(X))\,| \le L_f\,L_g\,\mathsf{Var}(X)$. This combined with (8) for $p = 2$ completes the proof of (6).

Second from the estimate $\mathsf{Var}(f(X)) \le \mathsf{E}\big(f(X) - c\big)^2$ for any $c \in \mathbb{R}$ we obtain for the special value $c_0 = (\inf_{a \le x \le b} f(x) + \sup_{a \le x \le b} f(x))/2$ that $\sup_{a \le x \le b} |\,f(x) - c_0\,| \le \mathrm{osc}_a^b(f)/2$ and thus

$$\sqrt{\mathsf{Var} f(X)} \le \mathrm{osc}_a^b(f)/2$$

proving (7). □

*Proof of Theorem* 3. We rewrite and estimate $\mathsf{E}|\,X - \mu\,|^p$ as follows

$$\begin{aligned}\mathsf{E}|\,X - \mu\,|^p &= \int_a^\mu (\mu - x)^p\,P_X(\mathrm{d}x) + \int_\mu^b (x - \mu)^p\,P_X(\mathrm{d}x) \\ &\le (\max\{\mu - a, b - \mu\})^{p-2}\,\mathsf{Var}(X) .\end{aligned}$$

To complete the proof of (8) we show that

$$\begin{aligned}\mathsf{Var}(X) &= \mathsf{E}\Big(X - \frac{a+b}{2}\Big)^2 + \mathsf{E}\Big[\,(X-\mu)^2 - \Big(X - \frac{a+b}{2}\Big)^2\Big] \\ &= \mathsf{E}\Big(X - \frac{a+b}{2}\Big)^2 - \Big(\mu - \frac{a+b}{2}\Big)^2 \\ &\le \Big(\frac{b-a}{2}\Big)^2 - \Big(\mu - \frac{a+b}{2}\Big)^2 = (b-\mu)(\mu-a)\,.\end{aligned}$$

On the other hand, the relation

$$0 = (b-\mu)(\mu-a) - \mathsf{E}(X-\mu)^2 = \mathsf{E}(X-a)(b-X)$$

is valid iff $\mathsf{P}(X \in \{a,b\}) = 1$. This means that $\alpha = \mathsf{P}(X = a) = 1 - \mathsf{P}(X = b)$ and $\mu = \alpha\, a + (1-\alpha)\, b$ for some $0 \le \alpha \le 1$. A short calculation reveals that equality in (8) is only possible for $\alpha = 1/2$. □

## References


[1] DRAGOMIR, S.S. *Operator Inequalities of the Jensen, Čebyšev and Grüss Type*, Springer, New York, 2012

[2] GRÜSS, G. Über das Maximum des absoluten Betrages von

$$\tfrac{1}{b-a}\int_a^b f(x)\,g(x)\,\mathrm{d}x - \tfrac{1}{(b-a)^2}\int_a^b f(x)\,\mathrm{d}x \int_a^b g(x)\,\mathrm{d}x,$$

*Math. Z.* **39** (1935), 215–226.

[3] KANTOROVICH, L.V. Functional analysis and applied mathematics (in Russian), *Uspechi matem. nauk* **3** (1948), 89–185.

[4] LANDAU, E. Über einige Ungleichungen von Herrn G. Grüss, *Math. Z.* **39** (1935), 742–744.

[5] MARSHALL, A.W. AND OLKIN, I. *Inequalities: Theory of Majorization and Its Applications*, Academic Press, New York, 1979.

[6] PTÁK, V. The Kantorovich inequality, *Amer. Math. Monthly* **102** (1995), 820–821.

[7] SCHWEITZER, P. An inequality concerning the arithmetic mean (in Hungarian), *Math.-Phys. Lapok* **23** (1914), 257–261.

[8] STEELE, J.M. *The Cauchy–Schwarz Master Class*, Cambridge University Press, New York, 2004.



Lothar Heinrich
Augsburg University
Institute of Mathematics
D-86135 Augsburg, Germany
e-mail: `heinrich@math.uni-augsburg.de`