

Distinct distances between lattice points

Autor(en): **Erdős, P. / Guy, R.K.**

Objektyp: **Article**

Zeitschrift: **Elemente der Mathematik**

Band (Jahr): **25 (1970)**

Heft 6

PDF erstellt am: **20.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-27359>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

ELEMENTE DER MATHEMATIK

Revue de mathématiques élémentaires – Rivista di matematica elementare

*Zeitschrift zur Pflege der Mathematik
und zur Förderung des mathematisch-physikalischen Unterrichts*

Publiziert mit Unterstützung des Schweizerischen Nationalfonds
zur Förderung der wissenschaftlichen Forschung

El. Math.

Band 25

Heft 6

Seiten 121-144

10. November 1970

Distinct Distances Between Lattice Points

How many points (x_i, y_i) , $1 \leq i \leq k$, with integer coordinates $0 < x_i, y_i \leq n$, may be chosen with all mutual distances distinct? By counting such distances, and pairs of differences of coordinates, we have

$$\binom{k}{2} \leq \binom{n+1}{2} - 1, \quad (1)$$

so that $k \leq n$, and for $2 \leq n \leq 7$ such a bound can be attained; e.g. for $2 \leq n \leq 5$, by the points (1,1), (1,2), (3,1), (4,4) and (5,3); for $n = 6$ by (1,1), (1,2), (2,4), (4,6), (6,3) and (6,6); and for $n = 7$ by (1,1), (1,3), (2,3), (3,7), (4,1), (6,6) and (7,7).

However, the fact that numbers may be expressed in more than one way as the sum of two squares indicates that this bound cannot be attained for $n > 15$. A result of LANDAU [4] states that the number of integers less than x expressible as the sum of two squares is asymptotically $c_1 x (\log x)^{-1/2}$, so we can replace the right member of (1) by $c_2 n^2 (\log n)^{-1/2}$ and we have the upper bound

$$k < c_3 n (\log n)^{-1/4}, \quad (2)$$

where c_i is in each case a positive constant.

A heuristic argument can be given to support the conjecture

$$(?) \quad k < c_4 n^{2/3} (\log n)^{1/6}, \quad (3)$$

but it lacks conviction since the corresponding argument in one dimension gives a false result.

On the other hand we can show

$$k > n^{2/3-\varepsilon} \quad (4)$$

for any $\varepsilon > 0$ and sufficiently large n , by means of the following construction. Choose points successively; when k points have been chosen, take another so that

(a) it does not lie on any circle having one of the k points as centre and one of the $\binom{k}{2}$

distinct distances determined by these points as radius.

(b) it does not form, with any of the first k points, a line with slope b/a , $(a, b) = 1$, $|a| < n^{1/3}$, $|b| < n^{1/3}$. Note that in particular no two points determine a distance less than $n^{1/3}$.

(c) it is not equidistant from any pair of the first k points.

We may choose such a point provided that all n^2 points are not excluded by these conditions.

Condition (a) excludes at most $k \binom{k}{2} n^{c_s/\log \log n}$ points, since there are $\binom{k}{2}$ circles round each of k points, and each circle contains at most $n^{c_s/\log \log n}$ lattice points¹⁾.

Condition (b) excludes at most

$$k \sum_{a=1}^{n^{1/3}} 4 \varphi(a) \frac{n}{a} < c_6 k n^{4/3}$$

points, since a line with slope b/a , $b < a$, $(a, b) = 1$, contains at most n/a lattice points.

Condition (c) excludes at most $\binom{k}{2} n^{2/3}$ points, since there are $\binom{k}{2}$ lines of equidistant points, each of which has slope b/a , $(a, b) = 1$, $|a| \geq n^{1/3}$ and such a line contains at most $n/|a| \leq n^{2/3}$ lattice points.

Hence, so long as

$$\frac{1}{2} k^3 n^{c_s/\log \log n} + c_6 k n^{4/3} + \frac{1}{2} k^2 n^{2/3} < n^2,$$

there remain eligible points, and this is the case if $k \leq n^{2/3-\epsilon}$. The lower bound (4) is thus established.

For the corresponding problem in one dimension, the existence of perfect difference sets [6] shows that for n an even power of a prime,

$$k \geq n^{1/2} + 1,$$

so that generally

$$k > n^{1/2} (1 - \epsilon). \quad (5)$$

On the other hand it is known [2, 5] that

$$k < n^{1/2} + n^{1/4} + 1. \quad (6)$$

In d dimensions, $d \geq 3$, we may replace Landau's theorem by the theorems on sums of three or four squares, giving an upper bound

$$k < c_7 d^{1/2} n, \quad (7)$$

while the corresponding heuristic argument suggests the conjecture

$$(?) \quad k < c_8 d^{2/3} n^{2/3} (\log n)^{1/3}. \quad (8)$$

The construction, with (hyper)spheres and (hyper)planes, corresponding to that given above, yields the same lower bound (4) as before.

One can also ask for configurations containing a *minimum* number of points, determining distinct distances, so that *no* point may be added without duplicating

¹⁾ It is well known that the number of solutions of $n = x^2 + y^2$ is less than or equal to $d(n)$, the number of divisors of n [3] and $d(n) < nc/\log \log n$ by a well known result of WIGERT [3].

a distance. Can this be done with as few as $O(n^{1/2})$ points; or with $O(n^{1/3})$ points in one dimension?

Another open problem [1] is given any n points in the plane (not necessarily lattice points) [or in d dimensions], how many can one select so that the distances which are determined are all distinct? P. ERDÖS and R. K. GUY, Budapest

REFERENCES

[1] ERDÖS, P., *Néhány geometriai problémáról* (in Hungarian), Mat. Lapok 8, 86–92 (1957); M. R. 20, 6056 (1959).
 [2] ERDÖS, P. and TURAN, P., *On a Problem of Sidon in Additive Number Theory and Some Related Problems*, J. London Math. Soc. 16, 212–215 (1941); M. R. 3, 270 (1942).
 [3] HARDY, G. H. and WRIGHT, E. M., *Introduction to the Theory of Numbers*, 4th ed. (Oxford 1960).
 [4] LANDAU, E., *Handbuch der Lehre von der Verteilung der Primzahlen* (Leipzig 1909), II, 643.
 [5] LINDSTRÖM, B., *An Inequality for B_2 -sequences*, J. Combinatorial Theory 6, 211–212 (1969).
 [6] SINGER, J., *A Theorem in Finite Projektive Geometry and Some Applications to Number Theory*, Trans. Amer. Math. Soc. 43, 377–385 (1938).

Note on a Diophantine Equation

SCHINZEL and SIERPIŃSKI [1] have given the general solution of the diophantine equation

$$(x^2 - 1)(y^2 - 1) = \left[\left(\frac{x - y}{2} \right)^2 - 1 \right]^2,$$

and SZYMICZEK [2] has given the general solution of

$$(x^2 - z^2)(y^2 - z^2) = \left[\left(\frac{y - x}{2} \right)^2 - z^2 \right]^2.$$

The purpose of this paper is to obtain a complete solution of the diophantine equation

$$(x^2 + a)(y^2 + a) = \left[a \left(\frac{y - x}{2b} \right)^2 + b^2 \right]^2, \tag{1}$$

where a and b are any two given integers.

Let $X = x - y$, $Y = x + y$; then $X \equiv Y \pmod{2}$ and (1) becomes

$$b^4 (X^2 + 2XY + Y^2 + 4a)(X^2 - 2XY + Y^2 + 4a) = (aX^2 + 4b^4)^2.$$

This equation reduces to

$$b^4 ((Y^2 - X^2)^2 + 8a(Y^2 - X^2) + 16a^2) = (aX^2 + 4b^4)^2 - 16ab^4X^2$$

and we have

$$b^2 (Y^2 - X^2 + 4a) = \pm (aX^2 - 4b^4).$$