

# Convexity of elementary functions with applications to inequalities

Autor(en): **Jovanovi, Milan / Jovanovi, Vladimir**

Objektyp: **Article**

Zeitschrift: **Elemente der Mathematik**

Band (Jahr): **66 (2011)**

PDF erstellt am: **20.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-283489>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

---

---

## Convexity of elementary functions with applications to inequalities

---

---

Milan Jovanović and Vladimir Jovanović

Milan Jovanović received his Ph.D. from the University of Belgrade (Serbia) in 1989. He is now full professor at the University of Banja Luka (Bosnia and Herzegovina). His main interest is convex optimization.

Vladimir Jovanović received his Ph.D. from the Albert-Ludwigs Universität Freiburg (Germany) in 2004. At present he is assistant professor at the University of Banja Luka (Bosnia and Herzegovina). His research focuses on numerical methods for partial differential equations.

### 1. Introduction

Let  $I \subset \mathbb{R}$  be an interval. The function  $f : I \rightarrow \mathbb{R}$  is *convex*, if the inequality

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y) \quad (1)$$

holds for all  $\lambda \in [0, 1]$  and  $x, y \in I$ . If instead  $\leq$  in (1) we have  $\geq$ , such a function is *concave*.

In most textbooks convexity is treated within differential calculus. However, we will see how one can single out convex functions among the basic elementary ones without exploiting differential calculus. This will be achieved by using several inequalities and simple assertions. In addition, some elementary convex functions, as it was pointed out in Jensen's

Konvexität bzw. Konkavität differenzierbarer Funktionen lässt sich bekanntlich leicht mit den Mitteln der Differentialrechnung nachweisen. In der vorliegenden Arbeit wird ein Werkzeugkatalog zusammengestellt, der es erlaubt, Konvexität bzw. Konkavität von Funktionen zu zeigen, die nicht notwendigerweise differenzierbar sind. Beispielsweise genügt es für den Nachweis der Konvexität einer stetigen Funktion  $f$  die Gültigkeit der Ungleichung  $f((x + y)/2) \leq (f(x) + f(y))/2$  für alle  $x, y \in \mathbb{R}$  zu verifizieren. Mit ihrem Katalog untersuchen die Autoren die Konvexität bekannter elementarer Funktionsklassen in systematischer Weise. Mit Hilfe dieses Formalismus gelingt es ihnen überdies, die Konvexität komplexerer Funktionen relativ leicht einzusehen und daraus bekannte Ungleichungen erneut abzuleiten.

fundamental paper [5], are not differentiable, such as

$$x \mapsto \sum_{i=1}^n |x - x_i| \quad (x_i \in \mathbb{R}, \quad i = 1, \dots, n).$$

Moreover, in the same paper Jensen provided an example (attributed to U. Dini and J. Lüroth) of a convex function defined on  $(0, 1)$ , which is not differentiable at any rational point:

$$f(x) = \sum_{i=1}^{\infty} c_i |x - x_i|.$$

Here  $c_i > 0$ , the series  $\sum_{i=1}^{\infty} c_i$  is convergent and  $\{x_i : i \in \mathbb{N}\} = \mathbb{Q} \cap (0, 1)$ . The proof follows from the relation

$$f'_+(x_i) - f'_-(x_i) = 2c_i,$$

where  $f'_\pm$  are the one-sided derivatives of  $f$ .

It should be mentioned that for any convex function  $f$  the derivatives  $f'_\pm$  exist and  $f'_- \leq f'_+$  in the interior of  $I$ . Furthermore,  $f'_+ = f'_-$  up to a countable set (see, e.g., [8]).

Besides having the one-sided derivatives, convex functions have another natural property: continuity. More precisely, if  $f : I \rightarrow \mathbb{R}$  is convex, then it is continuous in the interior of  $I$  (see also [8]).

Instead of convex functions defined by (1), Jensen in [5] considered the functions satisfying (1) only for  $\lambda = \frac{1}{2}$ :

$$f\left(\frac{x+y}{2}\right) \leq \frac{f(x) + f(y)}{2} \quad (x, y \in I). \quad (2)$$

It turns out that these functions, which are called *Jensen-convex* (or *J-convex*) functions, are also convex, provided they are continuous. In the proof of this fact Jensen used the inequality

$$f\left(\frac{x_1 + \dots + x_n}{n}\right) \leq \frac{f(x_1) + \dots + f(x_n)}{n}, \quad (3)$$

which today bears his name.

**Example 1.** The function  $f(x) = \frac{1}{x}$  is convex on  $(0, \infty)$ . Indeed, due to continuity it suffices to prove that  $f$  is *J-convex*, namely,

$$\frac{2}{x+y} \leq \frac{\frac{1}{x} + \frac{1}{y}}{2} \quad (x, y > 0)$$

and this is obviously true.

## 2. Operations with convex functions

We start with an evident assertion.

**Theorem 1.** *If  $f, g : I \rightarrow \mathbb{R}$  are convex and  $\alpha \geq 0$ , then  $f + g$  and  $\alpha f$  are also convex.*  $\square$

The next two simple theorems are stated for the first time in [5].

**Theorem 2.** *Let  $f : I \rightarrow J$  and  $g : J \rightarrow \mathbb{R}$ , where  $J \subset \mathbb{R}$  is also an interval. Then:*

$$\begin{aligned} f \text{ convex, } g \text{ convex and increasing} &\Rightarrow g \circ f \text{ convex;} \\ f \text{ convex, } g \text{ concave and decreasing} &\Rightarrow g \circ f \text{ concave;} \\ f \text{ concave, } g \text{ concave and increasing} &\Rightarrow g \circ f \text{ concave;} \\ f \text{ concave, } g \text{ convex and decreasing} &\Rightarrow g \circ f \text{ convex.} \end{aligned}$$

*Proof.* Let us prove only the first statement. For all  $x, y \in I$ , we have in order

$$\begin{aligned} f(\lambda x + (1 - \lambda)y) &\leq \lambda f(x) + (1 - \lambda)f(y), \\ g(f(\lambda x + (1 - \lambda)y)) &\leq g(\lambda f(x) + (1 - \lambda)f(y)) \\ &\leq \lambda g(f(x)) + (1 - \lambda)g(f(y)). \end{aligned} \quad \square$$

**Theorem 3.** *Let  $f : I \rightarrow J$ ,  $J \subset \mathbb{R}$  with  $J = f(I)$ . Then:*

- (i)  $f^{-1}$  is convex, if  $f$  is convex and decreasing or concave and increasing;
- (ii)  $f^{-1}$  is concave, if  $f$  is convex and increasing or concave and decreasing.

*Proof.* Suppose that  $f$  is convex and increasing. Then  $f^{-1}$  is increasing. Consequently, for all  $x_1, x_2 \in I$  and  $y_1 = f(x_1)$ ,  $y_2 = f(x_2)$ , it follows that

$$f^{-1}(f(\lambda x_1 + (1 - \lambda)x_2)) \leq f^{-1}(\lambda f(x_1) + (1 - \lambda)f(x_2)),$$

that is,

$$\lambda f^{-1}(y_1) + (1 - \lambda)f^{-1}(y_2) \leq f^{-1}(\lambda y_1 + (1 - \lambda)y_2).$$

One similarly proceeds with the remaining statements.  $\square$

As an immediate consequence of Theorem 2 and Example 1, we have

**Theorem 4.** *If  $f : I \rightarrow (0, \infty)$  is concave, then  $\frac{1}{f}$  is convex.*  $\square$

An analogous assertion for a convex function does not hold in general. For example,  $f(x) = x^2 + 1$  is convex on  $\mathbb{R}$ , but  $\frac{1}{f}$  is neither convex nor concave.

### 3. Some important inequalities

In the sequel we will employ the following inequalities:

$$\sqrt[n]{a_1 \dots a_n} \leq \frac{a_1 + \dots + a_n}{n}, \quad a_i \geq 0 \quad (i = 1, \dots, n); \quad (4)$$

$$a^\lambda b^{1-\lambda} \leq \lambda a + (1 - \lambda)b, \quad \lambda \in [0, 1], \quad a, b \geq 0; \quad (5)$$

$$\left(\frac{a+b}{2}\right)^\alpha \leq \frac{a^\alpha + b^\alpha}{2}, \quad \alpha \geq 1, \quad a, b \geq 0. \quad (6)$$

The first inequality is the celebrated inequality between the arithmetic and geometric mean (AM–GM). More than one hundred proofs are known, and one of them, Cauchy's original proof is included in the beautiful book [1].

The second inequality, often called Young's inequality, is a consequence of (4): assuming first that  $\lambda = \frac{m}{n}$  ( $m, n \in \mathbb{N}$ ,  $m \leq n$ ) is rational, we obtain

$$\sqrt[n]{a^m b^{n-m}} \leq \frac{ma + (n-m)b}{n},$$

or equivalently,

$$a^{\frac{m}{n}} b^{1-\frac{m}{n}} \leq \frac{m}{n} a + \left(1 - \frac{m}{n}\right)b. \quad (7)$$

Passing to the limit in (7), we conclude that (5) holds also for irrational  $\lambda \in [0, 1]$ .

Concerning the proof of the third inequality, note that

$$a_1^\lambda b_1^{1-\lambda} + a_2^\lambda b_2^{1-\lambda} \leq (a_1 + a_2)^\lambda (b_1 + b_2)^{1-\lambda} \quad (8)$$

for all positive  $a_i, b_i$  ( $i = 1, 2$ ). Indeed, applying inequality (5), we have

$$\begin{aligned} & \left(\frac{a_1}{a_1 + a_2}\right)^\lambda \left(\frac{b_1}{b_1 + b_2}\right)^{1-\lambda} + \left(\frac{a_2}{a_1 + a_2}\right)^\lambda \left(\frac{b_2}{b_1 + b_2}\right)^{1-\lambda} \\ & \leq \lambda \left(\frac{a_1}{a_1 + a_2}\right) + (1 - \lambda) \left(\frac{b_1}{b_1 + b_2}\right) + \lambda \left(\frac{a_2}{a_1 + a_2}\right) + (1 - \lambda) \left(\frac{b_2}{b_1 + b_2}\right) \\ & = 1. \end{aligned}$$

If we set

$$a_1 = \frac{1}{2} a^\alpha, \quad a_2 = \frac{1}{2} b^\alpha, \quad b_1 = b_2 = \frac{1}{2}, \quad \lambda = \frac{1}{\alpha}$$

in (8), we obtain (6).

In 1888, H. Simon published a paper with the proof of an inequality slightly more general than (6). There he made the remark that I.-J. Bienaymé had stated in 1840 similar results, however, without proofs.

### 4. Convexity of elementary functions

Since elementary functions are continuous on their domain of definition, convexity in this case reduces to  $J$ -convexity. We will see now how one can establish convexity (or concavity) of the most common elementary functions by applying the results from the previous sections.

### 1. Power function

Let  $f(x) = x^n$  and  $n \in \mathbb{N}$ . Then

$$\left(\frac{x+y}{2}\right)^n \leq \frac{x^n + y^n}{2}$$

holds on  $[0, \infty)$  by virtue of (6). Now, it is a simple matter to prove that  $f$  is convex on the whole of  $\mathbb{R}$  for  $n$  even and concave on  $(-\infty, 0]$  for  $n$  odd.

### 2. Root function

Taking into account that  $f(x) = \sqrt[n]{x}$  is the inverse of the power function, Theorem 3 applies.

### 3. Exponential function

Thanks to the AM–GM inequality for  $n = 2$ , we have

$$e^{\frac{x+y}{2}} = \sqrt{e^x e^y} \leq \frac{e^x + e^y}{2},$$

and, consequently,  $\exp$  is convex on  $\mathbb{R}$ .

The function  $f(x) = a^x$  ( $a > 0$ ) is also convex on  $\mathbb{R}$  as being a composition of the increasing convex function  $\exp$  and the convex (linear) function  $x \mapsto (\ln a)x$ .

### 4. Logarithmic function

Again, since  $f(x) = \log_a x$  ( $a > 0$ ,  $a \neq 1$ ) is the inverse of the convex function  $x \mapsto a^x$ , we simply employ Theorem 3 to deduce concavity for  $a > 1$  and convexity for  $0 < a < 1$ .

### 5. Generalized power function

For  $f(x) = x^\alpha$  ( $x > 0$ ) we distinguish the following cases:

(a)  $\alpha \geq 1$ :

The corresponding inequality (2) for  $f$  is exactly (6).

(b)  $0 < \alpha < 1$ :

In this case  $f$  is concave, since  $f^{-1}(x) = x^{1/\alpha}$  is increasing and, according to (a), convex.

(c)  $\alpha < 0$ :

Utilizing convexity of  $x \mapsto \alpha \ln x$ , it follows from  $f(x) = e^{\alpha \ln x}$  that  $f$  is convex.

## 6. Trigonometric functions

- (a) The function  $\sin$  is concave on  $[0, \pi]$ . Indeed, for  $x, y \in [0, \pi]$ , we have  $\sin \frac{x+y}{2} \geq 0$ . Therefore,

$$\sin x + \sin y = 2 \sin \frac{x+y}{2} \cos \frac{x-y}{2} \leq 2 \sin \frac{x+y}{2}.$$

Hence,

$$\sin \frac{x+y}{2} \geq \frac{\sin x + \sin y}{2}.$$

Applying  $\sin(x + \pi) = -\sin x$ , we see that  $\sin$  is convex on  $[\pi, 2\pi]$ .

- (b) From (a) and

$$\cos x = \sin \left( x + \frac{\pi}{2} \right),$$

we conclude that  $\cos$  is convex on  $[\frac{\pi}{2}, \frac{3\pi}{2}]$ , and concave on  $[0, \frac{\pi}{2}]$  and  $[\frac{3\pi}{2}, 2\pi]$ .

- (c)  $\tan$  is convex on  $[0, \frac{\pi}{2})$ . To see this, we utilize concavity of  $\cos$  on  $[0, \frac{\pi}{2})$  and the AM–GM inequality to obtain

$$\sqrt{\cos x \cos y} \leq \frac{\cos x + \cos y}{2} \leq \cos \frac{x+y}{2}.$$

Finally, for  $x, y \in [0, \frac{\pi}{2})$ , it follows that

$$\tan \left( \frac{x+y}{2} \right) = \frac{\sin \frac{x+y}{2}}{\cos \frac{x+y}{2}} = \frac{2 \sin \frac{x+y}{2} \cos \frac{x+y}{2}}{2 \cos^2 \frac{x+y}{2}} \leq \frac{\sin(x+y)}{2 \cos x \cos y} = \frac{\tan x + \tan y}{2}.$$

As an odd function,  $\tan$  is concave on  $(-\frac{\pi}{2}, 0]$ .

## 7. Inverse trigonometric functions

Here the application of Theorem 3 and previous results allow us to deduce intervals of convexity and concavity for the functions  $\arcsin$ ,  $\arccos$ ,  $\arctan$ .

## 8. Hyperbolic functions

- (a) The function  $\sinh$  is convex on  $[0, +\infty)$ . This follows from

$$\sinh x + \sinh y = 2 \sinh \left( \frac{x+y}{2} \right) \cosh \left( \frac{x-y}{2} \right) \geq 2 \sinh \left( \frac{x+y}{2} \right).$$

$\sinh$  is concave on  $(-\infty, 0]$ , since it is an odd function.

- (b) The function  $\cosh$  is convex on  $\mathbb{R}$  being a sum of two convex functions. Note that  $\ln \cosh$  is also convex on  $\mathbb{R}$  by virtue of Theorem 2. Inequality (2) for this function implies that

$$\cosh \left( \frac{x+y}{2} \right) \leq \sqrt{\cosh x \cosh y}. \quad (9)$$

(c) From (9) one infers that for all  $x, y \geq 0$ ,

$$\tanh\left(\frac{x+y}{2}\right) = \frac{2 \sinh \frac{x+y}{2} \cosh \frac{x+y}{2}}{2 \cosh^2 \frac{x+y}{2}} \geq \frac{\sinh(x+y)}{2 \cosh x \cosh y} = \frac{\tanh x + \tanh y}{2},$$

which means that  $\tanh$  is concave on  $[0, \infty)$ . Convexity on  $(-\infty, 0]$  follows from the fact that it is an odd function.

## 5. Applications to inequalities

### 5.1 Henrici's and Ky Fan's inequalities

Two interesting inequalities can be derived by using concavity of  $\tanh$  on  $[0, \infty)$ . First, note that the function  $f(x) = \frac{1}{1+e^x}$  is convex on  $[0, \infty)$ , due to

$$f(x) = \frac{1}{2} - \frac{1}{2} \tanh \frac{x}{2}.$$

Jensen's inequality (3) for  $f$  reduces to the inequality of P. Henrici [4]

$$\frac{n}{1 + \sqrt[n]{x_1 \dots x_n}} \leq \sum_{i=1}^n \frac{1}{1 + x_i} \quad (x_i \geq 1, \quad i = 1, \dots, n).$$

Note further that  $f$  is decreasing. Therefore,

$$f^{-1}(x) = \ln \frac{1-x}{x}$$

is convex on  $(0, \frac{1}{2})$ . Jensen's inequality (3) applied to  $f^{-1}$  leads to the inequality of Ky Fan

$$\frac{\prod_{i=1}^n x_i}{(\sum_{i=1}^n x_i)^n} \leq \frac{\prod_{i=1}^n (1-x_i)}{(\sum_{i=1}^n (1-x_i))^n}, \quad 0 < x_i \leq \frac{1}{2} \quad (i = 1, \dots, n).$$

This shows how closely related the inequalities of Henrici and Ky Fan are.

Interestingly, Henrici's inequality was posed as problem No. 245 in *Elemente der Mathematik* in 1955. On the other hand, Ky Fan's inequality has originally been stated without proof in [2] as an unpublished result of Ky Fan, accompanied with a hint of using mathematical induction.

### 5.2 Several geometric inequalities

Let us denote by  $P$  the area of a triangle with the side lengths  $a, b, c$ , the angles  $\alpha, \beta, \gamma$ , and the height lengths  $h_a, h_b, h_c$ .

Thanks to concavity and positivity of the function  $\sin$  on  $(0, \pi)$ , we conclude by virtue of Theorem 4 that the function  $1/\sin$  is convex on the same interval. Jensen's inequality (3) with  $n = 3$  in this case yields

$$\frac{1}{\sin \frac{\alpha+\beta+\gamma}{3}} \leq \frac{\frac{1}{\sin \alpha} + \frac{1}{\sin \beta} + \frac{1}{\sin \gamma}}{3}.$$



Therefore,

$$\frac{1}{\sin \alpha} + \frac{1}{\sin \beta} + \frac{1}{\sin \gamma} \geq 2\sqrt{3}.$$

Now, from

$$a^2 + b^2 + c^2 \geq ab + bc + ca = 2P \left( \frac{1}{\sin \alpha} + \frac{1}{\sin \beta} + \frac{1}{\sin \gamma} \right),$$

we get Weitzenböck's inequality [9]

$$4\sqrt{3}P \leq a^2 + b^2 + c^2.$$

Note that from

$$6P = ah_a + bh_b + ch_c \leq ab + bc + ca \leq a^2 + b^2 + c^2$$

follows

$$4\sqrt{3}P \leq \frac{2}{\sqrt{3}}(a^2 + b^2 + c^2). \quad (10)$$

Consequently, Weitzenböck's inequality is an improvement of the trivial inequality (10).

Many estimates for  $4\sqrt{3}P$  are known. For example, Pólya and Szegő proved in [6] that

$$4\sqrt{3}P \leq 3\sqrt[3]{a^2b^2c^2}. \quad (11)$$

Having in mind that

$$3\sqrt[3]{a^2b^2c^2} \leq a^2 + b^2 + c^2,$$

we see that the estimate (11) is sharper than Weitzenböck's inequality. Here we show a proof of (11), which partially employs ideas of H. Flanders [3].

The function  $f = \ln \sin$  is a composition of two concave functions with  $\ln$  additionally being increasing. Thus,  $f$  is concave. Jensen's inequality applied to  $-f$  implies

$$\sin \alpha \sin \beta \sin \gamma \leq \frac{3}{8}\sqrt{3},$$

and (11) easily follows from the evident identity

$$P^3 = \frac{1}{8}(abc)^2 \sin \alpha \sin \beta \sin \gamma.$$

We leave as an exercise to prove the inequality (see [7]):

$$\sin \frac{\alpha}{2} \sin \frac{\beta}{2} \sin \frac{\gamma}{2} \leq \frac{1}{8}.$$

**References**

- [1] Aigner, M.; Ziegler, G.M.: *Proof from THE BOOK*. Springer-Verlag, 2003.
- [2] Beckenbach, E.F.; Bellman, R.: *Inequalities*. Springer-Verlag, Berlin, 1961.
- [3] Flanders, H.: Review: Problems and theorems in analysis. *Bull. Amer. Math. Soc.* 84 (1978), 53–62.
- [4] Neue Aufgaben. *Elem. Math.* 10 (1955), 92.
- [5] Jensen, J.L.W.V.: Sur les fonctions convexes et les inégalités entre les valeurs moyennes. *Acta Math.* 30 (1906), 175–193.
- [6] Pólya, G.; Szegő, G.: *Problems and theorems in analysis*. Vol. I. Springer-Verlag, 1972.
- [7] Radó, T.: On mathematical life in Hungary. *Amer. Math. Monthly* 37 (1932), 85–90.
- [8] Roberts, A.W.; Varberg, D.: *Convex Functions*. Academic Press, 1973.
- [9] Weitzenböck, R.: Über eine Ungleichung in der Dreiecksgeometrie. *Math. Z.* 5 (1919), 137–146.

Milan Jovanović and Vladimir Jovanović  
Faculty of Sciences  
University of Banja Luka  
Mladena Stojanovića 2  
51000 Banja Luka, Bosnia and Herzegovina  
e-mail: jaglikaj@yahoo.com  
e-mail: vlajov@blic.net