

**Zeitschrift:** Eclogae Geologicae Helvetiae  
**Herausgeber:** Schweizerische Geologische Gesellschaft  
**Band:** 20 (1926-1927)  
**Heft:** 3

**Artikel:** The Microspheric form of "Orthophragmina" (Discocyclina) Peruviana, Cushman  
**Autor:** Berry, Willard  
**DOI:** <https://doi.org/10.5169/seals-158616>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 28.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

Fig. 43 u. 44. Zwei Metatarsalia, 43 wahrscheinlich das erste.  
 Fig. 45 u. 46. Zwei Phalangen.  
 Fig. 47. Klaue.

### Erklärung der Tafel XV.

Neue Rekonstruktion von *Diplodocus Carnegii*. Zugrunde liegt das von Hatcher u. Holland abgebildete Skelett, aber mit berücksichtigt ist das in Frankfurt a./M. aufgestellte Original-Skelett. 1:40 nat. Grösse. Das als Masstab danebengestellte menschliche Skelett ist 1,75 m hoch.

### Erklärung der Tafel XVI.

Rekonstruktion von *Titanosaurus australis* aus der oberen Kreide Patagoniens nach dem im Museum zu La Plata befindlichen Material, das vom Verfasser bearbeitet worden ist. 1:12 nat. Grösse. Das als Masstab danebengestellte menschliche Skelett ist 1,75 m hoch.

Manuskript eingegangen am 12. Februar 1927.

---

## The Microspheric Form of "Orthophragmina" (Discocyclina) Peruviana, Cushman.<sup>1)</sup>

By E. WILLARD BERRY (Negritos).

With 1 Figur.

---

The megalospheric form of this species was described by J. A. CUSHMAN in 1923 from some material sent him by T. O. BOSWORTH. This material was "from the Lobitos Formation around Negritos".<sup>2)</sup> There is a species of *Orthophragmina*, which places this lot of material definitely in the Upper Eocene.

It has been my good fortune to do some work in the area (*i.e.* around Negritos) and have found the megalospheric form a very good guide fossil in well correlation. Recently I have found the microspheric form of this same species. This small form is scarce but looks much like its larger form and is only found associated with it (the megalospheric form).

Dr. CUSHMAN'S description of the megalospheric form is as follows<sup>3)</sup>: "Test circular, consisting of two portions, the

central raised, umbonate, occupying about one-third of the diameter of the test, surrounded by a thin flange-like portion of equal width; surface coarsely papillate, both the center and the peripheral portion almost equally so; diameter about 6 mm."

In equatorial sections there are to be seen the very narrow rectangular chambers of the "Orthophragmina" type.

The microspheric form is as follows: Test circular, consisting of two portions. A central, slightly raised portion, occupying about one-third of the test (like the larger form). This central portion is surrounded by a thin flange-like portion of about the same width; the surface coarsely papillate, much more so than in the megalospheric form. Diameter about 2 mm.

In the thin section of the equatorial region there are two unequal embryonic chambers. A large one (Fig. 1B) almost enclosing a smaller one (Fig. 1A). This pair of chambers is in turn surrounded by a circlet of chamberlets (Fig. 1C) consisting usually of 20 rectangular chamberlets. These are usually nearly square. Beyond these are the regular bands or rows of much smaller regular thin rectangular chambers (Fig. 1D). In the megalospheric form the chambers of this area are narrow rectangular (*i.e.* longer radially than tangentially) just the reverse of those in the microspheric form.

Diameter of chamber A about 0.168 mm.  
 " " " B " 0.252 mm.




Fig. 1

The scarcity of the microspheric form may be remarked from the fact that I worked in the area using the larger form often for almost two years before finding the microspheric form.

<sup>1</sup>) Published by permission Chief Geologist, International Petroleum Co. Ltd.

<sup>2</sup>) BOSWORTH, T. O. Geol. and Paleon. of N. W. Peru; 1922; p. 136.

<sup>3</sup>) Ibidem, p. 138; pl. XXIV, fig. 3.

