

Zeitschrift: Eclogae Geologicae Helvetiae
Herausgeber: Schweizerische Geologische Gesellschaft
Band: 64 (1971)
Heft: 3

Artikel: Rb-Sr age determinations on biotites and whole rock samples from the Mandi and Chor granites, Himachal Pradesh, India
Autor: Jäger, Emilie / Bhandari, Anil K. / Bhanot, V.B.
DOI: <https://doi.org/10.5169/seals-163993>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 22.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Rb-Sr Age Determinations on Biotites and Whole Rock Samples from the Mandi and Chor Granites, Himachal Pradesh, India

By EMILIE JÄGER¹⁾, ANIL K. BHANDARI²⁾ and V. B. BHANOT³⁾

ABSTRACT

Rb-Sr age determinations on three biotites and four whole rock samples from the Mandi granite, Himachal Pradesh, India, were performed. All the apparent biotite-ages fall in the range 24–31 m.y. pointing towards the intense phase of Himalayan metamorphism evidenced by their parent rocks during that period. The total rocks give an age of 500 ± 100 m.y. suggesting the time of formation of these rocks, which were later remobilized during Himalayan metamorphism.

One sample of biotite separated from a granitic sample from Chor area gave a mineral age of 50 ± 10 m.y.

Introduction

Himalayan granites pose a problem for geologists as regards their age and origin. Opinions differ as to the age and mode of origin of these rocks. Himalayan granites were assigned ages by different workers on the basis of various criteria such as nature of the xenoliths, field relationship with the country rock, petrographical similarities, structural trends, degree of metamorphism, geochronometry, etc.

Based on these evidences, earlier workers like MCMAHON (1897), GRIESBACH (1891), AUDEN (1933) and WADIA (1933) grouped the Himalayan granites into the following age groups:

- I. Late Palaeozoic granites
- II. Post Carboniferous-Early Tertiary granites
- III. Tertiary granites

KRUMMENACHER (1961) and SARKAR et al. (1964) have reported a few K-Ar ages for minerals separated from Himalayan granitic samples, but no Rb-Sr whole rock ages have been published for the Himalayan granites.

In this investigation Rb-Sr age determinations were made on three biotites and four whole rock samples from the Mandi granite. For comparison one biotite from the

¹⁾ Mineralogisch-petrographisches Institut der Universität Bern, Bern, Switzerland.

²⁾ Department of Physics, Panjab University, Chandigarh, India; present address: Geological Survey of India, Lucknow, U.P., India.

³⁾ Department of Physics, Panjab University, Chandigarh, India.




Fig. 1. Sketch map of the Himalayas, location of Fig. 2 is indicated by a black rectangle.




Fig. 2 Geological map of the Mandi and Chor area. Chandigarh is situated 220 km N of Delhi.

Chor granite was also dated. From the biotite analyses the influence of the Himalayan orogeny on these rocks can be studied. The total rock results could reveal the previous history of these rocks.

Geological setting and petrography

According to FUCHS (1967), the Mandi granite forms the front part of the crystalline nappe, other geologists believe that the Mandi granite occurs in the core of an anticline of Chail sediments and that it belongs to the Chail unit and not to the crystalline nappe. In a recent paper FUCHS (FUCHS and GUPTA 1971) also considers the Mandi granite to be part of the Chails. BHANDARI made several traverses through the northern part of this granitic body, along and parallel to Mandi-Kulu road, as well as across the granite body. A description of the observed features is given in the following paragraphs (Fig. 2 and 3).

The granitic body, as a whole, is oblong in shape with its elongation following the regional trend of the metasediments (NNW–SSE). There is no sign of gradation between the metasediments and the granite at the contacts where exposed. These contacts are quite sharp and there is little variation in the grain size of the granite near the contact zone. However, at the eastern contact on the Mandi-Kulu road, there is a complex series of granitic sheets intruding along the foliation planes of the metasediments. Sheets more than a few meters wide are normal porphyritic gneisses, but the smaller sheets tend to be fine grained.

Near the margin, the Mandi granite is a porphyritic gneiss which grades into a coarse-grained nearly non-porphyritic gneiss and then into a medium to-coarse-grained homogeneous granite towards the core. The porphyroblasts of feldspars occurring in the porphyritic gneiss are mostly microcline wrapped around by biotites and are of varying sizes. The foliation – although often indistinct – follows the structural pattern of the country rock. The most prominent direction of elongation of feldspars is NW–SE. The feldspar crystals are often fractured, a fact which points to their having been affected by the Himalayan Orogeny. Next variety of marginal granite is a coarse-grained rock with smaller feldspar crystals and a poorly defined foliation. This unit of the granite is well-jointed and traversed by cross-cutting veins of aplite and pegmatite which do not strictly follow the joint planes. This granite is moderately rich in biotite. Samples SMK 4 and SMK 5 are from this gneissose granite.

A third type of rock is a coarse-grained homogeneous granite consisting mainly of feldspars (plagioclase predominating over K-feldspar), quartz, biotite and muscovite. Muscovite occurs in two generations. Samples PHS1, PHS2, PHS3 and PHS4 were collected from this type.

Near the contact a number of xenoliths of biotite-schist and talc-schist, varying in size from a few centimeters to more than a few meters, were observed in the granite and porphyritic granite.

Microscopically, the central granitic rocks are generally recrystallized and are suggestive of a metamorphism at low temperatures which is indicated by the common occurrence of secondary sericite at the margins of perthite and muscovite. The biotite in these rocks has a “wavy” colour, usually pale brown to pale green, and has inter-

growths of secondary muscovite. Bands of recrystallized quartz and sericite occur between the feldspars and the mica grains and fill the fracture surfaces along which they have been broken.

According to PILGRIM and WEST (1928), the Chor granite (Fig. 2) is a granite of laccolithic nature, intruded into a syncline of Jutogh beds. This syncline and the granite intrusion within it have been overturned by a powerful movement from the NE. The Chor granite is a coarse-grained, usually foliated rock, with abundant phenocrysts of orthoclase, and can be called a porphyritic biotite granite. Inclusions of zircon and apatite are common. Though biotite is the predominating mica, it rarely occurs in large crystals. Muscovite in subordinate amounts is present, along with quartz, sphene and epidote. Sample CA 3 was collected from a place about 1 mile from Nauraghat village in the Chor area.

Analytical data and age results

Table 1 summarizes the analytical data and age results. The total rock analyses are presented in the NICOLAYSEN diagram (Fig. 4).

The young Rb-Sr ages on biotites show the influence of the young Himalayan metamorphism in the Mandi granite. A young Rb-Sr age result was found on the biotite from the Chor granite. This biotite concentrate was not very well separated, resulting in a high content of common Sr and a high error. But there is no doubt that this age is young, too.




Fig. 3. Geological map of the Mandi area, along the Mandi-Kulu road.

Table 1. Analytical data and age results

Sample No.	Rock	Analyzed sample	ppm Rb ⁸⁷	ppm Sr ⁸⁷ rad.	‰ rad.	ppm Sr comm.	Rb ⁸⁷ /Sr ⁸⁶	Sr ⁸⁷ /Sr ⁸⁶	Age in m.y.
PHS1	Mandi granite	Total rock	122	0.907	23.8	42.1	29.802	0.9322	
PHS2	Mandi granite	Total rock	120	0.921	23.9	42.6	28.967	0.9314	
PHS3	Mandi granite	Total rock	112	0.881	16.7	64.2	17.928	0.8502	500 ± 100
PHS4	Mandi granite	Total rock	114	0.866	17.6	58.6	19.986	0.8608	
PHS4	Mandi granite	Biotite	539	0.211	47.0	3.45	1 606.93	1.3379	20.5 ± 0.8
SMK4	Mandi granite	Biotite	249	0.0877	18.3	5.69			
SMK5	Mandi granite	Biotite	397	0.183	19.9	10.7			
CA3	Chor granite	Biotite	255	0.188	9.9	24.7			

Constants: $= 1.47 \times 10^{-11} \text{y}^{-1}$; Rb: $85/87 = 2.591$; Sr: $88/86 = 8.432$, $87/86 = 0.7091$

The age of biotite PHS4 represents the isochron with the total rock, biotite SMK4, SMK5 and CA3 were calculated with the common Sr-values mentioned above.

For total rock analyses 4 samples of 30 kg were collected; but they were not selected according to their Rb/Sr ratio. As Figure 4 shows, the 4 analyzed samples fall on a straight line which is not well defined: it is a 2-point "isochron". This is the reason for the large error limit of both apparent age result and the initial $\text{Sr}^{87}/\text{Sr}^{86}$.

But still, there is no doubt that the Mandi granite gives a pre-Himalayan age. It would be worthwhile to measure the total rock age more precisely by analyzing well selected rock samples. With a precise age result a minimum age for the Chail sedimentation of this area could be given. It would be interesting to prove more precisely that the intensive magmatic phase around 450 m.y., commonly found in Europe, is also present in the Himalayan region.




Fig. 4. Nicolaysen diagram on the total rock analyses from the Mandi granite.

Acknowledgments

We owe our thank to Prof. E. Wenk and Dr. W. Frank for reading the manuscript and for critical discussions. Thanks are also due to Mr. J. K. Bhalla for assistance in separation of minerals and for several useful discussions. We thank Professors I. C. Pande and B. C. Ghosh for permission to use the laboratory facilities available in their departments. One of us (A. K. B.) would like to thank the Council of Scientific and Industrial Research, India, for award of a Research Fellowship. We owe our thanks to the "Schweizerischer Nationalfonds zur Förderung der wissenschaftlichen Forschung" for financing the experimental work.

LITERATURE

AUDEN, J. B. (1933): *On the Age of Certain Himalayan Granites*. Rec. geol. Surv. India 66, 461–471.
 FUCHS, G. (1967): *Zum Bau des Himalaya*. Öst. Akad. Wiss., math.-naturw. Klasse, Denkschr. 113.
 FUCHS, G., and GUPTA, V. J. (1971): *Paleozoic Stratigraphy of Kashmir, Kishtwar and Chamba (Panjab Himalayas)*. Verh. geol. Bundesanst., Wien, in press.
 GANSSER, A. (1964): *Geology of the Himalayas*. Interscience Publishers, London, New York, Sidney.

GRIESBACH, C. L. (1891): *Geology of the Central Himalayas*. Mem. geol. Surv. India 23.

JÄGER, E. (1962): *Rb-Sr Age Determinations on Micas and Total Rocks from the Alps*. J. geophys. Res. 67, 5293.

KRUMMENACHER, D. (1961): *Déterminations d'âge isotopique faites sur quelques roches de l'Himalaya du Nepal par la méthode potassium-argon*. Schweiz. miner. petrogr. Mitt. 41, 273.

McMAHON, C. A. (1897): *Notes on the Age and Structure of the Gneissose-Granite of the Himalayas with Reference to Mr. Middlemiss's Memoir on the Geology of Hazara*. Geol. Mag., p. 304-313.

NICOLAYSEN, L. O. (1961): *Graphic Interpretation of Discordant Age Measurements on Metamorphic Rocks*. N.Y. Acad. Sci. 91, 198-206.

PILGRIM, G. E., and WEST, W. D. (1928): *Structure of the Simla Himalayas*. Mem. geol. Surv. India 53, 52-56.

SARKAR, S. N., POLKANOV, E. K., and GERLING, E. K. (1964): *Geochronology of the Precambrian of Peninsular India: A Synopsis*. Sci. Cult., p. 527-537.

WADIA, D. N. (1933): *Notes on the Geology of Nanga Parbat (Mt. Domir adjoining portions of Chilas, Gilgit dist. Kashmir)*. Rec. geol. Surv. India 66, 212-234.

