

**Zeitschrift:** Eclogae Geologicae Helvetiae  
**Herausgeber:** Schweizerische Geologische Gesellschaft  
**Band:** 64 (1971)  
**Heft:** 3

**Artikel:** Die Gattungen Hysterocheras Spath und Mortonicerias Meek (Ammonoidea) aus den Anden Venezuelas

**Autor:** Renz, Otto

### **Inhaltsverzeichnis**

**DOI:** <https://doi.org/10.5169/seals-163996>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 28.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## ABSTRACT

The Mojsisovicsiinae and Engonoceratidae from the uppermost layer of the La Puya Member have been described in two earlier publications (O. RENZ 1968, 1970). As a continuation the Brancoceratinae and Mortoniceratinae from the same layer are dealt with in the present article.

The La Puya Fauna is of special interest, as it derives from a single layer, reaching a thickness of only about 1 m. It forms the termination of a thick sequence consisting predominantly of carbonate rocks without cephalopods. As reworking or condensation might be excluded, this fauna represents a cross section during a short time interval within the Upper Albian. It apparently has been deposited within the passage zone from the southern rim of the slowly sinking Maracaibo Platform towards a deeper pelagic basin.

A correlation of the fauna with one of the European ammonite zones could not be achieved with the Mojsisovicsiinae and Engonoceratidae as these forms are found only occasionally in the classical sections of England and France. On the contrary most of the species here described of the genera *Hysterocheras* and *Mortoniceras* occur abundantly in the sections of Europe (table 1).

The absence of *Hysterocheras varicosum* (J. de C. Sow.) within the La Puya Limestone suggests a younger age for its fauna than the *varicosum* Zone in Europe. On the other hand *Hysterocheras bucklandi* (SPATH), occurring abundantly in Venezuela, remains restricted to the *auritus* Zone in England and France.

With one exception, all *Mortoniceras* from the La Puya Limestone are characterized by only two rows of tubercles. *Mortoniceras (M.) kiliani* (LASSWITZ), restricted to the *aequatoreale*- and the *substuderi* Zone in England and France, has so far not been detected in Venezuela. On the contrary *M. (M.) pricei intermedium* SPATH, which is known in England from the *auritus* Zone only, occurs rather abundantly in the La Puya Limestone. Moreover the occurrence of the subgenus *Deiradoceras* in the lower part of the Upper Albian (*orbigny*- and *varicosum* Zone) in England as well as in France has to be remarked. This confirms the assumption that the La Puya Fauna, containing abundant *Deiradoceras*, is not younger than the *auritus* Zone.

The above observations point to a correlation of the La Puya Fauna with the *auritus* Zone in England and with the lower part of the „sous-zone B” in France.

Amongst 11 species and subspecies of *Hysterocheras*, reaching 21% of the fauna, 4 are known from England and 5 from France. All forms rather suggest a connection to Europe.

The genus *Mortoniceras* is represented by 10 species and 1 subspecies, amounting only to 7% of the fauna. The majority of the species available point to a relation to West and South Africa, as well as to Madagascar.

## INHALT

1. Einleitung . . . . .	570
2. Anteil der Gattungen <i>Hysterocheras</i> und <i>Mortoniceras</i> an der Gesamtfauna aus dem La Puya-Member . . . . .	571
3. Alter der Fauna . . . . .	572
4. Paläontologischer Teil . . . . .	577
a) Subfamilie Brancoceratinae SPATH 1933 . . . . .	577
Gattung <i>Hysterocheras</i> HYATT 1900 . . . . .	577
b) Subfamilie Mortoniceratinae SPATH 1925 . . . . .	592
Gattung <i>Prohysterocheras</i> SPATH 1921 . . . . .	592
Gattung <i>Mortoniceras</i> MEEK 1876 . . . . .	594

## 1. Einleitung

In zwei früheren Mitteilungen (RENZ 1968 und 1970) wurden die Vertreter der Gattungen *Oxytropidoceras*, *Parengonoceras* und *Knemiceras* aus dem La Puya-Kalk in den venezolanischen Anden beschrieben. In vorliegender Arbeit sollen weitere Cephalopoden aus der Familie Brancoceratidae SPATH 1933 zur Darstellung gelangen. Es