

Zeitschrift:	Eclogae Geologicae Helvetiae
Herausgeber:	Schweizerische Geologische Gesellschaft
Band:	72 (1979)
Heft:	1
Artikel:	The Upper Jurassic coccoliths from the Haddenham and Gamlingay boreholes (Cambridgeshire, England)
Autor:	Medd, Alan W.
Kapitel:	Details of the assemblages
DOI:	https://doi.org/10.5169/seals-164830

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

assumption. The number of specimens also varies within the stratigraphical interval studied; there is an increase in such numbers in the lower part of the Kimmeridge Clay, to a maximum number per unit mass of rock found for the whole of the Jurassic.

The methods of study are as stated in MEDD (1971) but a Zeiss photomicroscope II has been used for the routine optical investigations. The Nomarski interference phase contrast accessory is occasionally of help in the study of coccoliths. A TV fast scanning accessory on the scanning electron microscope has also proved to be of value for a detailed study of coccolith strew mounts; the instrument used is a Cambridge Instruments Co. Stereoscan Mark 2A, which is located at the Institute of Geological Sciences, London. The technique of NOËL (1973, p. 98) in examining the rock surface with an SEM has also been used but without her remarkable successes. Other Kimmeridgian samples examined, however, have provided excellent results, with abundant assemblages and coccospores seen.

The problem of sample contamination is considerably diminished using bore-hole material and none of the problems associated with surface collected material (MEDD 1971, p. 823) was encountered. The problem of reworking of older coccoliths into the younger nannofloras has not been overcome as to its recognition, as opposed to these species having an extended range. It is difficult to envisage a method of erosion of the older sediments, which would preserve the coccoliths in such numbers as have been found at some levels in these boreholes. The relative proportions of the species are also different from that encountered in the older material. These suspect species appear most often in samples, which are already rich in numbers of indigenous species (i.e., not found in older sediments).

Figured or cited material prefixed with the letters "EM" are of micrographs taken with a Philips 75B transmission electron microscope at the Sedimentology Research Laboratory, University of Reading, and the material is lodged at that laboratory. If prefixed with the letters "SEM", the material was examined using the scanning electron microscope at IGS, London, where the material is now housed. If prefixed by the letters "MPK", the optical strew slides are stored in the MPK collection of type and figured specimens at the Institute of Geological Sciences, Leeds.

Details of the assemblages

The dominant species in all of the samples studied is *Ellipsagelosphaera britannica* (STRADNER) and particularly the small form of it. The only other species found, which occasionally constitute more than 25% of the nannoflora are: *Tetralithus gothicus* DEFLANDRE, *Palaeopontosphaera dubia* NOËL, *Zeugrhabdotus erectus* (DEFLANDRE) and small tremalith-like forms. The remainder of the species recorded can be found in varying degrees of abundance, but with no single species comprising more than 10% of the nannoflora.

Stratigraphical distribution

The distribution of the nannoflora of the Haddenham and Gamlingay boreholes are summarized in Tables 2 and 3, which record the species in the order of their