

Zeitschrift: Eclogae Geologicae Helvetiae
Herausgeber: Schweizerische Geologische Gesellschaft
Band: 77 (1984)
Heft: 3

Artikel: A Middle Jurassic-Early Cretaceous low-latitude radiolarian zonation based on unitary associations and age of Tethyan radiolarites
Autor: Baumgartner, Peter O.

Inhaltsverzeichnis

DOI: <https://doi.org/10.5169/seals-165530>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 31.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

CONTENTS

1. Introduction.....	731
2. Radiolarian biochronology	733
3. Comparison to other zonations and chronostratigraphic calibration	736
3.1 Comparison to earlier zonations	736
3.2 Chronostratigraphic calibration and distribution of Unitary Associations	741
4. Significance of dating radiolarites and conclusions	745
4.1 Chronostratigraphy: correlation of Atlantic and Tethys and timing of Middle–Late Jurassic siliceous sedimentation	745
4.2 Paleoceanographic conclusions.....	749
4.3 Radiolarian faunal changes and provincialism related to paleoceanography	750
4.4 Final conclusions and perspectives.....	751
5. Systematic paleontology.....	752
Explanatory notes.....	752
Alphabetic listing of genera and species	753
6. Locality descriptions.....	792
6.1 Introduction.....	792
6.2 Geographic/paleogeographic overview of the studied localities.....	793
6.3 Listing of localities included in the database	794
Acknowledgments.....	800
References	800
Appendix: Database.....	808

1. Introduction

Much of the present state of Mesozoic radiolarian paleontology and biostratigraphy is the result of the past ten years. After a period of active study of Mesozoic radiolarians in Europe at the turn of the century (RÜST 1885, 1898, PARONA 1890, SQUINABOL 1914, etc.) the interest declined and the biostratigraphic usefulness of radiolarians was questioned.

A number of favorable circumstances revitalized the field in the early seventies: The Deep-Sea Drilling Project was coring Cenozoic and Mesozoic sediments in the oceans which furnished well preserved radiolarian assemblages greatly stimulating biostratigraphic work on this group, as it did for other fossil groups. Radiolarian biostratigraphy, first worked out for the Cenozoic (summary in RIEDEL & SANFILIPPO 1978), rapidly was extended to the Cretaceous (FOREMAN 1971, 1973, 1975, 1978; MOORE 1973, RIEDEL & SANFILIPPO 1974, SCHAAF 1981). The use of hydrofluoric acid to extract fossils from siliceous rocks, known for a long time to palynologists (LEJEUNE 1936) was successfully applied to radiolarians (DUMITRICA 1970, PESSAGNO & NEWPORT 1972) and allowed the observation of isolated forms also from highly lithified siliceous limestones and cherts, leading to systematic work and first Late Jurassic–Cretaceous zonations mainly derived from land-based samples from California (PESSAGNO 1971, 1972, 1973, 1976, 1977a, b). Meanwhile, the Scanning Electron Microscope (SEM) began to be regularly used by micropaleontologists. It produced accurate illustrations even of internally recrystallized or opaque fossil material, inappropriate for transmitted light microscopy.

Only recently, Mesozoic radiolarian biostratigraphy became revitalized in the European area (see DE WEVER et al. 1979) and resulted in first zonations for Tethyan radio-