

Plates

Autor(en): **[s.n.]**

Objekttyp: **Appendix**

Zeitschrift: **Eclogae Geologicae Helvetiae**

Band (Jahr): **77 (1984)**

Heft 3

PDF erstellt am: **15.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

<http://www.e-periodica.ch>

Plate explanations

The illustrations are in accordance with the alphabetic listing of genera and species arranged in alphabetical order, where space on the plates permitted.

Each illustrated specimen is designated with two numbers of the following format: Sample number/year/SEM-negative number, C-number. The sample number refers to locality descriptions and Plate 12. Year and SEM-number refer to the collection of negatives stored at the *Labor für Raster-Elektronenmikroskopie*, Geological Institute, University of Basel. Negative numbers larger than 81/9000 and 82/9000 refer to negatives taken at the Scripps Institution of Oceanography and are stored in the author's collection.

C-numbers refer to the collection of all illustrated material deposited at the *Naturhistorisches Museum Basel, Switzerland*.

Magnifications are indicated for each illustration, they are standardized where possible to allow visual comparisons.

Plate 1

Scanning electron micrographs of Middle Jurassic to Early Cretaceous siliceous (si) and pyritized (py) Radiolaria from Blake-Bahama Basin (DSDP Site 534), Lombardy (POB 1205, 1330, 1341), Romania (MO, V), Greece (POB 899) and Japan (IN 7) (see locality descriptions).

- Fig. 1–2 *Acaeniotyle diaphorogona* FOREMAN, s.l.
 (data 59, range 77, pob 90, rk –), 1: 534A-106-1-29/81/9028, C 35739, py, ×100. 2: 534A-106-1-29/81/9002, C 35740, py, ×100.
- Fig. 3–4 *Acaeniotyle diaphorogona dentata* BAUMGARTNER, n. subsp.
 (data 94, range 99, pob 281, rk –), 3: holotype POB 1205/79/5254, C 35741, si, ×100; 4: paratype MO 22/79/4112, C 35742, broken specimen, note triradiate base of internal beam connecting to (lacking) medullary shell; external spines have no inward continuation as for all *Acaeniotyle*. py, ×100.
- Fig. 5 *Acaeniotyle umbilicata* (RÜST)
 (data 80, range 88, pob 92, rk 18), MO 46/79/4161, C 35743, py, ×100.
- Fig. 6 *Acanthocircus suboblongus* (YAO)
 (data 24, range 22, pob 85, rk 41), POB 899/78/6123, C 35744, si, ×75.
- Fig. 7 *Acanthocircus dicranacanthos* (SQUINABOL)
 (data 82, range 86, pob 87, rk 17), MO 46/79/3095, C 35745, py, ×75.
- Fig. 8–10 *Alievum helenaе* SCHAAF
 (data 104, range 103, pob 228, rk 20), 8, 10: V 34/80/2798, C 35746, py, 8, ×100, 10, ×165; 9: POB 1330/81/9086, C 35747, si, ×100.
- Fig. 11–12 *Andromeda podbielensis* (OZVOLDOVA)
 (data 16, range 43, pob 8, rk 87), 10: 534A-126-2-125/81/9142, C 35748, py, ×100; 11: 534A-125-5-111/81/9141, C 35749, py, ×100.
- Fig. 13–15 *Andromeda praepodbielensis* BAUMGARTNER n. sp.
 (data 3, range 2, pob 6, rk –), 13, 15: holotype POB 1341/81/2978, C 35750, si, ×100; 14: paratype IN 7/79/4431, C 35751, si, ×100.
- Fig. 16–18 *Andromeda praecrassa* BAUMGARTNER n. sp.
 (data 10, range 5, pob 7, rk –), 16: paratype POB 1341/81/2880, si, ×100; 17: holotype POB 1341/81/2975, C 35752, si, ×100; 18: paratype 534A-126-2-125/81/9143, C 35753, segmental divisions are marked with arrows, py, ×100.




Plate 2

Scanning electron micrographs of Middle Jurassic to Early Cretaceous siliceous (si) and pyritized (py) Radiolaria from Blake-Bahama Basin (DSDP Site 534), Lombardy (POB 1205, 1330, 1341), Romania (MO, V), Greece (POB 28, 1262) and western Switzerland (POB 1134) (see locality descriptions).

- Fig. 1–3** *Angulobracchia (?) portmanni* BAUMGARTNER n.sp.
(data 98, range 97, pob 285, rk –), 1: holotype POB 1330/81/9091, C 35754, si, ×100; 2: paratype MO 46/79/3121, C 35755, py, ×100; 3: paratype POB 1205/79/5741, C 35756, si, ×100.
- Fig. 4** *Angulobracchia purisimaensis* (PESSAGNO)
(data 67, range 57, pob 144, rk 42), POB 28/78/3762, C 34809, si, ×75.
- Fig. 5–6** *Archaeodictyomitra apiara* (RÜST)
(data 75, range 82, pob 263, rk 14), 5: MO 22/79/4101, C 35757, py, ×150; 6: 534A-81-2-64/81/9118, C 35758, py, ×150.
- Fig. 7–8** *Archaeodictyomitra excellens* (TAN SIN HOK)
(data 100, range 102, pob 287, rk –), 7: MO 46/79/4292, C 35759, py, ×150; 8: 534A-81-2-3/81/9101, C 35760, py, ×150.
- Fig. 9–13** *Archaeohagiastrum munitum* BAUMGARTNER n.gen. n.sp.
(data 92, range 40, pob 271, rk –), 9: holotype 534A-125-5-111/81/9140, C 35761, py, ×150; 10, 13: paratype 534A-126-2-125/81/9175, C 35762, py, 10: ×150, 13: ×500; 11–12: 534A-126-2-125/81/9151, C 35763, lateral view of a specimen with broken spine showing three primary canals arranged around primary beam and six external beams, note also highly raised nodes of central area, typical for this species; py, 11: ×150, 12: ×500.
- Fig. 14–15** *Bernoullius cristatus* BAUMGARTNER n.gen. n.sp.
(data 39, range 39, pob 221, rk 109), 14: holotype 534A-125-5-72/82/9197, C 35764, py, ×100; 15: paratype 534A-125-5-72/81/9198, C 35765, py, ×100.
- Fig. 16** *Bernoullius dicera* (BAUMGARTNER)
(data 35, range 56, pob 223, rk 69), 534A-125-5-72/81/9200, C 35766, py, ×100.
- Fig. 17–18** *Cecrops septemporatus* (PARONA)
(data 110, range 108, pob 229, rk 24), 17: POB 1134/80/2168, C 35767, py, ×100; 18: MO 46/79/4278, C 35768, py, ×100.
- Fig. 19** *Diboloachras chandrika* KOCHER
(data 55, range 75, pob 265, rk 43), 534A-106-1-29/81/9039, C 35769, py, ×100.
- Fig. 20** *Diacanthocapsa normalis* YAO
(data 34, range 10, pob 54, rk –), POB 1262/80/3961, C 35770, si, ×250.
- Fig. 21** *Ditrabs sansalvadorensis* (PESSAGNO)
(data 103, range 96, pob 227, rk 21), MO 46a'/81/0962, C 35771, py, ×100.




Plate 3

Scanning electron micrographs of Middle Jurassic to Early Cretaceous siliceous (si) and pyritized (py) Radiolaria from Blake-Bahama Basin (DSDP Site 534), Lombardy (POB 1341), Greece (POB 899, 986, 1262) (see locality descriptions)

- Fig. 1 *Emiluvia hopsoni* PESSAGNO
 (data 74, range 69, pob 225), POB 899/79/1656, C 35772, si, ×100.
- Fig. 2 *Emiluvia sedecimporata elegans* (WISNIEWSKI)
 (data 40, range 18, pob 216, rk –), POB 986/78/8107, C 35773, si, ×100.
- Fig. 3 *Emiluvia pessagnoi* FOREMAN s.l.
 (data 71, range 71, pob 226, rk 36), POB 986/78/8201, C 35774, si, ×100.
- Fig. 4, 7 *Emiluvia sedecimporata salensis* PESSAGNO
 (data 33, range 50, pob 215, rk 44 & 45), 4: 534A-126-2-125/81/9167, C 35775, py, ×100; 7:
 POB 899/78/8204, C 35776, si, ×100.
- Fig. 5 *Emiluvia orea* BAUMGARTNER
 (data 60, range 81, pob 224, rk 63), topotype POB 899/78/6106, C 35777, si, ×75.
- Fig. 6, 8–9, 11, 12 *Emiluvia premygii* BAUMGARTNER n. sp.
 (data 19, range 14, pob 210, rk 88), 6: paratype 534A-124-1-52/81/2423, C 35778, py, ×100;
 8, 11–12: holotype 534A-124-1-52/81/2424, C 35779, py, note perfect preservation in pyrite,
 showing the smooth surface of bars and spines and the knobby surface of nodes known
 from recent opaline polycystins, 8: ×100, 11: ×250, 12: ×1000; 9: paratype POB 899/79/
 1654, C 35780, si, ×100.
- Fig. 10 *Emiluvia (?)* sp.P.
 (data 41, range 59, pob 219, rk 90), 534A-125-2-36/81/1397, C 35781, py, ×150.
- Fig. 13–16 Eucyrtid gen. et sp. indet.
 (data 63, range 7, pob 74, rk –), 13–14: POB 1341/81/2958, C 35782, si, 13: ×250, 14: ×100;
 15: POB 1263/80/3787, C 35783, si, ×100, 16: POB 1341/81/2960, C 35784, si, ×100.




Plate 4

Scanning electron micrographs of Middle Jurassic to Early Cretaceous siliceous (si) and pyritized (py) Radiolaria from Blake-Bahama Basin (DSDP Site 534) and Greece (POB 325, 899) (see locality descriptions).

- Fig. 1-3 *Eucyrtidiellum ptyctum* (RIEDEL & SANFILIPPO)
 (data 56, range 66, pob 27, rk 46 [pars]), 1: POB 325/80/3798, C 35785, si, $\times 250$; 2: 534A-122-1-43/81/2217, C 35786, py, $\times 250$; 3: 534A-122-1-43/81/2212, C 35787, py, $\times 250$.
- Fig. 4-5 *Eucyrtidiellum pustulatum* BAUMGARTNER n. gen. n. sp.
 (data 91, range 44, pob 13, rk -), 4: holotype 534A-124-1-52/81/2428, C 35788, py, $\times 250$; 5: paratype 534A-124-1-52/81/2429, C 35789, py, $\times 250$.
- Fig. 6 *Eucyrtidiellum unumaensis* (YAO)
 (data 17, range 12, pob 12, rk 89), 534A-126-2-125/81/9144, C 35790, py, $\times 250$.
- Fig. 7 *Gorgansium pulchrum* (KOCHE)
 (data 11, range 28, pob 76, rk 105), 534A-125-2-36/81/1393, C 35791, py, $\times 250$.
- Fig. 8-9 Hagiastrid sp. A.
 (data 8, range 41, pob 153, rk 107 and 108), 534A-124-1-52/81/2795, C 35792, py, fragment!
 8: $\times 500$, 9: $\times 150$.
- Fig. 10-11 *Haliodictya (?) hojnosti* RIEDEL & SANFILIPPO
 (data 86, range -, pob 254, rk 3), 10: 534A-124-1-52/81/2672, C 35793, py, $\times 150$; 11: POB 899/78/6147, C 35794, si, $\times 200$.
- Fig. 12 *Higumastra* sp. aff. *H. inflata* BAUMGARTNER
 (data 66, range 15, pob 107, rk 47), 534A-126-2-125/81/9183, C 35795, py, $\times 150$.
- Fig. 13 *Higumastra imbricata* (OZVOLDOVA)
 (data 13, range 29, pob 110, rk 92), 534A-125-5-72/81/9210, C 35796, py, $\times 150$.
- Fig. 14 *Holocryptocanium barbui* DUMITRICA
 (data 108, range 106, pob 292, rk -), 534A-81-2-64/81/9119, C 35797, py, $\times 150$.
- Fig. 15 *Homoeoparonaella argolidensis* BAUMGARTNER
 (data 43, range 37, pob 103, rk 30), topotype POB 899/78/6201, C 35798, si, $\times 75$.
- Fig. 16 *Homoeoparonaella elegans* (PESSAGNO)
 (data 65, range 63, pob 104, rk 48), 534A-124-1-52/81/2417, C 35799, py, $\times 75$.
- Fig. 17 *Homoeoparonaella giganthea* BAUMGARTNER
 (data 70, range 68, pob 105, rk 37), topotype POB 899/78/6216, C 35800, si, $\times 75$.




Plate 5

Scanning electron micrographs of Middle Jurassic to Early Cretaceous siliceous (si) and pyritized (py) Radiolaria from Blake-Bahama Basin (DSDP Site 534), Lombardy (POB 1205), Greece (POB 28, 899, 986), western Switzerland (POB 1134) and Japan (IN 7) (see locality descriptions).

- Fig. 1–2** *Hsuum brevicostatum* (OZVOLDOVA)
 (data 23, range , pob 181, rk 49), 1: 534A-125-3-29/81/2440, C 35801, py, $\times 150$ 2: POB 28/79/0360 C 35802, si, $\times 150$.
- Fig. 3–4** *Hsuum maxwelli* PESSAGNO group.
 (data 47, range 42, pob 180, rk 93), 3: POB 28/78/3413, C 35803, si, $\times 150$; 4: 534A-122-1-43/81/2274, C 35804, py, $\times 150$.
- Fig. 5–7** *Guexella nudata* (KOCHE)
 (data 7, range 27, pob 61, rk 106), 534A-124-1-52/81/2667, C 35805, py, 5: $\times 300$, 6: $\times 150$, 7: $\times 450$.
- Fig. 8, 22** *Mirifusus guadalupensis* PESSAGNO
 (data 37, range 55, pob 160, rk 50), 8: POB 28 /78/3587, C 35806, si, $\times 75$; 22: POB 899/78/6261, C 35807, detail of inflated median part of test showing irregular outer layer, si, $\times 250$.
- Fig. 9, 15** *Mirifusus chenodes* (RENZ)
 (data 77, range 80, pob 162, rk –), POB 899/79/1642, C 35808, si, note irregular outer layer of branching diagonal bars covering inner layer of small uniform pores in 4–5 rows per segment, 9: $\times 75$, 15: $\times 250$.
- Fig. 10, 18** *Mirifusus mediociliatus baileyi* PESSAGNO
 (data 76, range 67, pob 161, rk 4 [pars]), 10: 534A-106-1-29/81/9053, C 35809, py, note slender conical proximal portion of test (above arrow) with several, well-defined segmental divisions, $\times 75$; 18: POB 986/78/8183, C 34867, si, detail of inflated median portion of test with regular triangular pore frames, two rows of pores per segment and no covering outer layer, $\times 250$.
- Fig. 11, 14** *Mirifusus mediociliatus minor* BAUMGARTNER n. subsp.
 (data 99, range 90, pob 286, rk 4 [pars]), 11: holotype POB 1205/79/5038, C 35810, si, note short, blunt proximal conical portion of test (above arrow) with only one externally visible segmental division, $\times 75$; 14: POB 1134/80/2182, C 35811, py, proximal conical portion of test shows outer layer entirely covering segmental divisions, $\times 75$.
- Fig. 12, 16–17, 20–21** *Mirifusus fragilis* BAUMGARTNER n. sp.
 (data 14, range 9, pob 159, rk –), 12, 17, 20: holotype IN 7/79/4419, C 35812, si, note almost complete absence of outer layer, 12: $\times 75$, 17: $\times 225$, 20: $\times 250$; 16: paratype 534A-126-2-125/81/9158, C 35813, py, note well-developed outer layer of diagonal bars, thin circumferential ridges, well visible inner layer of three rows of circular pores per segment, $\times 250$; 21: 534A-125-3-29, C 35814, py, fragment of median inflated portion with weakly developed outer layer, thin circumferential ridges, $\times 250$.
- Fig. 13, 19** *Mirifusus mediociliatus mediociliatus* (RÜST)
 (data 76, range 67, pob 161, rk 4 [pars]), 13: POB 28/78/3443, C 35815, si; 19: POB 899/78/6696, C 34866, si, detail of inflated median portion of test showing regular circular pore frames in two rows per segment and disappearing outer layer in upper part of Figure, $\times 250$.




Plate 6

Scanning electron micrographs of Middle Jurassic to Early Cretaceous siliceous (si) and pyritized (py) Radiolaria from Blake-Bahama Basin (DSDP Sites 5, 534), Lombardy (POB 1205, 1330) and Greece (POB 28, 783, 899, 986) (see locality descriptions)

- Fig. 1–2, 5** *Monotrabs plenoides* BAUMGARTNER n. gen. n. sp.
(data 42, range 54, pob 152, rk 91), holotype 534A-124-1-52/81/2686, C 35816, py; 2, 5: note long lateral spines and possible hagiastrid structure composed of beams and bars; 1: ×150, 2: ×250, 5: ×500.
- Fig. 3** *Napora deweeveri* BAUMGARTNER
(data 46, range 62, pob 35, rk 95), topotype POB 899/78/6462, C 35817, si, ×150.
- Fig. 4** *Napora bukryi* PESSAGNO
(data 73, range 61, pob 34, rk 31), POB 899/78/6456, C 35818, si, ×150.
- Fig. 6** *Napora lospensis* PESSAGNO
(data 72, range 76, pob 36, rk 32), POB 783/79/0105, C 35819, si, ×150.
- Fig. 7–9** *Obesacapsula rusconensis* BAUMGARTNER n. sp.
(data 95, range 100, pob 282, rk –), 7: paratype POB 1205/79/5039, C 35820, si, ×75; 8: holotype POB 1205/80/2996, C 35821, si, ×75; 9: paratype 534A-89-2-47/81/9060, C 35822, si, ×75.
- Fig. 11–12** *Napora pyramidalis* BAUMGARTNER n. sp.
(data 12, range 11, pob 33, rk 104), 11: holotype 534A-124-1-52/81/2704, C 35823, py, ×250; 12: paratype 534A-124-1-52/81/2656, C 35824, py, ×250.
- Fig. 13** *Obesacapsula rotunda* (HINDE)
(data 83, range 95, pob 202, rk 16), 5A-7-1/79/4232, C 35825, si, ×75.
- Fig. 14–15** *Pantanellium (?) berriasanum* BAUMGARTNER n. sp.
(data 93, range 92, pob 280, rk –), 14: holotype POB 1205/79/5265, C 35826, si, ×150; 15: paratype POB 1330/81/9085, C 35827, si, ×150.
- Fig. 16** *Paronaella bandyi* PESSAGNO
(data 58, range 21, pob 135, rk 51), POB 899/78/6218, C 35828, si, ×75.
- Fig. 17** *Paronaella broennimanni* PESSAGNO
(data 53, range 73, pob 137, rk 71), POB 28/78/3773, C 34792, si, ×75.
- Fig. 18** *Formanella diamphidia* (FOREMAN)
(data 79, range 85, pob 112, rk 13), POB 28/78/3811, C 35829, si, ×75.
- Fig. 19** *Formanella hipposidericus* (FOREMAN)
(data 78, range 83, pob 111, rk 12), POB 986/78/8152, C 34726, si, ×75.
- Fig. 20** *Paronaella kotura* BAUMGARTNER
(data 48, range 64, pob 140, rk 85), topotype POB 899/79/6217, C 35830, si, ×75.
- Fig. 21** *Paronaella mulleri* PESSAGNO
(data 38, range 32, pob 139, rk 96), POB 899/78/6229, C 35831, si, ×75.




Plate 7

Scanning electron micrographs of Middle Jurassic to Early Cretaceous siliceous (si) and pyritized (py) Radiolaria from Blake-Bahama Basin (DSDP Site 534), Lombardy (POB 1205) Greece (POB 28, 284, 899, 986), Sicily (S 4) and California (NSF 907) (see locality descriptions)

- Fig. 1 *Parvingula cosmoconica* (FOREMAN)
 (data 102, range 94, pob 255, rk 22), 534A-81-2-64/81/9111, C 35831, py, $\times 150$.
- Fig. 2–4 *Parvingula dhimenaensis* BAUMGARTNER n. sp.
 (data 90, range 33, pob 197, rk –), 2–3: holotype POB 284/79/0079, C 35833, si, 2: $\times 150$, 3: $\times 250$; 4: A-125-5-72/81/9214, C 35834, py, $\times 150$.
- Fig. 5–6 *Perispyridium ordinarium* (PESSAGNO)
 (data 31, range 48, pob 100, rk 53), 5: POB 986/78/8147, C 35835, si, $\times 100$; 6: 534A-124-1-52-81/2430, C 35836, py, $\times 100$.
- Fig. 7 *Podbursa helvetica* (RÜST)
 (data 18, range 13, pob 169, rk 98), POB 28/78/3551, C 35837, si, $\times 100$.
- Fig. 8 *Podbursa spinosa* (OZVOLDOVA)
 (data 64, range 78, pob 230, rk 54), S 4/79/4721, C 35838, si, $\times 100$.
- Fig. 9–10 *Podocapsa amphitreptera* FOREMAN
 (data 69, range 84, pob 171, rk 38), 9: 534A-106-1-29/81/9009, C 35839, py, $\times 100$; 10: POB 1205/80/2868, C 35840, si, $\times 100$.
- Fig. 11–14 *Praeconocaryomma* (?) *hexacubica* BAUMGARTNER n. sp.
 (data 87, range 31, pob 244, rk –), 11: holotype 534A-126-2-125/81/9154, C 35841, py, $\times 150$; 12: paratype 534A-126-2-125/81/9153, C 35842, py, spines supporting medullary shell are attached in center of squares of cortical shell, $\times 150$; 13: 534A-126-2-125/81/9203, C 35843, py, note characteristic hexagonal pore arrangement, $\times 265$; 14: paratype 534A-125-3-60/81/2451, C 35844, py, morphotype without spines, $\times 150$.
- Fig. 15 *Protunuma costata* (HEITZER)
 (data 21, range 35, pob 232, rk 67), 534A-106-1-29, C 35845, py, $\times 150$.
- Fig. 16 *Pseudocrucella adriani* BAUMGARTNER
 (data 52, range 34, pob 129, rk 72), topotype POB 899/78/6206, C 35846, si, $\times 75$.
- Fig. 17 *Pseudocrucella sanfilippoae* (PESSAGNO)
 (data 51, range 58, pob 126, rk 73), topotype NSF 907/79/1695, C 35847, si, $\times 75$.




Plate 8

Scanning electron micrographs of Middle Jurassic to Early Cretaceous siliceous (si) and pyritized (py) Radiolaria from Blake-Bahama Basin (DSDP Site 534), Lombardy (POB 1205), Greece (POB 899, 986, 1263), western Switzerland (POB 1134) and Romania (MO) (see locality descriptions).

- Fig. 1 *Pseudodictyomitra carpatica* (LOZNYAK)
 (data 107, range 105, pob 293, rk -), 534A-81-2-64/81/9121, C 35848, py, $\times 150$.
- Fig. 2, 7–8, 11 *Pseudodictyomitra depressa* BAUMGARTNER n. sp.
 (data 97, range 101, pob 284, rk -), 2, 11: holotype MO 22/79/0163, C 35849, py, 2: $\times 150$,
 11: $\times 250$; 7: paratype 534A-81-2-3/81/9099, C 35850, py, $\times 150$; 8: paratype 534A-81-2-3/
 81/9097, C 35851, $\times 150$.
- Fig. 3–4, 9 *Ristola altissima* (RÜST)
 (data 32, range 47, pob 164, rk 52), 3: 534A-106-1-29/81/9011, C 35852, py, $\times 100$; 4, 9:
 534A-126-2-125/81/9133, C 35853, py, note distally disappearing outer layer, 4: $\times 100$, 9:
 $\times 250$.
- Fig. 5, 10 *Ristola cretacea* (BAUMGARTNER)
 (data 101, range 93, pob 165, rk 23), MO 26/80/1857, C 35854, py, 5: $\times 100$, 6: $\times 250$.
- Fig. 6 *Ristola procera* (PESSAGNO)
 (data 45, range 72, pob 163, rk 97), POB 899/78/6275, C 35855, si, $\times 100$.
- Fig. 12 *Saitoum pagei* PESSAGNO
 (data 88, range 49, pob 20, rk 55), POB 986/78/8172, C 35/93, si, $\times 250$.
- Fig. 13 *Sethocapsa cetia* FOREMAN
 (data 68, range 87, pob 203, rk 39), POB 1205/79/5745, C 35856, si, $\times 75$.
- Fig. 14 *Sethocapsa trachyostraca* FOREMAN
 MO 46/79/4143, C 35857, py, $\times 150$.
- Fig. 15 *Sethocapsa uterculus* (PARONA)
 (data 111, range 109, pob 297, rk -), POB 1134/80/2671, C 35858, py, $\times 150$.
- Fig. 16 *Spongocapsula palmerae* PESSAGNO
 (data 50, range 38, pob 199, rk 76), 534A-125-5-72/81/9204, C 35859, py, $\times 100$.
- Fig. 17 *Spongocapsula perampla* (RÜST)
 (data 85, range -, pob 267, rk 9), POB 986/79/0202, C 35860, si, $\times 100$.
- Fig. 18 *Staurosphaera antiqua* RÜST
 (data 49, range 60, pob 218, rk 83), POB 899/78/6730, C 35861, si, $\times 100$.
- Fig. 19 *Stichocapsa convexa* YAO
 (data 61, range 16, pob 55, rk 56), 534A-125-3-29/81/2440, C 35862, py, $\times 150$.
- Fig. 20 *Stichocapsa* sp. aff. *S. japonica* YAO
 (data 4, range 3, pob 48, rk -), POB 1263/80/6730, C 35863, si, $\times 150$.




Plate 9

Scanning electron micrographs of Middle Jurassic to Early Cretaceous siliceous (si) and pyritized (py) Radiolaria from Blake-Bahama Basin (DSDP Site 534), Lombardy (POB 1205), Greece (POB 28, 899, 986), western Switzerland (POB 1134) and Japan (IN 7) (see locality descriptions).

- Fig. 1–2** *Stylocapsa oblongula* KOCHER
 (data 6, range 53, pob 59, rk 111), 1: 534A-125-3-29/81/2438, C 35864, py, $\times 250$; 2: POB 325/80/3802, C 35865, si, $\times 250$.
- Fig. 3–4** *Syringocapsa agolarium* FOREMAN
 (data 105, range 104, pob 291, rk –), 3: MO 22/79/3706, C 35866, py, $\times 150$; 3: 534A-81-2-64/81/9108, C 35867, py, $\times 150$.
- Fig. 5** *Syringocapsa lucifer* BAUMGARTNER n. sp.
 (data 96, range 91, pob 283, rk –), holotype POB 1205/79/5033, C 35858, si, $\times 75$.
- Fig. 6–7** *Tetradityma corralitosensis* (PESSAGNO)
 (data 20, range 17, pob 124, rk 58), 534A-126-2-125/81/9188, C 35869, py, 6: $\times 100$; 7: lateral view of same specimen as Figure 6, showing internal ray structure with three primary canals and cortical space, $\times 250$.
- Fig. 8–9, 13, 13a** *Tetradityma praeplena* BAUMGARTNER n. sp.
 (data 5, range 6, pob 125, rk –), 8: paratype IN 7/81/3027, C 35870, si, $\times 75$; 9, 13–13a: holotype IN 7/79/4404, C 35871, si, 9, 13: $\times 75$, 13a: note delicate porous cortical wall (arrow), $\times 250$.
- Fig. 10** *Tetratrabs zealis* (OZVOLDOVA)
 (data 36, range 24, pob 121, rk 61), POB 1341/81/2955, C 35872, si, small specimen! $\times 75$.
- Fig. 11** *Tetratrabs bulbosa* BAUMGARTNER
 (data 62, range 74, pob 122, rk 60), S 4/79/4700, C 35873, si, $\times 75$.
- Fig. 12, 14** *Tetradityma pseudoplena* BAUMGARTNER
 (data 57, range 36, pob 123, rk 59), 12: POB 28/78/3400, C 35874, si, $\times 75$; 14: holotype POB 899/79/1500, C 34760, si, $\times 75$.
- Fig. 15** *Thanarla pulchra* (SQUINABOL)
 (data 109, range 107, pob 296, rk –), MO 46a'/81/0948, C 35875, py, $\times 150$.
- Fig. 16–17** *Theocapsomma cordis* KOCHER
 (data 15, range 30, pob 277, rk 99), 534A-126-2-125/82/9094, C 35876, py, Figure 17 shows basal aperture, $\times 250$.






Plate 10

Scanning electron micrographs of Middle Jurassic to Early Cretaceous siliceous (si) and pyritized (py) Radiolaria from Blake-Bahama Basin (DSDP Site 534), Greece (POB 28, 899, 986, 1262), Romania (MO) and Sicily (S 4) (see locality descriptions).

- Fig. 1 *Triactoma cornuta* BAUMGARTNER
 (data 89, range 65, pob 166, rk 78), topotype POB 899/78/6085, C 35877, si, $\times 75$.
- Fig. 2 *Triactoma echiodes* FOREMAN
 (data 81, range 89, pob 94, rk 19), MO 46a'/81/0986, C 35878, si, $\times 100$.
- Fig. 3 *Triactoma blakei* (PESSAGNO)
 (data 25, range 46, pob 95, rk 64), 534A-126-2-125/81/9133, C 35879, py, $\times 75$.
- Fig. 4 *Triactoma jonesi* (PESSAGNO)
 (data 29, range 25, pob 96, rk 33), 534A-126-2-125/81/9131, C 35880, py, $\times 100$.
- Fig. 5 *Triactoma tithonianum* RÜST
 (data 30, range 52, pob 97, rk 40), POB 899/78/6173, C 35881, si, $\times 100$.
- Fig. 6–7 *Tricolocapsa plicarum* YAO
 (data 9, range 8, pob 51, rk –), 6: POB 1262/80/3954, C 35882, si, $\times 250$; 7: 534A-122-1-43/81/2242, C 35883, py, $\times 250$.
- Fig. 8 *Trillus* sp. cf. *T. seidersi* PESSAGNO & BLOME
 (data 1, range 1, pob 39, rk –), POB 1262/80/3957, C 35884, si, $\times 250$.
- Fig. 9 *Tritrabs casmiliaensis* (PESSAGNO)
 (data 26, range 45, pob 117, rk 81), POB 28/78/3777, C 35885, si, $\times 75$.
- Fig. 10 *Tritrabs ewingi* (PESSAGNO)
 (data 54, range 70, pob 113, rk 34), S 4/79/4689, C 35886, si, $\times 75$.
- Fig. 11 *Tritrabs exoticus* (PESSAGNO)
 (data 27, range 37, pob 118, rk 35), POB 899/78/6222, C 35887, si, $\times 75$.
- Fig. 12 *Tritrabs hayi* (PESSAGNO)
 (data 28, range 20, pob 116, rk 101), POB 899/78/6292, C 35888, si, $\times 75$.
- Fig. 13 *Tritrabs rhododactylus* BAUMGARTNER
 (data 27, range 26, pob 118, rk 35), POB 986/79/1631, C 35889, si, $\times 75$.
- Fig. 14–15 *Unuma echinatus* ICHIKAWA & YAO
 (data 2, range 4, pob 231, rk –), 14: POB 1262/80/2144, C 35890, si, $\times 150$; 15: POB 1262/80/2857, C 35891, si, $\times 150$.
- Fig. 16–17 *Xitus* sp. cf. *X. spicularius* ALIEV
 (data 106, range 98, pob 295, rk –), 16: MO 22/79/0177, C 35892, py, $\times 150$; 17: 534A-81-2-64/81/9104, C 35893, py, $\times 150$.







MIDDLE JURASSIC TO
EARLY CRETACEOUS
RADIOLARIAN RANGE CHART
AND ZONAL DEFINITIONS
BASED ON
UNITARY ASSOCIATIONS


(computed after GUEX & DAVID 1982, 1984)

CORRELATION TO
CHRONOSTRATIGRAPHY



CENTRAL ATLANTIC





NAMES OF FORMATIONS

A1	WILHELMSTADT CHART
B1	GRANADA CHART
B2	ALICE AND ALICE CHART
B3	GRANADA CHART
B4	GRANADA CHART
B5	GRANADA CHART
B6	GRANADA CHART
B7	GRANADA CHART
B8	GRANADA CHART
B9	GRANADA CHART
B10	GRANADA CHART
B11	GRANADA CHART
B12	GRANADA CHART
C1	GRANADA CHART
C2	GRANADA CHART
C3	GRANADA CHART
C4	GRANADA CHART
C5	GRANADA CHART
C6	GRANADA CHART
C7	GRANADA CHART
C8	GRANADA CHART
C9	GRANADA CHART
C10	GRANADA CHART
C11	GRANADA CHART
C12	GRANADA CHART
D1	GRANADA CHART
D2	GRANADA CHART
D3	GRANADA CHART
D4	GRANADA CHART
D5	GRANADA CHART
D6	GRANADA CHART
D7	GRANADA CHART
D8	GRANADA CHART
D9	GRANADA CHART
D10	GRANADA CHART
D11	GRANADA CHART
D12	GRANADA CHART
E1	GRANADA CHART
E2	GRANADA CHART
E3	GRANADA CHART
E4	GRANADA CHART
E5	GRANADA CHART
E6	GRANADA CHART
E7	GRANADA CHART
E8	GRANADA CHART
E9	GRANADA CHART
E10	GRANADA CHART
E11	GRANADA CHART
E12	GRANADA CHART
F1	GRANADA CHART
F2	GRANADA CHART
F3	GRANADA CHART
F4	GRANADA CHART
F5	GRANADA CHART
F6	GRANADA CHART
F7	GRANADA CHART
F8	GRANADA CHART
F9	GRANADA CHART
F10	GRANADA CHART
F11	GRANADA CHART
F12	GRANADA CHART
G1	GRANADA CHART
G2	GRANADA CHART
G3	GRANADA CHART
G4	GRANADA CHART
G5	GRANADA CHART
G6	GRANADA CHART
G7	GRANADA CHART
G8	GRANADA CHART
G9	GRANADA CHART
G10	GRANADA CHART
G11	GRANADA CHART
G12	GRANADA CHART
H1	GRANADA CHART
H2	GRANADA CHART
H3	GRANADA CHART
H4	GRANADA CHART
H5	GRANADA CHART
H6	GRANADA CHART
H7	GRANADA CHART
H8	GRANADA CHART
H9	GRANADA CHART
H10	GRANADA CHART
H11	GRANADA CHART
H12	GRANADA CHART
I1	GRANADA CHART
I2	GRANADA CHART
I3	GRANADA CHART
I4	GRANADA CHART
I5	GRANADA CHART
I6	GRANADA CHART
I7	GRANADA CHART
I8	GRANADA CHART
I9	GRANADA CHART
I10	GRANADA CHART
I11	GRANADA CHART
I12	GRANADA CHART
J1	GRANADA CHART
J2	GRANADA CHART
J3	GRANADA CHART
J4	GRANADA CHART
J5	GRANADA CHART
J6	GRANADA CHART
J7	GRANADA CHART
J8	GRANADA CHART
J9	GRANADA CHART
J10	GRANADA CHART
J11	GRANADA CHART
J12	GRANADA CHART
K1	GRANADA CHART
K2	GRANADA CHART
K3	GRANADA CHART
K4	GRANADA CHART
K5	GRANADA CHART
K6	GRANADA CHART
K7	GRANADA CHART
K8	GRANADA CHART
K9	GRANADA CHART
K10	GRANADA CHART
K11	GRANADA CHART
K12	GRANADA CHART
L1	GRANADA CHART
L2	GRANADA CHART
L3	GRANADA CHART
L4	GRANADA CHART
L5	GRANADA CHART
L6	GRANADA CHART
L7	GRANADA CHART
L8	GRANADA CHART
L9	GRANADA CHART
L10	GRANADA CHART
L11	GRANADA CHART
L12	GRANADA CHART
M1	GRANADA CHART
M2	GRANADA CHART
M3	GRANADA CHART
M4	GRANADA CHART
M5	GRANADA CHART
M6	GRANADA CHART
M7	GRANADA CHART
M8	GRANADA CHART
M9	GRANADA CHART
M10	GRANADA CHART
M11	GRANADA CHART
M12	GRANADA CHART
N1	GRANADA CHART
N2	GRANADA CHART
N3	GRANADA CHART
N4	GRANADA CHART
N5	GRANADA CHART
N6	GRANADA CHART
N7	GRANADA CHART
N8	GRANADA CHART
N9	GRANADA CHART
N10	GRANADA CHART
N11	GRANADA CHART
N12	GRANADA CHART
O1	GRANADA CHART
O2	GRANADA CHART
O3	GRANADA CHART
O4	GRANADA CHART
O5	GRANADA CHART
O6	GRANADA CHART
O7	GRANADA CHART
O8	GRANADA CHART
O9	GRANADA CHART
O10	GRANADA CHART
O11	GRANADA CHART
O12	GRANADA CHART
P1	GRANADA CHART
P2	GRANADA CHART
P3	GRANADA CHART
P4	GRANADA CHART
P5	GRANADA CHART
P6	GRANADA CHART
P7	GRANADA CHART
P8	GRANADA CHART
P9	GRANADA CHART
P10	GRANADA CHART
P11	GRANADA CHART
P12	GRANADA CHART
Q1	GRANADA CHART
Q2	GRANADA CHART
Q3	GRANADA CHART
Q4	GRANADA CHART
Q5	GRANADA CHART
Q6	GRANADA CHART
Q7	GRANADA CHART
Q8	GRANADA CHART
Q9	GRANADA CHART
Q10	GRANADA CHART
Q11	GRANADA CHART
Q12	GRANADA CHART
R1	GRANADA CHART
R2	GRANADA CHART
R3	GRANADA CHART
R4	GRANADA CHART
R5	GRANADA CHART
R6	GRANADA CHART
R7	GRANADA CHART
R8	GRANADA CHART
R9	GRANADA CHART
R10	GRANADA CHART
R11	GRANADA CHART
R12	GRANADA CHART
S1	GRANADA CHART
S2	GRANADA CHART
S3	GRANADA CHART
S4	GRANADA CHART
S5	GRANADA CHART
S6	GRANADA CHART
S7	GRANADA CHART
S8	GRANADA CHART
S9	GRANADA CHART
S10	GRANADA CHART
S11	GRANADA CHART
S12	GRANADA CHART
T1	GRANADA CHART
T2	GRANADA CHART
T3	GRANADA CHART
T4	GRANADA CHART
T5	GRANADA CHART
T6	GRANADA CHART
T7	GRANADA CHART
T8	GRANADA CHART
T9	GRANADA CHART
T10	GRANADA CHART
T11	GRANADA CHART
T12	GRANADA CHART
U1	GRANADA CHART
U2	GRANADA CHART
U3	GRANADA CHART
U4	GRANADA CHART
U5	GRANADA CHART
U6	GRANADA CHART
U7	GRANADA CHART
U8	GRANADA CHART
U9	GRANADA CHART
U10	GRANADA CHART
U11	GRANADA CHART
U12	GRANADA CHART
V1	GRANADA CHART
V2	GRANADA CHART
V3	GRANADA CHART
V4	GRANADA CHART
V5	GRANADA CHART
V6	GRANADA CHART
V7	GRANADA CHART
V8	GRANADA CHART
V9	GRANADA CHART
V10	GRANADA CHART
V11	GRANADA CHART
V12	GRANADA CHART
W1	GRANADA CHART
W2	GRANADA CHART
W3	GRANADA CHART
W4	GRANADA CHART
W5	GRANADA CHART
W6	GRANADA CHART
W7	GRANADA CHART
W8	GRANADA CHART
W9	GRANADA CHART
W10	GRANADA CHART
W11	GRANADA CHART
W12	GRANADA CHART
X1	GRANADA CHART
X2	GRANADA CHART
X3	GRANADA CHART
X4	GRANADA CHART
X5	GRANADA CHART
X6	GRANADA CHART
X7	GRANADA CHART
X8	GRANADA CHART
X9	GRANADA CHART
X10	GRANADA CHART
X11	GRANADA CHART
X12	GRANADA CHART
Y1	GRANADA CHART
Y2	GRANADA CHART
Y3	GRANADA CHART
Y4	GRANADA CHART
Y5	GRANADA CHART
Y6	GRANADA CHART
Y7	GRANADA CHART
Y8	GRANADA CHART
Y9	GRANADA CHART
Y10	GRANADA CHART
Y11	GRANADA CHART
Y12	GRANADA CHART
Z1	GRANADA CHART
Z2	GRANADA CHART
Z3	GRANADA CHART
Z4	GRANADA CHART
Z5	GRANADA CHART
Z6	GRANADA CHART
Z7	GRANADA CHART
Z8	GRANADA CHART
Z9	GRANADA CHART
Z10	GRANADA CHART
Z11	GRANADA CHART
Z12	GRANADA CHART