

Zeitschrift: Eclogae Geologicae Helvetiae
Herausgeber: Schweizerische Geologische Gesellschaft
Band: 78 (1985)
Heft: 2

Artikel: A correlation of the Tethyan Maiolica Formation of the Breggia section (southern Switzerland) with Early Cretaceous oozes of Site 534A, DSDP Leg 76 in the western Atlantic
Autor: Renz, Otto / Habicht, Konrad
Kapitel: F: Conclusions
DOI: <https://doi.org/10.5169/seals-165662>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

coincide with the 75 m (63 m) mark at the Maiolica section. Here, at the top of the Valanginian, a very conspicuous break in variability and abundance of aptychi can be observed, as displayed on the distribution chart (Fig. 2). This distribution of aptychi, we may assume, should also be reflected in ammonite assemblages from close to the Valanginian–Hauterivian boundary.

The intervals between 63 m and 92 m at the Maiolica section and between Core 67 to Core 68 in Hole 534A, and Core 24 to 25 in Hole 391C, both in the Blake-Bahama Basin, are dominated by aptychi distinguished by retroverse curving lamellae patterns. They belong to the group of *Lamellaptychus seranonis*, a form restricted according to present knowledge to the Hauterivian.

Barremian (102 to 130 m)

In the Blake-Bahama Basin (Hole 391A) and the Cape Verde Basin (Hole 367) considerable gaps without remains of Cephalopoda hamper the fixation of the Hauterivian–Barremian boundary.

Within the Maiolica section an interval (tunnel, Fig. 1) without aptychi of 9 m thickness separates the last *L. seranonis* (Hauterivian) from the first *L. angulocostatus*, a widely distributed form within the Tethys, indicating a Barremian age. The limit Hauterivian–Barremian thus remains unclear.

From 5 m below the top of the Maiolica Formation, near 125 m, an ammonite assemblage has been described by RIEBER (1977). The presence of *Pulchellia* and *Karsteniceras*, widely known from Europe (Spain) and northern South America (Colombia, Trinidad) represent a reliable indication for a Barremian age.

The discovery of *Pulchellia*, although flattened and poorly preserved, from Hole 534A in the Blake-Bahama Basin is of the greatest value for regional correlation. Our present correlation is based mainly on the occurrence of *Pulchellia* in Hole 534A, and in the Maiolica section.

F. Conclusions

The present study on aptychi assemblages from the Maiolica Formation in the river Breggia, based on detailed stratigraphic collecting of the fossils, permits a preliminary correlation with holes drilled in the northern Atlantic by the Deep Sea Drilling Project. Regional correlations, based on aptychi from sections exposed within the Tethyan realm of Europe have so far not been attempted, the main reason being the consecutive repetition of nearly identical lamellae patterns of aptychi during the Jurassic–Cretaceous Periods.

Specimens with very similar morphological features (homeomorph) occur within levels of quite different ages. In most publications on aptychi little attention has been paid to the stratigraphical order in spite of its significance for taxonomy. This is demonstrated by the synonymy lists included in the present paper.

The aptychi obtained from the Deep Sea Drilling Project provided an ideal opportunity for investigations of aptychi, this being a main reason for carrying out the present study.

At present we just begin to understand the biochronologic significance of aptychi assemblages, especially in sediments from the Tethys.

The results, evidently, need confirmation at other sections. We propose to study additional sections of deep water Tethyan sediments containing aptychi. Of special interest in this respect are the Jurassic and Cretaceous deposits in the Betic Cordillera in southern Spain, as shown by a few specimens from old collections figured here.

Acknowledgments

The paper has been prepared in the Museum of Natural History, Basel. The authors thank Edith Müller-Merz for some assistance in preparing the manuscript and Urs Marrer for helping to draft the text-figures. The photographs on Plates 1–5 were made by Wolfgang Suter of the Museum's photographic laboratory.

REFERENCES

- BACHMAYER, F. (1963): Beiträge zur Paläontologie oberjurassischer Riffe. I. Die Aptychen (Ammonoidea) des Oberjura von Stramberg (ČSSR). II. Die Aptychen der Klentnitzer Serie in Österreich. – *Ann. nathist. Mus. Wien* 66, 125–138.
- BERGER, W. H. (1970): Planktonic Foraminifera: selective solution and the lysocline. – *Marine Geol.* 8, 111–138.
- (1973): Deep-Sea carbonates: evidence for a coccolith lysocline. – *Deep Sea Res.*, 20, 917–921.
- (1976): Biogenous deep sea sediments. Production, preservation and interpretation. – In: RILEY, J. P., & CHESTER, R. (Ed.): *Chem. Oceanogr.* 5, 266–388.
- BERNOULLI, D. (1964): Zur Geologie des Monte Generoso (Lombardische Alpen). – *Beitr. geol. Karte Schweiz [N.F.]* 118.
- (1966): Pontegana-Konglomerat. – *Lex. stratigr. int.*, I, Europe, Fasc. 7, Suisse, Fasc. 7c, Alpes suisses et Tessin méridional, II, p. 881.
- (1972): North Atlantic and Mediterranean Mesozoic facies. – *Init. Rep. DSDP 1*, 801–871.
- (1980a): Southern Alps of Ticino. In: *Geology of Switzerland, a Guide Book, part A*, TRÜMPY, R.: An outline of the geology of Switzerland (p. 80–82). – Wepf & Co., Basel.
- (1980b): Mendrisio–Breggia valley–Chiasso. In: *Geology of Switzerland, a Guide Book, Part B, Geological Excursions*, p. 207–209. – Wepf & Co., Basel.
- BERNOULLI, D., BICHSEL, M., BOLLI, H. M., HÄRING, M. O., HOCHULI, P. A., & KLEBOTH, P. (1981): The Missaglia Megabed, a catastrophic deposit in the Upper Cretaceous Bergamo Flysch, northern Italy. – *Eclogae geol. Helv.* 74/2, 421–442.
- BERNOULLI, D., & JENKYN, H. C. (1974): Alpine, Mediterranean and Central Atlantic Mesozoic facies in relation to the early evolution of the Tethys. – *Spec. Publ. Soc. econ. Paleont. Mineral.* 19, 129–160.
- BICHSEL, M., & HÄRING, M. O. (1981): Facies evolution of Late Cretaceous Flysch in Lombardy (northern Italy). – *Eclogae geol. Helv.* 74/2, 383–420.
- BITTERLI, P. (1965): Bituminous intercalations in the Cretaceous of the Breggia river, S. Switzerland. – *Bull. Ver. schweiz. Petroleum-Geol. u. -Ing.* 31/81, 179–185.
- BIJU-DUVAL, B., DERCOURT, J., & LE PICHON, X. (1977): From the Tethys ocean to the Mediterranean seas: A plate-tectonic model of the evolution of the western Alpine system. In: BIJU-DUVAL, B., & MONTADERT, L. (Ed.): *International Symposium on the structural history of the Mediterranean basins, Split (Yugoslavia)* (p. 143–164). – Technip, Paris.
- BLASCHKE, F. (1911): Zur Tithonfauna von Stramberg in Mähren. – *Ann. nathist. Mus. Wien* 25, 143–222, Pl. 1–6.
- COQUAND, M. (1841): *Mémoire sur les Aptychus*. – *Bull. Soc. géol. France* 12, 376–391.
- DURAND-DELGA, M., & GASIOROWSKI, S. M. (1970): Les niveaux à *Aptychus* dans les pays autour de la Méditerranée occidentale et dans les Carpathes. – *C. R. Acad. Sci. (Paris) (D)* 270, 767–770.
- ERRICO, G., GROPPI, G., SAVELLI, S., & VAGHI, G. C. (1980): Malossa Field: A deep discovery in the Po valley, Italy. In: HALBOUTY, M. T. (Ed.): *Giant Oil Fields of the Decade: 1968–1978* (p. 525–538). – *Mem. amer. Assoc. Petroleum Geol.* 30.
- FAVRE, E. (1875): Description des fossiles du terrain jurassique de la Montagne des Voirons (Savoie). – *Mém. Soc. paléont. suisse* 2, 1–77.
- FAZZINI, P. (1965): Sulla presenza del *Lamellaptychus angulocostatus* Peters al tetto dei diaspiri nei Monti di Poggiano presso Montepulciano (Siena) (p. 1–10). – *Inst. geol. Modena*.