

Zeitschrift: Eclogae Geologicae Helvetiae
Herausgeber: Schweizerische Geologische Gesellschaft
Band: 79 (1986)
Heft: 2

Artikel: Mineralostratigraphy, litho- and biostratigraphy combined in correlation of the Oxfordian (Late Jurassic) formations of the Swiss Jura range

Autor: Gygi, Reinhart A. / Persoz, Francis

Inhaltsverzeichnis

DOI: <https://doi.org/10.5169/seals-165840>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 12.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

CONTENTS

1. Introduction	387
1.1 Previous work	387
1.2 Purpose of this paper	390
2. Methods	391
3. Results	394
3.1 Formations, members, and beds	394
3.2 Main types of facies	409
3.3 Lateral facies transitions	417
3.4 The ammonite succession	420
3.5 The nature and distribution of siliciclastic minerals	424
4. Interpretation and discussion	437
4.1 Provenance and variation in siliciclastic minerals	437
4.2 Correlation	441
4.3 Depositional sequences	443
5. Conclusions	447

1. Introduction

1.1 Previous work

Stratigraphic work in the Swiss Jura range began before 1820. MERIAN (1821) correlated the Early and Middle Oxfordian marl-clays (Renggeri Member and Terrain à Chailles Member) of northwestern Switzerland with the Effingen Member in canton Aargau, and the coral limestones of the St-Ursanne Formation in the “basinal” realm of canton Aargau (see Pl. 1A). GRESSLY (1838–41) carried out extensive geological mapping in canton Solothurn and in adjacent areas. Close observation of the coral bioherms and the coeval fine-grained sediments of the St-Ursanne Formation at La Caquerelle near St-Ursanne inspired this author to introduce the concept of facies into the scientific literature. Ironically, the great effort in mapping, practical use of the new stratigraphic method, and ample fossil collecting did not allow this distinguished geologist to arrive, in his own opinion, at a satisfactory time correlation between the deposits from shallow water and those of deeper marine origin. The progress of paleontology as pioneered by OPPEL (1856–8, 1862–3) in the Oxfordian of this region led to an important revision of Merian’s correlation. On the evidence of ammonites, OPPEL (1857, p. 626) recognized that the thick Renggeri Member and part of the Terrain à Chailles Member in the northwest thin out to the southeast and grade into what is now called the Schellenbrücke Bed (Table 2 and Pl. 1A). This is a ferruginous marly limestone with iron ooids. The thickness of the bed is normally less than 10 cm. ROLLIER (1888, p. 87) correlated the Liesberg Member with the Birmenstorf Member, and the St-Ursanne Formation with the Effingen and Geissberg Members. He had no ammonites from the platform deposits to support his assertion. Later, ROLLIER (1911, Fig. 54) reaffirmed his view, and it remained unchallenged until 1967, when BOLLIGER & BURRI proposed another correlation which was based on the distribution of detrital quartz. The significance of the important paleontological work by de Loriol in relation to Rollier’s correlation was not appreciated by most stratigraphic workers, perhaps because de Loriol paid so little attention to stratigraphy. In a short review, ARKELL (1956, p. 95–96) threw some light on the stratigraphic