

Zeitschrift: Eclogae Geologicae Helvetiae
Herausgeber: Schweizerische Geologische Gesellschaft
Band: 79 (1986)
Heft: 3

Artikel: The foraminifera in the lower Cretaceous of Trinidad, W.I. Part 5, Maridale formation, upper Part : Hedbergella rohri zone
Autor: Bartenstein, Helmut / Bolli, Hans M.
Kapitel: Index to genera and species. Part 1-5
DOI: <https://doi.org/10.5169/seals-165857>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

below the *Hedbergella rohri* Zone, or *Spiroplectinata annectans* and the genera *Pleurostomella* and *Pseudosigmoilina* of which the earliest occurrences are from the Middle Albian.

Based on the above evidence, the previously held view (BARTENSTEIN & BOLLI 1977, p. 561) that the *Hedbergella rohri* Zone occurred in the range Lower to Upper Albian has to be revised.

Index to genera and species, Part 1–5

(Part 1: 1957, Part 2: 1966, Part 3: 1973, Part 4: 1977, Part 5: 1986)

Legend

- 1 = Barremian Toco Formation, *Lenticulina barri* Zone
 Cuche Formation, *Lenticulina ouachensis ouachensis* Zone
 2 = Late Aptian Maridale Formation, Type locality, *Planomalina maridalensis* Zone
 3 = Late Aptian Maridale Formation, Co-Type locality, *Planomalina maridalensis* Zone
 4 = Early Aptian Cuche Formation, *Leupoldina protuberans* Zone
 5 = Late Aptian to earliest Albian Maridale Formation, *Hedbergella rohri* Zone.
 1:145 = number of paper and page. – New taxa are indicated with name and first citation in bold face.

- Agathammina*
 sp.? 5: 954
- Ammobaculites*
euides 3: 394, 5: 949
goodlandensis 2: 139, 3: 393
 cf. *goodlandensis* 3: 394
reophacoides 4: 546, 562–563, 5: 949
 sp. 1: 17
subcretaceus 1: 17, 2: 139, 3: 394, 4: 546, 5: 950
torosus 2: 139, 3: 394
***trinidadensis* 1: 17**
- Ammodiscus*
gaultinus 2: 140, 3: 394; see *Glomospirella gaultina*
tenuissimus 5: 947
- Arenobulimina* 5: 978
- Bigenerina*
clavellata 3: 395, 5: 950
 cf. *clavellata* 2: 141
- Biglobigerinella*
barri 2: 164, 3: 411
- Bolivina*
textilarioides 1: 42
- Bullopora*
laevis 5: 971
- Choffatella*
decipiens 4: 549
- Citharina* 5: 976
acuminata 1: 39, 4: 554, 562–563
- Clavihedbergella*
subcretacea 4: 559
- Conorotalites*
aptiensis 2: 162, 3: 411, 4: 558, 5: 974
bartensteini intercedens 1: 48, 4: 562, 564
- Cornuspira*
cretacea 3: 409
- Dentalina* 5: 955
aequivoca 5: 955
***bonaccordensis* 5: 955**
catenula 5: 956
communis 1: 34, 2: 153, 3: 404, 5: 956
cylindroides 2: 153, 3: 405, 5: 957
debilis 1: 35, 5: 957
 cf. *deflexa* 2: 153
distincta 2: 153, 3: 405, 5: 957
expansa 5: 957
filiformis 5: 957
gracilis 1: 34, 2: 153, 3: 404, 5: 958
guttifera 2: 154; see *subguttifera* 3: 405, 5: 959
linearis 1: 35, 2: 153, 3: 405, 5: 958; see *Nodosaria linearis*
nana 1: 35, 2: 152, 3: 404, 5: 958
 aff. *oligostegia* 5: 958
 cf. *porcatulata* 2: 154; see ***bonaccordensis* 5: 955**
soluta 2: 152, 3: 404, 5: 959
subguttifera 1: 34, 3: 405, 5: 959
 cf. *terquemi* 5: 959
- Dorothia*
 cf. *conula* 2: 144; see *gradata*
filiformis 2: 144, 3: 397, 4: 562, 564, 5: 950
gradata 3: 397, 4: 549, 5: 950
- Epistomina* 5: 977
caracolla caracolla 1: 46, 4: 557
***hechti* 1: 46**
ornata 1: 46, 4: 557
spinulifera spinulifera 4: 557
- Falsoguttulina*; see *Guttulina*
vandenboldi 2: 158, 3: 407, 5: 972
- Falsopalmula*
 sp. 2: 157

*Flabellinella**didyma* 5: 959*Fronicularia**gaultina* 2: 156, 3: 406, 5: 960

sp. 1 1: 39

sp. 2 1: 39

*Gaudryina**compacta* 5: 950*dividens* 2: 141, 3: 396, 4: 562, 564, 5: 951*klamathensis* 5: 951*reicheli* 2: 142, 3: 396, 5: 951*Gaudryinella**hannoverana* 4: 547, 562–563*sherlocki* 2: 141, 3: 395, 4: 547, 562–563, 5: 951*Gavelinella*cf. *ammonoides* 3: 410*barremiana* 1: 47, 4: 558, 562, 564*intermedia* 2: 161, 3: 410, 4: 558, 562, 564, 5: 975

Genus et Species incertae sedis 2: 159, 3: 398

Globigerina see *Hedbergella**Globigerinelloides*cf. *blowi* 5: 979*ferreolensis* 5: 979*gyroidinaeformis* 5: 979*Globorotalites* see *Conorotalites**Globulina**prisca* 2: 158, 3: 407, 4: 555, 5: 972cf. *prisca* 1: 41*Glomospira**charoides* 5: 947*gordialis* 2: 140, 3: 395, 5: 947*Glomospirella**gaultina* 5: 947; see *Ammodiscus gaultinus**Guttulina*aff. *symploca* 5: 972***vandenboldi* 1: 40**; see *Falsoguttulina vandenboldi**Haplophragmium*cf. *aequale* 1: 18*Haplophragmoides**concaus* 1: 16, 2: 138, 3: 393, 4: 545, 5: 952*nonioninoides* 5: 952

(?) sp. 4: 545

*Hedbergella**delrioensis* 2: 164, 3: 412, 5: 979*infracretacea* 4: 559*kugleri* 4: 559*planispira* 4: 559*rohri* 5: 980*Hippocrepina**depressa* 5: 947*Hormosina**ovulum* 5: 948*Hyperammia**gaultina* 2: 137, 3: 392, 5: 948*Kalamopsis**grzybowski* 5: 948*Lagena**apiculata* 5: 960*globosa* 5: 960*laevis* 1: 40, 2: 157, 3: 407, 5: 960aff. *oxystoma* 5: 960*Lamarckina**lamplughii* 3: 411, 4: 562, 564*Lenticulina* 5: 961*acuta* 2: 146, 3: 400, 5: 961cf. *acuta* 2: 147, 3: 400***antillica* 5: 961*****barri* 1: 28**cf. *bronni* 1: 33*calliopsis* 2: 149, 3: 402, 5: 962***caribica* 5: 962***cephalotes* 2: 150, 3: 403, 5: 962*complanata* 5: 962*crepidularis* 1: 29, 4: 551, 562–563, 5: 976*cultrata* 1: 23*eichenbergi* 1: 27, 4: 562, 564*excentrica* 2: 149, 3: 402*frankei* 1: 33, 4: 552, 562–563*gaultina* 5: 963*gracilissima* 4: 552cf. *gracilissima* 1: 31*grata* 2: 148, 3: 401, 5: 963cf. *grata* 1: 30*guttata* 1: 27, 2: 146 (see *meridiana*), 4: 562, 564*harpa* 2: 150, 3: 402*inaequalis* 5: 963*incurvata* 1: 30, 2: 150, 3: 402, 5: 963***kugleri* 1: 27**, 4: 551, 562–563cf. *kugleri* 1: 28*lituola* 2: 150, 3: 403, 5: 964***maridalensis* 3: 401**, 4: 552, 5: 977*matutina* 1: 31*meridiana* 2: 146, 3: 399, 4: 562, 564, 5: 964*muensteri* 1: 22, 4: 550, 5: 964*nodosa* 1: 24, 4: 550, 562–563, 5: 977*ouachensis ouachensis* 1: 25, 4: 550, 562–563, 5: 977cf. *ouachensis* 1: 27***ouachensis multicella* 1: 26***ouachensis wisselmanni* 1: 26*perobliqua* 5: 964*planiuscula* 2: 148, 3: 402***praegaultina* 1: 24***prima* 1: 31; see *excentrica**pulchella* 2: 147, 3: 400*robusta* 5: 965*roemeri* 1: 23, 5: 965*saxocretacea* 2: 146, 3: 399, 5: 965*schloenbachi* 5: 965cf. *schreiteri* 2: 145*scitula* 2: 149, 3: 402, 5: 965*secans* 5: 966*sigali* 1: 32

- (*Lenticulina*, cont.)
 sp.1 1: 23, 32, 33
 sp.2 1: 29
 sp. (n.sp.) 2: 148 (see *maridalensis*)
spinosa 2: 151, 3: 403, 4: 553, 5: 977
strombecki 2: 147, 3: 400, 5: 966, 977
subalata 1: 23
subaperta 3: 400
subgaultina 2: 147, 3: 401
tricarinnella 1: 30, 4: 562, 5: 976
turgidula 2: 147, 3: 400, 5: 966
vocontiana 3: 398, 5: 966
- Leupoldina*
protuberans 4: 559
- Lingulina*
loryi 2: 155, 3: 406, 5: 966
praelonga 1: 38
- Marginulina*
bullata 5: 967
linearis 2: 151, 3: 403; see *Nodosaria linearis*
 5: 967
pyramidalis 1: 34, 2: 151, 3: 403, 5: 967
- Marssonella*
kummi 4: 548
oxycona 2: 144 (see *praeoxycona*), 3: 396, 4: 562,
 564, 5: 952
 cf. *oxycona* 1: 20; see *kummi*
praeoxycona 3: 397, 4: 548, 562–563, 5: 976
subtrochus 2: 144, 3: 397, 4: 548, 5: 952
 cf. *trochus* 1: 20; see *subtrochus*
- Nodosarella*
rohri 1: 43
- Nodosaria* 5: 955
jonesi 5: 967
 cf. *chapmani* 1: 36
linearis 5: see *Dentalina linearis*
nuda 5: 968
obscura 1: 36, 5: 968
orthopleura 2: 152, 5: 968
paupercula 1: 36, 5: 968
sceptrum 1: 35, 3: 404, 4: 553, 5: 969
 sp. 2: 152
 cf. *zippei* 1: 36
- Orthokarstenia*
shastaensis 4: 554, 5: 969
- Patellina*
subcretacea 1: 45, 4: 556, 5: 976
- Planomalina*
blowi 4: 560; see *Globigerinelloides* cf. *blowi*
chenioureensis 5: 980
maridalensis 2: 163, 3: 411
saundersi 2: 163, 3: 411
 sp. 2: 163
- Plectorecurvoides*
alternans 5: 952
- Pleurostomella* 5: 978
- Praeglobotruncana* see *Hedbergella*
- Proteonina*
ampullacea 1: 15
- Psammosphaera*
fusca 1: 14; see *scruposa*
scruposa 3: 392
 sp. 5: 948
- Pseudoglandulina*; see *Pseudonodosaria*, *Rectoglandulina*
humilis 1: 37
mutabilis 1: 37
- Pseudonodosaria*; see *Pseudoglandulina*, *Rectoglandulina*
humilis 5: 969
mutabilis 5: 970
- Pseudosigmoilina* 5: 978
- Pyrulina*
cylindroides 2: 158, 3: 407, 5: 973
exserta 2: 158, 3: 407, 5: 973
 cf. *exserta* 1: 41
- Quadratina*
maertensi 2: 157
- Quinqueloculina* 5: 954
sabella 5: 955
 sp.? 5: 955
- Ramulina* 2: 159
aculeata 2: 159 (pars: *berthelini*), 3: 408, 4: 555,
 5: 973
berthelini 3: 408, 5: 973
fusiformis 1: 41
globulifera 2: 159, 5: 973
grandis 3: 407, 5: 974
laevis 2: 159, 5: 974
spandeli 1: 42
 sp. 3: 407
- Rectoglandulina*; see *Pseudoglandulina*, *Pseudonodosaria*
humilis 2: 155, 3: 406
mutabilis 2: 154, 3: 405
 cf. *mutabilis* 2: 154
- Reophax*
guttifer 1: 16, 2: 137, 3: 393, 5: 949
guttifer Form a 1: 16
minutus 5: 949
scorpiurus 1: 15
pilulifer 1: 15, 2: 137, 3: 392, 5: 949
 sp. 4: 545
subfusiformis 1: 16
- Rhizammina*
indivisa 1: 14; see *Hyperammina gaultina*
 sp. 1: 14
- Schackoina*
pustulans pustulans 2: 163, 3: 411, 4: 560
pustulans quinquecamerata 4: 560
reicheli 5: 981

*Spirillina**minima* 1: 44, 2: 161, 3: 409, 4: 556, 5: 976*Spiroplectinata**annectens* 5: 978*Textularia**bettenstaedti* 5: 953

sp. 2: 140

*Triplasia**emslandensis acuta* 1: 18 (new name: *georgsdor - fensis*)*emslandensis emslandensis* 1: 18 (new name: *georgsdorfensis*)*Tristix**acutangula* 1: 37, 2: 157, 3: 406, 5: 970*globulifera* 5: 970*Tritaxia**plummerae* 5: 953*pyramidata* 1: 19, 4: 546, 5: 977*tricarinata* 5: 977*Trochammina**depressa* 4: 549, 5: 953

sp. (n. sp.) 1: 21

(?)sp. 4: 545

sp. 5: 954

*Trocholina**infragranulata* 1: 44*Vaginulina**arguta* 1: 38, 4: 553, 5: 970*geinitzi* 2: 156, 3: 406*geisendoerferi* 5: 971*kochi* 1: 38*procera* 1: 39, 4: 562, 564*recta* 1: 38, 2: 155, 3: 406, 4: 553, 5: 971cf. *recta tenuistriata* 2: 156

sp. 2: 155, 3: 406

striolata 5: 971*Valvulina**fusca* 2: 145, 3: 398, 5: 954*Valvulineria**gracillima* 2: 161*loetterlei* 3: 410, 4: 556, 5: 975*Verneuilina*

sp. 3: 395

*Verneulinoides**neocomiensis* 1: 19*subfiliformis* 1: 19, 4: 546, 562–563, 5: 954

Total number: 78 genera, 185 species and subspecies, 13 new species or new subspecies, and 52 forms with open nomenclature.

Other microfossils

Holothuri 4: 560

Ostracoda 2: 165

Radiolaria 4: 561

Acknowledgments

The authors wish to express their sincere thanks to F. Allemann and to F. Zweili (both University of Bern) for the preparation of the SEM photographs of the fauna reproduced in this paper. They are grateful to J. P. Beckmann (ETH Zürich) and to J. B. Saunders (Museum of Natural History, Basel) for discussing and reviewing the manuscript.

REFERENCES

- BARTENSTEIN, H. (1976): Foraminiferal zonation of the Lower Cretaceous in North West Germany and Trinidad, West Indies. – *N. Jb. Geol. Paläont. [Mh.]* 3, 187–192.
- (1977): *Falsogaudryinella* n. g. (Foraminifera) in the Lower Cretaceous. – *N. Jb. Geol. Paläont. [Mh.]* 7, 385–401.
- (1979): Worldwide zonation of the Lower Cretaceous using benthonic Foraminifera. – *Newsl. Stratigr.* 7/3, 142–154.
- (1985): Stratigraphic pattern of index foraminifera in the Lower Cretaceous of Trinidad. – *Newsl. Stratigr.* 14/2, 110–117.
- BARTENSTEIN, H., BETTENSTAEDT, F., & BOLLI, H. M. (1957): Die Foraminiferen der Unterkreide von Trinidad, B.W.I. Erster Teil: Cuche- und Toco-Formation. – *Eclogae geol. Helv.* 50/1, 5–67.
- (1966): Die Foraminiferen der Unterkreide von Trinidad, W.I. Zweiter Teil: Maridale-Formation (Typlokalität). – *Eclogae geol. Helv.* 59/1, 129–177.
- BARTENSTEIN, H., & BOLLI, H. M. (1973): Die Foraminiferen der Unterkreide von Trinidad, W.I. Dritter Teil: Maridaleformation (Co-Typlokalität). – *Eclogae geol. Helv.* 66/2, 389–418.
- (1977): The Foraminifera in the Lower Cretaceous of Trinidad, W.I. Part 4: Cuche Formation, upper Part; *Leupoldina protuberans* Zone. – *Eclogae geol. Helv.* 70/2, 543–573.