

Zeitschrift: Eclogae Geologicae Helvetiae
Herausgeber: Schweizerische Geologische Gesellschaft
Band: 87 (1994)
Heft: 1

Artikel: Depositional trends in the Valdorbia Section (central Italy) during the Early Jurassic, as revealed by micropaleontology, sedimentology and geochemistry

Autor: Monaco, P. / Nocchi, M. / Ortega-Huertas, M.

Inhaltsverzeichnis

DOI: <https://doi.org/10.5169/seals-167447>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 25.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Depositional trends in the Valdorbia Section (Central Italy) during the Early Jurassic, as revealed by micropaleontology, sedimentology and geochemistry

P. MONACO¹, M. NOCCHI¹, M. ORTEGA-HUERTAS², I. PALOMO², F. MARTINEZ²
& G. CHIAVINI¹

Key words: Depositional trends, micropaleontology, sedimentology, geochemistry, Early Jurassic, Umbria-Marche basin, Central Italy

Abstract, Riassunto

| | |
|--|-----|
| 1 Introduction | 159 |
| 2 Geological setting | 161 |
| 3 Stratigraphy | 161 |
| 4 Ages | 167 |
| 5 Micropaleontology | 167 |
| 5.1 Data presentation | 167 |
| 5.1.1 Microfacies | 167 |
| 5.1.2 Microfossils | 170 |
| 5.2 Data discussion | 179 |
| 5.2.1 Paleoecological inferences | 179 |
| 6 Sedimentology | 185 |
| 6.1 Calcareous turbidites and associated gravity-flow deposits | 185 |
| 6.1.1 Description | 185 |
| 6.1.2 Age | 185 |
| 6.1.3 Interpretation | 188 |
| 6.2 Hummocky cross-stratified (HCS) deposits | 188 |
| 6.2.1 Description | 188 |
| 6.2.2 Age | 189 |
| 6.2.3 Interpretation | 189 |
| 6.3 Winnowed beds (WB) | 191 |
| 6.3.1 Description | 191 |
| 6.3.2 Age | 191 |
| 6.3.3 Interpretation | 191 |
| 6.4 A vertical trend from turbidites to HCS and WB deposits | 192 |

¹ Dipartimento di Scienze della Terra, Piazza dell'Università, Università degli Studi di Perugia, I-06100 Perugia

² Dpto. Mineralogía y Petrología, Fuentenueva, s/n. Universidad de Granada, E-18002 Granada

| | |
|--|-----|
| 7 Trace fossil assemblages | 192 |
| 7.1 Burrowing during authigenic sedimentation | 192 |
| 7.2 The significance of burrowing during turbiditic deposition | 193 |
| 7.3 The significance of burrowing during HCS deposition | 196 |
| 8 Mineralogy and geochemistry | 197 |
| 8.1 Methods | 197 |
| 8.2 Results and anoxic interval of deposition | 197 |
| 9 Discussion and conclusions | 199 |
| 9.1 The Lower Toarcian anoxic event | 199 |
| 9.2 Reworking | 203 |
| 9.3 Depositional trends | 204 |
| 9.4 Tectonics and eustacy | 206 |
| Acknowledgements | 206 |
| References | 207 |

ABSTRACT

In the Umbria-Marche basin open marine Jurassic sediments are well exposed in the Valdorbia section (ENE of Gubbio, Central Italy). The time interval considered here is from the Carixian to the Early Aalenian. The depositional units, already dated by means of ammonites and calcareous nannofossils, are: limestones and cherty limestones, Pliensbachian in age (“Corniola” = COR); marls of Early – Middle Toarcian age (“Marne del Monte Serrone” = MS), including black shales in the Tenuicostatum Zone; reddish nodular calcareous marls and limestones, Middle Toarcian to Early Aalenian in age, which constitute the “Rosso Ammonitico Umbro-Marchigiano” (= RAUM); and bivalve-bearing cherty limestones, Aalenian in age (“Calcari a Posidonia” = CP). Micropaleontological, sedimentological, trace fossil and geochemical-mineralogical analyses have been carried out. The microfossil study has revealed changes in the microforaminiferal assemblages, corresponding to changes in both oxygen conditions and depth of the sea floor: Miliolina, Textulariina and Lagenina are common in the Carixian; Textulariina and Lagenina in the Domerian and Lagenina, Spirillinina in the Toarcian/Aalenian. Opportunistic small species bloom in the most anoxic levels of the black shales.

The sedimentological study reveals two peaks in the detrital sedimentation. The first – probably connected with local tectonics (without excluding sea-level changes) – is found in the interval from the Carixian to the lower part of the Lower Toarcian. Metre-scale cycles of fine-grained calcareous turbidites, due to low-density flows, evolve gradually into coarse-grained, metre-thick turbidites often amalgamated and containing reworked skeletal grains of a carbonate platform environment, and gravity flow deposits. The second peak occurs in the Middle-Upper Toarcian. Fine-grained turbidites are overlain by hummocky cross-stratified (HCS) deposits and winnowed beds (WB), with large and pervasive bioturbation. The vertical transition from turbidites to sharp-based HCS deposits and WB is probably indicative of a general regressive trend and of a depositional environment above major storm wave base. This trend has also been indicated from the microfossil study.

Geochemical analysis of the Lower Toarcian (Tenuicostatum Zone) has revealed strong positive anomalies in Ba, V, Cr, Ni, Co, Cu, Zn, As, Sb and Pb, elements which are characteristic of black shale episodes. Weaker positive anomalies occur in similar sediments of the lower part of MS Formation, while such positive anomalies are absent in the largely bioturbated sediments deposited below (COR) and above (RAUM) the MS.

Depositional trends related to tectonic-eustatic variations in the depositional environment are suggested on the basis of information provided by the study of the Valdorbia Section and of other Umbria-Marche sections.

RIASSUNTO

Sedimenti pelagici del Giurassico sono ben esposti lungo la sezione della Valdorbia (ENE di Gubbio, Appennino Centrale). Tale sezione è ben nota in quanto è indicativa della sedimentazione giurassica di mare aperto che ha sostituito, nel Lias inferiore dell'area Umbro-Marchigiana, la sedimentazione tipica di una piattaforma