Zeitschrift:	Eclogae Geologicae Helvetiae
Herausgeber:	Schweizerische Geologische Gesellschaft
Band:	87 (1994)
Heft:	1
Artikel:	Depositional trends in the Valdorbia Section (central Italy) during the Early Jurassic, as revealed by micropaleontology, sedimentology and geochemistry
Autor:	Monaco, P. / Nocchi, M. / Ortega-Huertas, M.
Kapitel:	4: Ages
DOI:	https://doi.org/10.5169/seals-167447

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

4. Ages

The biostratigraphy is based on ammonite horizons within the Tenuicostatum and Serpentinus Zones (Venturi, in press) and on provisional horizons within the other zones studied by Venturi and not yet published (personal communication). The ammonite horizons are summarized in Table 1 and ammonite zones have been indicated, for graphic convenience, with age initials followed by a number. Each horizon is reported in metres in Figure 4, 5 and 6, following the Channell et al. (1984) sampling. Recently published data of the Valdorbia Section have been used as well (Cresta et al. 1989, Fig. 37 and 38). Calcareous nannofossil events, reported by Reale (1989), are located on the right side of the four partial sections. The lower part of the Valdorbia section (530 m to 510 m) has been attributed to the Carixian on the base of the first occurrence of Mitrolithus jansae at 499.8 m (Fig. 4) which falls in the upper part of the Davoei Zone (Carixian) according to Reale et al. (1991).

5. Micropaleontology

An important micropaleontological study of the Valdorbia Jurassic microfacies has been carried out by Centamore et al. (1971) in which the Valdorbia section was indicated as "Sezione del Sentino Ovest". A preliminary study of the microfossils occurring in the Valdorbia section both in thin section and in the sample washed residues was illustrated by Cresta et al. (1988), by means of a schematic distribution chart. No distinction between microfacies (observed in thin section) and microfossils (separated from the sediments) was indicated and a further study has allowed a better understanding of the microfaunal distribution and of the microfossil reworking.

In this work the micropaleontological observations concern both the microfaunal content of the microfacies analysed in thin section and the assemblages of the separated microfossils from the washed residues. 160 samples have been collected in stratigraphic order at 1 m interval roughly spanning the period of time from the Carixian to the Early Aalenian. 92 samples, mainly limestones have been processed for microfacies study from the COR, while 68 samples, mainly argillaceous and marly sediments, about 400 gr. each, have been collected mainly from the MS and RAUM units.

5.1 Data presentation

5.1.1 Microfacies

The fossil content is summarised in Figure 20 where the microfacies are arranged in groups with different characteristics.

Microfacies 1 (Carixian). The most characteristic fossils are stout calcareous sponge spicules and recrystallized echinoid remains (Pl. 1, Fig. 1, 2 and 3) while other organisms are scattered discontinuously. Radiolarians are associated with gastropods and other microfossils. This microfacies contains Miliolina (Pl. 1, Fig. 8), simple agglutinated foraminifers (Pl. 1, Fig. 5), plurilocular agglutinated foraminifers and Lagenina, both sculptured and smooth (Pl. 1 Fig. 7, Fig. 1, 2, 12, 13 and 14) of which Lenticulina is the dominant taxon. Besides Ophthalmidium (Pl. 1, Fig. 11) and Agerina martana (Pl. 1, Fig. 10), there are some tubular, porcellaneous, biloculine forms interpreted here as Planiinvoluta (Pl. 1, Fig. 6 and 9) with a porcellaneous test, according to Koehn Zaninetti (1969). The latter are characteristic but of controversial interpretation be-