

**Zeitschrift:** Eclogae Geologicae Helvetiae

**Herausgeber:** Schweizerische Geologische Gesellschaft

**Band:** 87 (1994)

**Heft:** 3: Concepts and controversies in phosphogenesis : proceedings of the symposium and workshop held on 6-10 September 1993

**Artikel:** Phosphorite geochemistry : state-of-the-art and environmental concerns

**Autor:** Jarvis, Ian / Burnett, William C. / Nathan, Yaacov

**DOI:** <https://doi.org/10.5169/seals-167474>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 12.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Phosphorite geochemistry: State-of-the-art and environmental concerns

IAN JARVIS<sup>1</sup>, WILLIAM C. BURNETT<sup>2</sup>, YAACON NATHAN<sup>3</sup>, FAYEZ S. M. ALMBAYDIN<sup>4</sup>,  
ABDEL K.M. ATTIA<sup>5</sup>, LILIANA N. CASTRO<sup>6</sup>, RENÉ FLICOTEAUX<sup>7</sup>, M. EZZELDIM HILMY<sup>8</sup>,  
VIQAR HUSAIN<sup>9</sup>, ABDEWAHAB A. QUTAWNAH<sup>4</sup>, AFAT SERJANI<sup>10</sup> & YURI N. ZANIN<sup>11</sup>

**Key words:** Phosphorite, phosphate, francolite, mineralogy, geochemistry, trace elements, REEs, stable-isotopes, Sr-isotopes, Nd-isotopes, uranium, diagenesis, palaeoceanography, environment, toxic metals, fertiliser, phosphogypsum, radionuclide

## ABSTRACT

Major advances in our understanding of the genesis and diagenesis of marine phosphate deposits (phosphorites) have been made by recent geochemical studies. This work is reviewed and considered in relation to environmental issues arising from the exploitation of phosphorite deposits. Unaltered phosphorites contain francolite, a highly substituted carbonate fluorapatite, as the only phosphate mineral. It is the mineralogy of this phase, together with kinetic and thermodynamic factors, which govern many of the primary geochemical characteristics of phosphorites, although metamorphism or weathering may fundamentally alter the composition of a deposit. Francolite precipitation is inhibited by Mg in seawater, but occurs nonetheless in PO<sub>4</sub>-rich porewaters via the formation of a metastable precursor. Phosphate originates largely from the breakdown of organic matter in surficial sediments, apatite precipitating within cm–dm of the sediment/water interface. Coupled redox-cycling of Fe and P provides one means of maintaining high-PO<sub>4</sub> levels in sediments. Incorporation of trace elements, including Cd, U, Y, Zn and the rare-earth elements (REEs), occurs in association with the bacterially and redox-mediated degradation of particulate phases in the sediment, during or soon after francolite precipitation. REE distributions commonly reflect those in seawater. Uranium is incorporated largely as U(IV), but may be oxidised to U(VI) by seafloor weathering. Oxygen ( $\delta^{18}\text{O}$ ), carbon ( $\delta^{13}\text{C}$ ) and sulphur ( $\delta^{34}\text{S}$ ) stable-isotope studies of francolites provide a wealth of palaeoenvironmental information, and confirm that precipitation generally occurs under oxic to suboxic conditions in surficial sediments. Uranium- and  $^{14}\text{C}$ -dating together with

<sup>1</sup> School of Geological Sciences, Kingston University, Penrhyn Road, Kingston upon Thames, Surrey KT1 2EE, UK

<sup>2</sup> Department of Oceanography, Florida State University, Tallahassee, Florida 32306, USA

<sup>3</sup> Geological Survey of Israel, 30 Malkhe Israel Street, Jerusalem 95501, Israel

<sup>4</sup> Jordan Phosphate Mines Co Ltd, PO Box 30, Amman, Jordan

<sup>5</sup> Central Metallurgical Research Institute, PO Box 87, Helwan, Cairo, Egypt

<sup>6</sup> Departamento Ciencias Geológicas, Universidad Buenos Aires, Pab II, I Piso, 1428 Buenos Aires, Argentina

<sup>7</sup> Laboratoire de Géosciences de l'Environnement, Université d'Aix-Marseille III, St. Jérôme, F-13397 Marseille Cedex 20

<sup>8</sup> Department of Geology, Ain Shams University, Cairo 11566, Egypt

<sup>9</sup> Pakistan Council of Scientific and Industrial Research Laboratories, Jamrud Road, Peshawar, Pakistan

<sup>10</sup> Instituti i Studineve dhe Projekteve të Grjeologjisë, Blloku "Vasil Shasto", Tirana, Albania

<sup>11</sup> United Institutes of Geology, Geophysics and Mineralogy, Siberian Branch, Russian Academy of Sciences, 630090 Novosibirsk, Russia

Sr-isotope ratios ( $^{87}\text{Sr}/^{86}\text{Sr}$ ) may be used to measure nodule and grain growth rates and precisely date phosphogenic episodes. Nodules accumulate slowly at  $< 0.1\text{--}1 \text{ cm kyr}^{-1}$  while phosphate grains may form in  $< 10$  years. Variations in  $\epsilon_{\text{Nd}}$  values of francolite provide a basis for palaeoceanographic studies. Environmental issues associated with phosphate fertiliser production and use are discussed. Toxic metal, particularly Cd, and radio-nuclide contamination of the environment from phosphogypsum and fertilisers are considered. It is concluded that insufficient data are currently available for an accurate assessment to be made of the environmental impact and consequent human health implications of such contamination. Future advances in phosphorite geology will necessitate further experimental work, well-integrated field and laboratory studies using established geological and geochemical methods, and the application of new high-sensitivity spatially resolved analytical geochemical techniques.

## ZUSAMMENFASSUNG

Neue geochemische Studien haben unser Verständnis für die Genese und Diagenese mariner phosphatischer Ablagerungen (Phosphorite) stark erweitert. Diese Arbeiten sollen in der vorliegenden Studie im Überblick dargestellt werden; insbesondere soll auch der Aspekt der Umweltbelastung betrachtet werden, welcher bei der Nutzung von Phosphatsedimenten entsteht. Einziges Phosphatmineral in unveränderten Phosphoriten ist der Frankolith, ein Karbonat-Fluorapatit mit hoher Ionen-Austauschfähigkeit. Es ist vor allem die Mineralogie in dieser Phase, zusammen mit kinetischen und thermodynamischen Faktoren, welche für viele der primären geochemischen Charakteristika von Phosphoriten verantwortlich ist, obwohl Metamorphose oder Verwitterung die Zusammensetzung solcher Sedimente massgeblich verändern können. Die Ausfällung von Frankolith wird durch Mg im Meerwasser verzögert; Frankolith erscheint aber trotzdem in PO<sub>4</sub>-reichen Porenwässern auf dem Umweg über einen metastabilen Vorläufer. Phosphat bildet sich hauptsächlich aus dem Zerfall organischen Materials in Oberflächensedimenten, wobei Apatit innerhalb weniger cm bis dm unterhalb der Sediment/Wasser-Grenze ausfällt. Ein gekoppelter Redox-Zyklus von Fe und P ist eines der Mittel, um im Sediment hohe PO<sub>4</sub>-Werte aufrecht zu erhalten. Der Einbau von Spurenelementen wie Cd, U, Y, Zn und seltenen Erden (REE) geschieht zusammen mit dem bakteriellen und Redox-gesteuerten Abbau von Feststoffen im Sediment während oder kurz nach der Bildung von Frankolith. Die Verteilung der REE's entspricht üblicherweise jener im Meerwasser. Uranium wird häufig als U(IV) eingebaut, kann aber durch die Verwitterung am Meeresboden auch zu U(VI) oxydiert werden. Studien über die Verteilung der stabilen Isotope von Sauerstoff ( $\delta^{18}\text{O}$ ), Kohlenstoff ( $\delta^{13}\text{C}$ ) und Schwefel ( $\delta^{34}\text{S}$ ) in Frankolith sagen sehr viel aus über das Paläomilieu; sie bestätigen aber auch, dass die Ausfällung im Allgemeinen unter oxydischen bis suboxydischen Bedingungen in den obersten Schichten des Sediments stattfindet. Die Datierung mit Hilfe von Uranium und  $^{14}\text{C}$  zusammen mit dem Sr-Isoptopenverhältnis ( $^{87}\text{Sr}/^{86}\text{Sr}$ ) kann benutzt werden, um Wachstumsraten an Körnern und Knollen zu messen und phosphogene Episoden genau zu datieren. Knollen wachsen langsam mit  $< 0.1\text{--}1 \text{ cm}/1000 \text{ J}$ , während Körner sich schon in weniger als 10 Jahren bilden können. Unterschiede im Gehalt an  $\epsilon_{\text{Nd}}$  im Frankolith bilden eine Basis für paläoozeanographische Studien. Die Umweltbelastungen, welche durch die Produktion und Verwendung von Phosphat-Düngemitteln entstehen, werden diskutiert. Durch giftige Metalle, speziell Cd und Radio-nuklide ergeben sich Umweltverschmutzungen, welche ebenfalls betrachtet werden. Sie wirken sowohl beim Gebrauch von Düngemitteln wie auch bei der Erzeugung von „Phosphorgips“ (Abfallprodukt der Düngemittelproduktion). Es zeigt sich, dass für eine genaue Abschätzung der Umweltbelastung und der mit einer solchen Kontamination verbundenen menschlichen Gesundheitsprobleme noch nicht genügend Daten vorhanden sind. Zukünftige Fortschritte bei der Phosphorit-Geologie werden weitere Experimente, gut aufeinander abgestimmte Feld- und Laboruntersuchungen mit bekannten geologischen und geochemischen Methoden, sowie die Anwendung von neuen, hochsensitiven, räumlich auflösenden analytischen geochemischen Techniken notwendig machen.

## Introduction

Phosphorus, potassium and nitrogen are all nutrients critical for plant growth. Of these, it may be argued that phosphorus is of primary concern in terms of long-term availability. Nitrogen is present in abundance as N<sub>2</sub> in the atmosphere, from which it is extracted naturally by certain plants. If necessary, it may be fixed artificially by well-known industrial processes. Potassium is available in many large-scale geological evaporite deposits, and can

be recovered from modern seawater if necessary. Only phosphorus, a critical life-sustaining nutrient, is unavailable except via the exploitation of geological phosphate deposits.

Agricultural crops are harvested, so to maintain yields phosphate must be added continually to replace that lost from the soil. With human population growth close to 2%, a doubling of agricultural output is required every 35 years, just to maintain current levels of nutrition. Thus, there remains a need to continually expand the phosphate raw-material resource base. The problem lies not so much in the limited nature of World resources, but in the geographic and economic distribution of producers and users. A decline in the use of fertilisers in recent years has seen a marked fall in World production of phosphate rock, from 156 Megatonnes in 1988 to only some 135 Mt in 1992, and an estimated 117 Mt in 1993. About 74% of the World output is provided by China, the FSU (countries of the former Soviet Union), Morocco, and the USA. Much of the production from Morocco, by far the World's largest exporter, and from other major suppliers in the Middle East, North and West Africa, is destined for Western European markets. The use of phosphate fertilisers in the undeveloped nations of the Third World remains minute, yet it is these very nations which have the most pressing need to increase crop yields. It remains essential, therefore, to establish programmes of co-operation between developed and developing countries in order to initiate improved exploration, mining and processing strategies which will locate and produce new local phosphate reserves at low cost.

Exploitable phosphate deposits (see Notholt et al. 1989 for a comprehensive review) may be subdivided into three types: (1) marine sedimentary rocks or "phosphorites", such as those of the Cretaceous-Palaeocene rocks of North Africa and the Middle East, and the Miocene-Pliocene of the SE United States; (2) apatite-bearing igneous rocks, particularly the carbonatite and other alkaline igneous complexes of the Kola Peninsula NW Russia, Goiás and São Paulo Brazil, and Phalaborwa South Africa; (3) insular deposits found on certain tropical islands of the Pacific and Indian Oceans. Of these, sedimentary phosphorites are by far the most important, constituting more than 80% of the total ore produced.

Commercial phosphorite deposits invariably have elevated phosphate contents, commonly expressed as  $P_2O_5$  concentrations. The detailed mineralogical and geochemical composition of a deposit are fundamental in determining its commercial viability, controlling the extraction, beneficiation and processing which must be adopted, and strongly influencing the environmental impact of its development. Furthermore, phosphorite geochemistry provides unique information concerning the age and genesis of a deposit, and the palaeoceanographic and other palaeoenvironmental conditions which prevailed during its formation.

Over the last two decades, major advances have been made in the acquisition and interpretation of elemental and isotopic geochemical data from sediments. This paper aims to present an overview of present knowledge concerning the inorganic geochemistry of marine phosphorites. Biological processes and aspects of the organic geochemistry of phosphorites are discussed in a companion paper by Krajewski et al. (1994), while sedimentological and palaeoceanographic factors are reviewed in more detail by Glenn et al. (1994b). Here, we will describe how geochemical methods may be employed to understand the processes which lead to their formation, we will consider how these data may be used to constrain regional and global palaeoceanographic and tectonic models, and we will discuss current environmental concerns relating to their exploitation.

## Mineralogy

Phosphorite is defined as a marine sedimentary rock with  $\geq 18\%$   $\text{P}_2\text{O}_5$  (Slansky 1979, 1989), so by definition, phosphate minerals are its major constituents. In practice, francoite, a carbonate-fluorapatite with  $>1\%$  F and appreciable amounts of  $\text{CO}_2$  (Sandell et al. 1939; McConnell 1973), is essentially the only phosphate mineral occurring in phosphorites unaltered by metamorphism or weathering.

Francolites are structurally and chemically complex (Gruner & McConnell 1937; Borneman-Starinkevich & Belov 1940, 1953; Altschuler et al. 1952; Smith & Lehr 1966; Lehr et al. 1968; McClellan & Lehr 1969; McArthur 1978a, 1985, 1990; McClellan 1980; Kolodny 1981; Nathan 1984; McClellan & Van Kauwenbergh 1990). The group may be represented by the general formula:  $\text{Ca}_{10-a-b}\text{Na}_a\text{Mg}_b(\text{PO}_4)_{6-x}(\text{CO}_3)_{x-y-z}(\text{CO}_3 \cdot \text{F})_y(\text{SO}_4)_z\text{F}_2$ . However, other substitutions (Tab. 1) occur in all of the apatite sites (the two Ca sites, the  $\text{PO}_4$  site and the F site). Not all substituents have the same valency as the original ion (e.g.  $\text{Na}^+$  substituting for  $\text{Ca}^{2+}$ ;  $\text{CO}_3^{2-}$  or  $\text{SO}_4^{2-}$  for  $\text{PO}_4^{3-}$ ), and coupled substitutions (e.g.  $\text{NaCO}_3^-$  for  $\text{CaPO}_4^-$  or  $\text{CO}_3 \cdot \text{F}^{3-}$  for  $\text{PO}_4^{3-}$ ) are generally proposed to maintain charge balance. Calcium deficiency has also been suggested (Gulbrandsen et al. 1966; Bonel et al. 1973) as an additional means of accomplishing the balance of charge. The substitution of  $\text{CO}_3^{2-}$  in apatites and the replacement of F by hydroxyl ions has been discussed by Vignoles et al. (1982). McArthur (1990) speculated that francoite may form with a  $\text{Na}/\text{OH}$  ratio of 1, with  $1/2$  of the OH being replaced by  $\text{H}_2\text{O}$ , and suggested that in unweathered francoites half of the F<sup>-</sup> atoms in the structural site are replaced by OH<sup>-</sup>. High F contents are then caused by different degrees of coupled substitution of  $\text{CO}_3 \cdot \text{F}^{3-}$  for  $\text{PO}_4^{3-}$ . However, such a high level of hydroxyl substitution has yet to be proven analytically. Finally, there are other substitutions which do not change the charge balance, such as  $\text{Mg}^{2+}$  and  $\text{Sr}^{2+}$  for  $\text{Ca}^{2+}$ , which nevertheless have considerable crystallographic and geochemical importance.

Minerals such as quartz (e.g. North Carolina, Florida phosphorites of the SE United States) and calcite (Tethyan deposits of North Africa) are major components in most phosphorite deposits. Silica also commonly occurs as opal-CT and chalcedony, forming

Tab. 1. Possible substitutions in the francoite structure

Constituent ion	Substituting ion
$\text{Ca}^{2+}$	<b><math>\text{Na}^+</math>, <math>\text{K}^+</math>, <math>\text{Ag}^+</math></b> <b><math>\text{Mg}^{2+}</math>, <math>\text{Sr}^{2+}</math>, <math>\text{Ba}^{2+}</math>, <math>\text{Cd}^{2+}</math>, <math>\text{Mn}^{2+}</math>, <math>\text{Zn}^{2+}</math></b> $\text{Bi}^{3+}$ , $\text{Sc}^{3+}$ , $\text{Y}^{3+}$ , REE <sup>3+</sup> $\text{U}^{4+}$
$\text{PO}_4^{3-}$	<b><math>\text{CO}_3^{2-}</math>, <math>\text{SO}_4^{2-}</math>, <math>\text{CrO}_4^{2-}</math></b> <b><math>\text{CO}_3 \cdot \text{F}^{3-}</math>, <math>\text{CO}_3 \cdot \text{OH}^{3-}</math>, <math>\text{AsO}_4^{3-}</math>, <math>\text{VO}_4^{3-}</math></b> $\text{SiO}_4^{4-}$
$\text{F}^-$	$\text{OH}^-$ , $\text{Cl}^-$ , $\text{Br}^-$ $\text{O}^{2-}$

REE = rare-earth elements; compositionally significant substitutions are indicated in bold

porcelanites and cherts which together with organic matter and phosphorite, constitute the so-called P-C-Si trilogy (Bentor 1980). Many other phases occur in phosphorites, notably dolomite, glauconite, sulphides (pyrite), sulphates (gypsum), clinoptilolite, clay minerals (illite, kaolinite, palygorskite, smectites) and organic matter. Clearly, the proportions, distributions, grain sizes and textural characteristics of such accessory phases have important implications for the extraction and beneficiation of a deposit.

## Chemistry

Francolite chemistry may be affected by a number of factors, including: (1) kinetic effects due to the rate of formation; (2) thermodynamic factors; (3) precipitation from solutions of different composition due to secular variation in seawater or evolving porewater chemistry; (4) mechanism of formation, such as differences resulting from precipitation directly from solution or via dissolution and replacement of a pre-existing mineral (most commonly calcite), or bacterially mediated versus 'inorganic' precipitation; (5) post-precipitation burial diagenetic or metamorphic alteration; (6) weathering. The relative importance of these factors is still not well-constrained, but their significance will become apparent in the ensuing discussion.

Over the last decade, new information concerning the geochemical processes which lead to francolite precipitation and the formation of phosphorites has been derived from three main sources: (1) pore-water and solid-phase geochemical studies of sediments from modern phosphogenic areas, particularly the Mexican, Peru and East Australian shelves (Froelich et al. 1983; Jahnke et al. 1983; Glenn et al. 1988, 1994a; Van Cappellen & Berner 1988; Glenn 1990a; Heggie et al. 1990; O'Brien et al. 1990); (2) oxygen, carbon and sulphur stable-isotope studies of modern and ancient phosphorites (see Kolodny & Luz 1992a for a recent review); (3) geochemical and mineralogical data, and modelling of inorganic and bacterially-mediated apatite precipitation under experimental conditions (Lucas & Prévôt 1981, 1984, 1985; Jahnke 1984; Prévôt & Lucas 1986; Van Cappellen & Berner 1988, 1991; Prévôt et al. 1989).

### *Major elements*

The major-element geochemistry of unaltered francolites display surprisingly little variation; unweathered Cenozoic francolites (McArthur 1978a, 1980, 1985, pers. comm. 1994) contain 32%  $P_2O_5$ , 52%  $CaO$ , 4% F, and typically include (%):  $1.2 \pm 0.2 Na$ ;  $0.25 \pm 0.02 Sr$ ;  $0.36 \pm 0.03 Mg$ ;  $6.3 \pm 0.3 CO_2$ ;  $2.7 \pm 0.3 SO_4$ . In general, the level of substitution shown by the mineral decreases progressively with increasing age, burial diagenesis and/or weathering, which promote a transition towards unsubstituted fluorapatite  $[Ca_{10}(PO_4)_6F_2]$  compositions. With the onset of greenschist facies metamorphism, hydroxyfluorapatites may develop (Da Rocha Araujo et al. 1992; Girard et al. 1993). Subsequently, under extreme weathering conditions, Fe- and Al-phosphate minerals such as crandallite, millisite, wavellite and strengite commonly form (Zanin 1968; Altschuler 1973; Lucas et al. 1980; Schwab & Oliveira 1981; Flicoteaux 1982; Flicoteaux & Lucas 1984; Bonnot-Courtois & Flicoteaux 1989).

The major-element chemistry of phosphorites reflects both the composition of francolite and that of accessory minerals. Consequently,  $SiO_2$  for example, is highest in quartz-

rich deposits, such as the Miocene phosphorites of the SE United States. This is exemplified in Table 2, which shows the major-element composition of three phosphorite ore concentrates: Western Phosphate Rock from the US Permian Phosphoria Formation (NIST 694); Moroccan Phosphate Rock (BCR 32, Eocene); Florida Phosphate Rock (NIST 120c, Miocene). The much higher proportions of quartz and clay minerals in the Florida sample are clearly indicated by elevated levels of  $\text{SiO}_2$  and, to a lesser extent,  $\text{Al}_2\text{O}_3$ .

#### Carbonate and fluorine substitution

The vast majority of unweathered seafloor phosphorites have francolite- $\text{CO}_2$  contents (generally determined using the XRD-method of Gulbrandsen 1970; see Schuffert et al. 1990 for a revised equation) of around 6.3%, close to the maximum substitution allowable by the mineral structure (McClellan & Lehr 1969; McClellan 1980; Jahnke 1984; McClellan & van Kauwenbergh 1990). However, significant variation in francolite- $\text{CO}_2$  contents have been observed in many ancient and some modern phosphorites. The cause of this variation has been the focus of considerable debate.

Gulbrandsen (1970), studying the Phosphoria Formation of Idaho and Wyoming, noted that the average  $\text{CO}_2$ -contents of francolites taken from carbonate-phosphorite deposits in central Wyoming are moderately high (3.4%  $\text{CO}_2$ ), but they decrease progressively westwards towards SE Idaho with the transition to mudstone-chert-phosphorite facies (1.2%). He suggested that this systematic variation related to primary differences in dep-

Tab. 2. Major-element composition of phosphorite ore concentrates

Element	NIST 120c Western	BCR 32 Moroccan	NIST 694 Florida
	(%)		
$\text{SiO}_2$	5.5	2.09	11.2
$\text{TiO}_2$	0.103	0.0285	0.11
$\text{Al}_2\text{O}_3$	1.30	0.55	1.8
$\text{Fe}_2\text{O}_3$	1.08	0.231	0.79
$\text{MnO}$	0.027	0.0024	0.0116
$\text{MgO}$	0.32	0.403	0.33
$\text{CaO}$	48.02	51.76	43.6
$\text{Na}_2\text{O}$	0.52	n.d.	0.86
$\text{K}_2\text{O}$	0.147	n.d.	0.51
$\text{P}_2\text{O}_5$	33.34	32.98	30.2
$\text{CO}_2$	3.27	5.10	n.d.
F	3.82	4.04	3.20
$\text{SO}_3$	0.92	1.84	n.d.

Data from Potts et al. 1992b; BCR = Community Bureau of Reference, Belgium; NIST = National Institute of Standards and Technology, USA; total Fe expressed as  $\text{Fe}_2\text{O}_3$ ; n.d. = not determined

ositional environments, predominantly temperature differences causing vary  $\text{CO}_2$ -saturation in the water column. Later, McArthur (1985) expressed the view that this regional variation could be ascribed solely to late diagenetic effects having caused greater loss of structural carbonate from more deeply buried western sites.

There is no doubt that burial diagenesis, metamorphism and weathering generally cause the loss of  $\text{CO}_2$  and other constituents from the francolite structure (Jacob et al. 1933; Bliskovskiy 1976; Gusev et al. 1976, Zanin & Krivoputskaya 1976; Matthews & Nathan 1977; Lucas et al. 1980; McArthur 1980, 1985; McClellan 1980; Zanin et al. 1985; McClellan & Saavedra 1986; Panczer et al. 1989; Da Rocha Araujo et al. 1992). During deep-burial (km-scale), an inverse relationship between  $\text{CO}_2$ -content and temperature develops (Kastner et al. 1990), although later stage,  $\text{CO}_2$ -rich secondary carbonate-hydroxyfluorapatites have been reported (Da Rocha Araujo et al. 1992) from deposits affected by low-grade metamorphism. In the case of Phosphoria Formation francolites, a strong correlation between  $\text{SO}_4^{2-}$  concentration and  $\delta^{34}\text{S}$ , and the consistency of  $\delta^{18}\text{O}$  data (Piper & Kolodny 1987; see below for further discussion of isotopic data), both suggest that these isotopes reflect initial conditions of deposition, so the degree of alteration cannot be high. Furthermore, Upper Cretaceous-Palaeogene phosphorites from Israel (Nathan et al. 1990) display similar variations in structural  $\text{CO}_2$  to the Phosphoria Formation, with francolites from well-oxygenated carbonate-rich depositional environments exhibiting greater substitution (5.5%  $\text{CO}_2$ ) than those from granular phosphorites associated with anoxic facies (3.7%) in the same region. Differential weathering does not appear to be responsible for these trends, pointing to a primary geochemical difference between the areas.

Namibian seafloor phosphorites exhibit a range of phosphate- $\text{CO}_2$  contents from 1.4% in poorly consolidated nodules to 4.2% in indurated nodules and pellets (Romankevich & Baturin 1972; Price & Calvert 1978). This may be due to the fact that these young phosphorites include varying amounts of a francolite precursor phase (see below) which contains little or no structural  $\text{CO}_2$ . Another potential cause of such variation has been identified from studies of Holocene phosphorites off Peru (Glenn et al. 1988; Glenn 1990a), which contain between 2.1 and 6.0%  $\text{CO}_2$ . A correlation between lighter carbon stable-isotope compositions (more negative  $\delta^{13}\text{C}$ ; see below) with increasing  $\text{CO}_2$  contents in francolites, led Glenn and co-workers to conclude that carbonate substitution depends mainly on the carbonate alkalinity (and hence  $\text{CO}_3^{2-}$  concentrations) of the pore waters, which increases progressively with depth in the sediment. Rather surprisingly, pH is relatively constant despite the rapid generation of metabolic  $\text{CO}_2$ , because pH is buffered by ammonia production and increases in total alkalinity. The constancy of pH observed in the Peru margin sediments is important, not least because  $\text{HCO}_3/\text{CO}_3^{2-}$  ratios are affected considerably by pH, and are not just dependant on total alkalinity. Indeed, a relatively small fall in pH may reverse the relationship seen between total alkalinity and  $\text{CO}_3^{2-}$ .

In the organic-rich (up to 20%  $\text{C}_{\text{org}}$ ) sediments of the Peru-Chile and Mexican margins, francolite formation is restricted to within 5–20 cm of the sediment/water interface (Fig. 1, 2), because precipitation ceases at the high carbonate alkalinities which develop rapidly in deeper sediments. Precipitation is probably limited by the ability of the mineral to incorporate carbonate, so elsewhere, in less organic-rich sediments with shallower alkalinity gradients, francolite may form at greater depths. In one Peru margin nodule,




Fig. 1. Porewater and solid-phase geochemistry for slowly accumulating ( $1.7 \text{ cm kyr}^{-1}$ ) sediments in a modern upwelling area on the Peru continental margin, 230 km NW of Lima. Black arrows at the tops of graphs are measured seawater values; white arrows are estimated concentrations. The porewater  $\text{PO}_4$  profile indicates production of dissolved phosphate in the uppermost 10 cm of sediment and active phosphate precipitation around and below that depth (stippled areas). Fluorine decreases progressively below the sediment/water interface, suggesting that it is supplied to the precipitating phosphate largely by diffusion from seawater. Calcium and Mg contents do not differ significantly from those in seawater. Nitrate,  $\text{SO}_4$  and total alkalinity (TA) profiles demonstrate that phosphate is precipitating in suboxic (i.e.  $\text{NO}_3^-$ ,  $\text{NO}_2^-$  reducing, non-sulphidic), mildly alkaline conditions. Significant levels of Mn, Fe or  $\text{H}_2\text{S}$  were not detected. Solid-phase geochemical profiles (dashed lines) are consistent with rapid degradation of organic matter (declining  $\text{C}_{\text{org}}$  contents) and precipitation of phosphate (increasing total P and F) below 5 cm. Biogenic opal is virtually absent in this core. Precipitation is apparently occurring both within and above a bed of phosphorite clasts and grains (data from: Froelich et al. 1988; Glenn 1990a).

increasing  $\text{CO}_2$  substitution could be correlated with decreasing growth rate. This may simply reflect the increasing solubility of more carbonate-substituted apatite (Chien & Black 1976; Jahnke 1984), but it may also point to a kinetic effect limiting substitution in rapidly precipitated francolites.

The availability of dissolved fluoride undoubtedly plays an important role in regulating the precipitation of francolite (Froelich et al. 1983, 1988; Jahnke et al. 1983; Van Cappellen & Berner 1988; Ruttenberg & Berner 1993). As with structural  $\text{CO}_2$ , the F contents of francolites formed in organic-poor glauconitic carbonate sediments of low productivity areas such as the East Australian Shelf and Agulhas Bank off South Africa, are significantly higher than those typical of francolites from organic-rich muds from high productivity areas off Namibia, Peru and Mexico (Romankevich & Baturin 1972; Price & Calvert 1978; O'Brien et al. 1990). Average F/ $\text{P}_2\text{O}_5$  ratios of phosphorites from carbonate-dominated areas are close to those occurring in 'ideal' francolite; in addition to containing less  $\text{CO}_2$ , those associated with more organic-rich sediments typically contain up to one-third less F. This broad correlation between carbonate and fluoride contents is readily explained by the substitution of  $\text{CO}_3 \cdot \text{F}^{3-}$  for  $\text{PO}_4^{3-}$  in francolite (Borneman-Starinkevich & Belov 1940, 1953; Smith & Lehr 1966; McClellan & Lehr 1969; Price & Calvert 1978; Bacquet et al. 1980; McArthur 1990). Indeed, the relationship appears to be linear in unweathered francolites (McArthur 1990 Fig. 1), scatter being introduced by alteration causing the preferential loss of  $\text{CO}_2$  (and  $\text{OH}^-$  ?) from the mineral.


It is evident that the amount of  $\text{CO}_2$  and F substitution in francolite does indeed vary with depositional/early diagenetic environment, and it is speculated that high carbonate alkalinites and slow precipitation rates favour the incorporation of both constituents. In general, carbonate-associated phosphorites contain more structural  $\text{CO}_2$  and F than those forming in organic-rich mudrocks. This may be due to differences in kinetics, 'replacement' being much slower than 'direct' precipitation. However, both components may be lost during deep-burial diagenesis, metamorphism and/or weathering which can substantially modify the primary signature.

#### Magnesium incorporation and inhibition effects

In contrast to  $\text{CO}_3$  and F, unaltered francolites display very little variation in their Mg, Na, Sr and  $\text{SO}_4$  contents (McArthur 1978a, 1985). Unweathered francolites typically contain around  $3600 \mu\text{g g}^{-1}$  Mg, but early experimental work (Bachra et al. 1965; Martens & Harris 1970; Atlas 1975; Atlas & Pytkowicz 1977; Nathan & Lucas 1976; Savenko 1978) clearly demonstrated that apatite precipitation is inhibited by Mg concentrations present in seawater and marine porewaters ( $1300 \mu\text{g g}^{-1}$ ). This effect has been explained as the result of distortion of the embryonic apatite lattice by  $\text{Mg}^{2+}$  ions replacing  $\text{Ca}^{2+}$  promoting a wagnerite rather than apatite structure, a distortion strong enough to inhibit further crystal growth. Recent experimental data (Van Cappellen & Berner 1991), however, indicate that it is more likely to be caused by blocking of Ca sites by hydrated Mg-complexes.

Burnett (1977), amongst others, has argued that the removal of Mg by the formation of authigenic minerals such as dolomite or palygorskite must be necessary for francolite precipitation. However, many phosphorites contain no significant Mg-bearing phase, while dolomite may itself be replaced by apatite (Cook pers. comm. 1994), suggesting that Mg removal is not essential. Magnesium inhibition has been shown to be pH dependent (Nathan & Lucas 1976; Van Cappellen & Berner, 1991) with an increase in growth rate kinetics at pH 7 compared with pH 8, despite the thermodynamic disadvantage of increasing apatite solubility with decreasing pH (Atlas 1975; Atlas & Pytkowicz 1977).

Porewater data from modern phosphogenic sediments (Fig. 1, 2) show no depletion in  $Mg^{2+}$  below seawater values (Baturin, 1978; Jahnke et al. 1983; Froelich et al. 1988). Furthermore, although low pH of around 7.5 have been reported from both Namibian and Peru-Chile sediments (Baturin et al. 1970; Manheim et al. 1975), no relationship between low pH and sites of apatite precipitation has been established (Jahnke et al. 1983; Glenn et al. 1988).



Recent work has resolved some of these apparent inconsistencies. It has been demonstrated experimentally that apatite may precipitate from seawater via an amorphous, F-poor precursor such as struvite ( $\text{NH}_4\text{MgPO}_4 \cdot 6\text{H}_2\text{O}$ ; Malone & Towe 1970; Lucas & Prévôt 1985; Prévôt et al. 1989), amorphous calcium magnesium phosphate (ACMP; Froelich et al. 1988; Krajewski et al. 1994) and/or octacalcium phosphate ( $\text{OCP}_p$ ,  $\text{Ca}_4\text{H}(\text{PO}_4)_3 \cdot x\text{H}_2\text{O}$ ; Van Cappellen & Berner 1991). A precursor appears to be involved whether precipitation occurs directly from seawater or via the dissolution and 'replacement' of an existing mineral, such as calcite.

Support for a precursor in modern phosphogenic environments is provided by pore-water data from the Peru margin which display uncoupled  $\text{PO}_4$  and F removal (Froelich et al. 1988). Once precipitated, the metastable precursor gradually dissolves and acts as a substrate for apatite precipitation. Thus seawater is probably saturated with respect to francolite but the kinetics are too slow to allow its precipitation. The opposite is true of precursor phases which precipitate rapidly but only at elevated  $\text{PO}_4$  levels. Supersaturation with respect to the precursor most likely occurs intermittently within the interfacial pore-water phosphate maximum that characterises many phosphogenic areas (Fig. 1, 2; Krajewski et al. 1984 Fig. 8). The mechanism producing this maximum is not well understood, but may be related to the redox-sensitive uptake and release of phosphorus by microbial communities proliferating in the top sediment, the metabolic activity of sulphide-oxidising or other bacteria, suboxic bacterial degradation of organic matter, the dissolution of fish debris, and/or iron-redox cycling (see below). Organic compounds may also be instrumental in catalysing the precipitation process.

Temporary supersaturation will cause precipitation of a phosphate precursor which, once seeded, will promote apatite growth at much lower  $\text{PO}_4$  concentrations. Nonetheless, experimental studies consistently demonstrate that such apatites display poor crystallinity, and 'ageing' on a time scale of 10–100 years (Gulbrandsen et al. 1984) seems to be necessary to produce a typical francolite structure. Furthermore, growth kinetics

---

Fig. 2. Porewater and solid-phase geochemistry for rapidly accumulating ( $230 \text{ cm kyr}^{-1}$ ) sediments in a modern upwelling area on the Peru continental margin, 60 km west of Lima. Black arrows at the tops of graphs are measured seawater values; white arrows are estimated concentrations. Porewater profiles indicate intense anoxic diagenesis below 10 cm, with high rates of sulphate reduction ( $\text{SO}_4$  depletion, evolution of  $\text{H}_2\text{S}$ ). Disequilibrium suboxic nitrate and Fe are present in the upper well-mixed 5–10 cm. An interfacial high in  $\text{PO}_4$ , combined with a marked minimum in the porewater profile around the base of the bacterial mat at 3–4 cm, suggest active phosphate regeneration above and precipitation within this interval (stippled area). Fluorine is sourced largely by diffusion from seawater. Calcium and Mg contents do not differ significantly from those in seawater at this depth. Below 5 cm, intense sulphate reduction produces exponential downward increases in total alkalinity (TA), total dissolved carbon (TDC), and by difference, total dissolved carbonate-ion ( $\text{CO}_3^{2-}$ ) concentrations. Porewaters rapidly develop highly depleted carbon stable-isotope signatures, down to  $-18\text{‰ } \delta^{13}\text{C}$ , indicating domination of the dissolved carbon reservoir by degrading organic matter. Phosphate precipitation is confined to the uppermost 5 cm of the sediment column and may be limited, at least in part, by high carbonate-ion concentrations preventing growth at greater depths (see text for discussion). Solid-phase  $\text{C}_{\text{org}}$  data demonstrate progressive regeneration of organic matter in the sediment under both suboxic and anoxic conditions. Variations in solid-phase inorganic phosphorus ( $\text{P}_{\text{in}}$ ) and F are caused by fluctuations in biogenic silica, which constitutes a major component of the sediment at this site. Phosphate grains were not identified in the core, the high sedimentation rate precluding the development of a phosphorite deposit (data from: Froelich et al. 1988; Glenn et al. 1988; Glenn 1990a).

initially limit crystal sizes to the order of 0.1–10 µm (Van Cappellen & Berner 1991). The multi-phase aspect of francolite precipitation, in particular, makes the experimental determination of thermodynamic and kinetic factors extremely problematic.

It must be concluded that although Mg depletion is helpful for the formation of crystalline apatite because it shortens the time required for crystallisation, it is not necessary and is probably uncommon during the formation of phosphorites.

### Strontium

Unweathered francolites typically contain around 2500 µg g<sup>-1</sup> of Sr (McArthur 1978a, 1985), although “average” phosphorite (Altschuler 1980) has only 750 µg g<sup>-1</sup>. Lucas et al. (1990) demonstrated that, under experimental conditions, the Sr content of apatite is largely thermodynamically controlled with varying compositions reflecting equilibrium precipitation from fluids of varying Sr content. The small variation in Sr and other major elements displayed by marine francolites probably results from the relatively constant composition of seawater (~7.6 µg g<sup>-1</sup> Sr) and shallow porewaters, and the absence of significant kinetic effects. Strontium isotope data (see below) confirm that although francolites undergo negligible diagenetic exchange of Sr, weathering promotes rapid Sr loss (Bliskovskiy et al. 1967; McArthur 1978a, 1985; Panczer et al. 1989); Na and SO<sub>4</sub> appear to be less mobile but are also lost.

McArthur et al. (1988) observed from their study of western South African phosphorites that although modern francolites all have consistent Sr/Ca ratios, significant differences could be identified in age groups of Tertiary samples. Graham et al. (1982), based on geochemical variation in the tests of Cenozoic planktonic foraminifera, suggested that the Sr/Ca ratio of seawater varied significantly during the Tertiary. This offers the possibility that francolite major-element geochemistry may provide information on secular variation in seawater elemental, as well as isotopic, composition. Unfortunately, to date, subsequent workers have been unable to duplicate Graham et al.’s (1982) curve, and published data for phosphorites are insufficient to test the hypothesis.

### Iron geochemistry – a key to phosphogenesis?

Francolite incorporates little Fe in its structure, and dissolved Fe-concentrations in oxic seawater are also very low, typically less than 40 µg g<sup>-1</sup>. However, iron oxyhydroxide-rich Neogene phosphorite nodules occur on a number of present-day continental shelves, including the Agulhas Bank, South Africa (Parker 1975; Parker & Siesser 1972), Danois Bank, Spain (Lucas et al. 1978), East Australian Shelf (Marshall & Cook 1980; O’Brien & Veeh 1980; O’Brien et al. 1986, 1987, 1990), and off Morocco (McArthur 1978b; Summerhayes & McArthur 1990). A close relationship between iron minerals and francolite occurs in many modern and ancient phosphorites (Glenn 1990b; Glenn et al. 1994a, b), but glauconite is generally the dominant iron mineral present, forming a characteristic greensand – nodular phosphorite facies association. Most Neogene Fe-rich nodules occur in glauconitic sands.

The strong positive correlation between Fe and P in Quaternary phosphorite-bearing sediments from the East Australian Shelf (Heggie et al. 1990; O’Brien et al. 1990), together with sedimentological, petrographic and porewater geochemical data, have led to the realisation that the diagenetic behaviour of the two elements may be closely linked.

The iron-redox model suggests that ferric oxyhydroxides scavenge F from seawater, and  $\text{PO}_4^{3-}$  released from organic matter by microbial decomposition under oxic and suboxic conditions within the top few cm of sediment (Fig. 3). These FeOOH particles are subsequently transferred deeper into the suboxic zone by burial and sediment-mixing processes where they dissolve and both F and  $\text{PO}_4^{3-}$  are released. Some of these are then taken up by apatite precipitation, much of the remainder diffusing upwards to be refixed by ferric oxyhydroxides in the oxic layer to become available for further recycling. Glauconite formation may be linked to the same process (Froelich et al. 1988; Carson & Crowley 1993), with a proportion of the porewater  $\text{Fe}^{2+}$  and  $\text{Si}(\text{OH})_4$  derived from the dissolution of biogenic opal, together with  $\text{K}^+$  and  $\text{Mg}^{2+}$  ions diffusing into the sediment from seawater, reacting with the aluminosilicate fraction to precipitate glauconite.




Fig. 3. Schematic diagram of the Fe-redox cycle. Light-stippled areas represent solid phases, black arrows are solid-phase fluxes. White-outlined black arrows indicate reactions, white arrows are diffusion pathways. Ferric oxyhydroxides precipitated in the water column and at the sediment/water interface scavenge  $\text{PO}_4$  and some F from seawater. During burial and mixing, microbial decomposition of organic matter utilises a sequence of electron acceptors in order of decreasing thermodynamic advantage. Oxygen is used first, followed by nitrate (and nitrite), manganese- and iron-oxyhydroxides and sulphate, in that order (Froelich et al. 1979). Degradation of organic matter liberates  $\text{PO}_4^{3-}$  to solution, and dissolution of FeOOH liberates  $\text{Fe}^{2+}$ ,  $\text{PO}_4^{3-}$  and  $\text{F}^-$ , causing elevated concentrations of these ions (e.g. Fig. 2) in porewaters. If sufficient levels are attained,  $\text{PO}_4^{3-}$  reacts with  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{SO}_4^{2-}$  and  $\text{F}^-$  ions diffusing into the sediment from seawater, and  $\text{CO}_3^{2-}$  derived from the oxidation of organic matter, to precipitate a precursor phosphate which subsequently recrystallises to francolite (see text for discussion). Excess  $\text{PO}_4^{3-}$  diffuses upwards towards the sediment/water interface where it is resorbed by ferric oxyhydroxides. Iron diffuses both downwards to be fixed as FeS under anoxic conditions, and upwards to the oxic layer where it is reprecipitated as FeOOH. The Fe-redox cycle, therefore provides an effective means of trapping  $\text{PO}_4^{3-}$  in the sediment and promotes the precipitation of francolite.

Iron-redox cycling of phosphorus probably occurs more generally in marine shelf sediments (Ruttenberg & Berner 1993), but does not normally lead to the precipitation of significant amounts of francolite. However, the permanent burial of iron-bound phosphorus probably takes place in a wide range of sedimentary environments, and may represent a major, and underestimated, phosphorus flux from the oceans (Berner et al. 1993; Krajewski et al. 1994). In the case of the East Australian Shelf, it is noteworthy that although sulphate reduction has been detected in anoxic microniches, francolite precipitation is essentially a suboxic (i.e. non-suphidic) process. Similarly, since glauconites contains both  $\text{Fe}^{2+}$  and  $\text{Fe}^{3+}$  (the latter predominating), precipitation of that mineral probably also occurs close to the oxic/suboxic interface.

The coupled cycling of Fe with  $\text{PO}_4^{3-}$  across the oxic/suboxic redox boundary (Fig. 3) means that off East Australia, apatite precipitation occurs almost entirely in the uppermost 20 cm of the sediment column, and only proceeds while nodules remain in the mixed layer, continuing for periods of more than 60 kyr. This process can produce phosphorite nodules containing up to 10%  $\text{Fe}_2\text{O}_3$ . Seafloor weathering of glauconites and other iron-bearing phases plus goethite precipitation cause further enrichment, with some seafloor conglomeratic deposits attaining 50%  $\text{Fe}_2\text{O}_3$ . However, such nodules display Na- and  $\text{SO}_3$ -depleted 'altered' francolite compositions.

It appears likely that Fe-redox cycling plays a major role in francolite precipitation in many organic-poor environments. The importance of such cycling is less clear in organic-rich deposits such as those on the Peru margin where  $\text{Fe}_2\text{O}_3$  contents are typically less than 0.5% (Glenn & Arthur 1988), and in ancient Fe-poor phosphorites such as the Upper Cretaceous phosphatic chalks of NW Europe (Jarvis 1992). Iron-redox cycling undoubtedly provides one method of concentrating  $\text{PO}_4^{3-}$  in porewaters. Whether absorption by ferric oxyhydroxides always provides the primary means of extracting  $\text{PO}_4^{3-}$  from seawater seems unlikely, given that high levels of: (a) phosphatic fish debris; (b) planktonic organic matter; (c) benthonic microbial organic matter, have been identified in many areas (see Krajewski et al. 1994 for further discussion), and all of these are capable of yielding high levels of  $\text{PO}_4^{3-}$  to the sediment.

### *Trace elements*

The 'open' apatite structure facilitates numerous trace-element substitutions (Tab. 1), and many elements are known to be enriched in phosphorites relative to "average" shale (Tab. 3). It is widely recognised (Gulbrandsen 1966; Tooms et al. 1969; Nathan et al. 1979; Altschuler 1980; Prévôt 1990) that an enriched trace-element group comprising Ag, Cd, Mo, Se, U, Y, Zn and the rare-earth elements (REEs: La, Ce, Pr, Nd, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, Lu) characterise the phosphorite facies. Several other elements including Br, Cu, Cr, I, Pb & V (Prévôt & Lucas 1979, 1980, 1985; Pacey 1985; Piper 1991; Jarvis 1992) also commonly show strong affiliation to the apatite fraction in sediments. Recently, mineral structural studies using electron-spin resonance (ESR) have confirmed that many trace elements may replace  $\text{Ca}^{2+}$  in the apatite structure, including  $\text{Pb}^{3+}$  (Gilinskaya 1993),  $\text{VO}^{2+}$  (Gilinskaya & Zanin 1983),  $\text{U}^{4+}$  and  $\text{Th}^{4+}$  (Gilinskaya 1991, 1993; Gilinskaya et al. 1993). However, two points have to be made:

Tab. 3. Concentration of trace elements ( $\mu\text{g g}^{-1}$ ) in marine phosphorites relative to "average" shale

Element	Average shale	Enrichment factor	Normal abundance	Depletion factor
Ag	0.07	30		
As	13		N	
B	100			6
Ba	580		N	
Be	3		N	
Cd	0.3	60		
Co	19			3
Cr	90		N	
Cu	45		N	
Ga	19			5
Hg	0.4			7
La	40	4		
Li	66			13
Mo	2.6	4		
Ni	68		N	
Pb	20	2		
Sc	13		N	
Se	0.6	8		
Sn	6			2
Sr	300	2		
U	3.7	30		
V	130		N	
Y	26	10		
Yb	2.6	5		
Zn	95	2		
Zr	160			2

After Altschuler (1980, Tab. 5); average shale values from Turekian & Wedepohl (1961);

N = 'normal' abundance, within a factor of 2 of the concentration in shale

- (1) trace elements are not necessarily located within the apatite structure; they may be adsorbed onto crystal surfaces or may be related to another mineral or phase, particularly organic matter and sulphides;
- (2) average values may be misleading, since very wide ranges of concentration exist in different deposits.

The study of trace elements is of commercial interest because of their importance as saleable by-products (U; REEs) or because of their toxicity (Cd; U-derived radionuclides), and their consequent detrimental influence on the environment either in fertilisers or in phosphogypsum. The following discussion will concentrate on members of these groups.

## Cadmium and Zinc

Cadmium and Zn constitute a pair of trace elements which are closely associated in most rock types. Cadmium is the trace-element which above all others is most enriched in phosphorites compared with other lithologies (Altschuler 1980; Slansky 1980, 1986; Batutin & Oreshkin 1984), with concentrations in the order of  $20 \mu\text{g g}^{-1}$  being common, compared with  $0.3 \mu\text{g g}^{-1}$  in "average" shale. Some deposits have even higher Cd contents. Upper Cretaceous-Palaeogene phosphorites from Senegal, Togo and parts of the Gafsa Basin of Tunisia (Menor 1975; Baechle & Wolstein 1984; Johnson 1987) contain averages of  $53\text{--}84 \mu\text{g g}^{-1}$  Cd, while the Middle Miocene phosphorites of the US Monterey Formation (Altschuler 1980) yield  $200 \mu\text{g g}^{-1}$  Cd!

Cadmium correlates strongly with Zn in many phosphorites, although the former element shows considerably less enrichment in comparison to other sedimentary rocks (Tab. 3), typically occurring at around  $200 \mu\text{g g}^{-1}$  (compared to  $95 \mu\text{g g}^{-1}$  for "average" shale). The Cd-Zn relationship is in some, but certainly not all, cases attributable to the presence of sphalerite (Altschuler 1980; Minster & Flexer 1991). Generally, the two elements do not correlate with either the francolite or organic-matter fractions (Prévôt 1990; Nathan et al. 1991; Piper 1991). Altschuler (1980) and Middelburg & Comans (1991), amongst others, have argued that Cd enrichment in phosphorites is due in large part to its substitution for  $\text{Ca}^{2+}$  in the apatite structure; in contrast, the much smaller ionic radius of  $\text{Zn}^{2+}$  makes incorporation of this element less favoured. However, it should be noted that Cd has a tendency to form covalent bonds, so it does not easily replace Ca in franco-lite.

Cadmium and Zn, and to a lesser extent many other trace-elements including Cu and Ni, display nutrient-like  $[\text{PO}_4^{3-}, \text{NO}_3^-, \text{Si(OH)}_4]$  distribution in the oceans. All of these elements are highly depleted in surface waters because of their preferential extraction by marine phytoplankton during photosynthesis (Martin & Knauer 1973; Boyle et al. 1976; Boyle 1981; Bruland 1983; Price & Morel 1990; Piper 1994), but are enriched at depth as bacterial degradation of organic matter in the water column returns them to solution. It has been shown that Cd may substitute for Zn in some metalloenzymes and can promote the growth of phytoplankton in Zn-limited environments. Cadmium shows the closest correspondence to  $\text{PO}_4$  profiles amongst all of the trace elements, and this has led to the suggestion that the Cd content of foraminiferal calcite may be used as a palaeoproductivity indicator.

The above observations, together with evidence that Cd may be concentrated further by marine bacteria (Gauthier et al. 1986), make a biogenic source of Cd in phosphorites very likely. Elemental ratios for the marine fractions of Cd, Cu, Ni and Zn in the Tertiary phosphatic sediments of both the US Monterey Formation, California (Piper & Isaacs 1994), and the San Gregorio Formation of Baja California, Mexico (Piper 1991), approach plankton values. Interestingly, the accumulation rates for these minor elements require a high rate of primary productivity, comparable to the modern Peru Shelf, which is compatible with the operation of an active upwelling regime during phosphogenesis. Interelement relationships and calculated accumulation rates for Cd, Cu, Zn, and possibly Ni, in the US Permian Phosphoria Formation of Idaho (Piper & Medrano 1994) are also consistent with a biological origin. Nonetheless, despite strong evidence for a marine plankton source of trace elements in these and other phosphorites, there remains consid-

erable uncertainty regarding their mineralogical affinities and the incorporation mechanisms involved.

### Rare-earth elements and yttrium

For more than a century (Cossa 1878), the REEs have been recognised as generally being enriched in phosphate deposits (McKelvey 1950; Altschuler et al. 1967) compared to other sedimentary rock types. "Average" phosphorite (Altschuler 1980) contains  $462 \mu\text{g g}^{-1}$  REEs plus  $275 \mu\text{g g}^{-1}$  Y, while "average" shale contains only  $220$  and  $30 \mu\text{g g}^{-1}$  respectively. The bulk of the REEs and Y apparently reside in francolite, with the substitution of  $\text{REE}^{3+}$  and  $\text{Y}^{3+}$  for  $\text{Ca}^{2+}$  (Tab. 1) in one or both of the 7- and 9-fold co-ordination sites. Generally, granular phosphorites exhibit higher REE/P ratios than their nodular equivalents (Ilyin & Ratnikova 1976; McArthur & Walsh 1984; Jarvis 1984, 1992; Piper et al. 1988). Moreover, a range of distribution patterns is known, and differences in concentration of more than two order of magnitudes have been reported.

Traditionally, phosphorites have been considered to be characterised by a so-called "seawater" pattern (Goldberg et al. 1963; Altschuler et al. 1967), which is strongly depleted in Ce and enriched in the heavy REEs (HREEs; Ho-Lu) when compared with shales. This pattern, and particularly Ce depletion, is a feature of many oceanic sediments, and for comparative purposes is commonly expressed as an anomaly, calculated (Elderfield & Greaves 1982) as:

$$\text{Ce}^* = \log [3\text{Ce}_{\text{sn}}/(2\text{La}_{\text{sn}} + \text{Nd}_{\text{sn}})] \quad (1)$$

where "sn" represents a concentration normalised to shale.

The typical seawater-like pattern of phosphorites (Fig. 4) would suggest that REEs in francolites were incorporated from seawater with no significant fractionation. Many important deposits, including the western US Permian Phosphoria Formation, Upper Cretaceous and Palaeogene Abu Tartur and Nile Valley deposits of Egypt, Gafsa-Metlaoui Basin of Tunisia, and Ganntour Basin of Morocco, and the late Tertiary Bone Valley Formation of Florida, (Altschuler et al. 1967; El-Kammar et al. 1979; McArthur & Walsh 1984; Tlig et al. 1987), do indeed have markedly negative Ce-anomalies, although not all display HREE enrichment.

Much current phosphorite research is dealing with REE distributions and their significance. Jordanian (Ruseifa and Eshidiya deposits; Jallad et al. 1989, Abu Murry 1993) and French (Picardy phosphatic chalks; Jarvis 1984, 1992) Upper Cretaceous phosphorites (Fig. 4), for example, are enriched by a factor of two in REEs relative to 'average' shale, and have typical seawater-like distributions, with average Ce-anomalies of  $-0.5$  to  $-0.6$  and enrichment in HREEs. Eocene-Miocene deep-water seamount phosphorites from the equatorial Pacific (Hein et al. 1993) also exhibit large negative Ce-anomalies. Phosphatic rocks from the western part of the Palaeogene Senegal Basin display seawater-like patterns, the details of which appear to correlate to palaeowater-depth (Bonnot-Courtois & Flicoteaux 1989; Flicoteaux et al. 1993). Cape de Naze samples of earliest Eocene age have the largest negative Ce-anomaly ( $-0.78$ ) and represent a deep-water facies. Lam Lam – Taiba samples of Middle Eocene to Oligocene age seem to be related to transitional environments between platform and deep water, and have lower Ce-




Fig. 4. Shale-normalised rare-earth element (REE) distributions for selected phosphorites. Seawater-like patterns with negative Ce-anomalies and enrichment in the heavy REEs (top left) are typical of many phosphorites, but flat shale-like patterns (bottom left) are also common (see text for discussion). Shale values (USGS shale reference material SCo-1) from Potts et al. (1993b). Seawater data from Elderfield & Greaves (1982); phosphorite data from McArthur & Walsh (1984), Jarvis (1992; N. France samples) and Martin-Algarra et al. (1993; Spanish sample); conodont data from Grandjean-Lécuyer et al. (1993).

anomalies of between  $-0.62$  and  $-0.72$ . Similar depth-related trends have been identified (Grandjean et al. 1987) in REE data from Upper Cretaceous – Middle Eocene fish teeth from Moroccan phosphorites, and have been suggested, but not yet corroborated, for the Oligo-Miocene deposits of Baja California Sur, Mexico (Piper 1991). These trends are consistent with those shown by dissolved REEs in modern seawater. Modern coastal and ocean-surface waters generally have little or no negative Ce-anomaly and little HREE enrichment (e.g. Elderfield & Greaves 1982; Bertram & Elderfield 1993), but both features develop progressively with depth (Fig. 4) to form more typical “seawater” patterns by around 1 km in the open ocean.

With the notable exception of some phosphatised limestones, most Miocene – Holocene offshore phosphorites from Namibia, Cape Province and Agulhas Bank of South Africa (McArthur & Walsh 1984; Nathan et al. 1993), and the Peru margin (Piper et al. 1988), have shale-like REE distributions (Fig. 4) with no Ce anomaly. Phosphatic micro-

stromatolites from the Lower Cambrian in the Algerian Sahara (Bertrand-Sarfati et al. 1993) and coeval deposits at Karatau, Kazakhstan (Semenev et al. 1962) have similar patterns. Unusually, stromatolite-related Upper Jurassic phosphorites from Spain (Martin-Algarra et al. 1993) display a variety of REE distributions ranging from relatively flat shale-like patterns to those with large positive Ce-anomalies (Fig. 4) of up to +1.27. In the latter case, the REEs appear to be related to iron and manganese oxyhydroxide phases in the phosphorite.

Variations in Ce anomalies might in some cases be controlled by oxygenation levels (Elderfield & Pagett 1987; Wright et al. 1987), since fish remains from anoxic basins and shelves exhibit very small negative to moderate positive Ce-anomalies, while well-oxygenated outer shelf and deep-ocean deposits show moderate to large negative Ce-anomalies. However, local positive Ce-anomalies of +0.5 have also been identified in suboxic porewater underlying seawater with a negative anomaly (Elderfield & Sholkovitz 1987), so the degree to which Ce anomalies are controlled by seawater as opposed to porewater composition and redox-state remains open to debate.

Many biogenic phosphates, including Upper Palaeozoic conodont elements (Wright et al. 1987; Grandjean & Albarède 1989; Grandjean-Lécuyer et al. 1993), and both Triassic and Eocene fish teeth from France, show unusual REE patterns. These are strongly depleted by up to one order of magnitude in both the heavy and light REEs compared to the intermediate REEs, producing strongly concave-down patterns (Fig. 4) centred on Eu and Gd. A Devonian phosphate nodule from New York State in the US (McArthur & Walsh 1984) and Lower Carboniferous (Tournaisian; Notholt pers. comm. 1994) nodules from the Montagne Noire, France (Tlig et al. 1987), have similar patterns. Concave-down REE distributions have been reported (Hoyle et al. 1984) for  $\mu\text{m}$ -scale particulate and colloidal matter from modern river-seawater mixing zones, offering the possibility that the Palaeozoic phosphates acquired their REEs during diagenesis in estuarine or coastal settings. Unfortunately, this suggestion does not appear to be consistent with geological evidence, which indicates that the majority of samples originated from well-oxygenated shallow-water (<100 m) carbonate-shelf environments of normal marine salinity. Whether these patterns reflect unmodified seawater compositions remains controversial, but Nd-isotope studies (see below) support such a conclusion.

Recently, Grandjean-Lécuyer et al. (1993) have argued that prior to the evolution of modern open-ocean plankton during the early Cretaceous, the REE distribution of seawater was not controlled by ocean-surface biological productivity, but by inorganic processes associated with the scavenging of REEs by ferric oxyhydroxides. These authors argued that fractionation associated with these processes led to a characteristic concave-downwards profile in pre-Cretaceous seawater which was transferred unaltered to biogenic apatites. This suggestion seems oversimplistic. Not all pre-Cretaceous phosphorites display such patterns. Phosphoria Formation grains of Permian age, for example, have slightly HREE-enriched patterns with large negative Ce-anomalies (Fig. 4). On the other hand, Eocene fish teeth have been described (Grandjean & Albarède 1989) with strongly concave-downwards patterns. It must be concluded that although removal processes may promote different REE patterns in phosphate, such patterns are not necessarily global in distribution.

The REEs in francolite may be derived from: (1) seawater directly; (2) ferromanganese oxide-organic matter particles; (3) biogenic silica; (4) siliciclastic debris, particularly

clay minerals; (5) precursor carbonates. Simple mass-balance arguments indicate that given the relatively large size (and proportionally small surface area) of biogenic phosphate and phosphorite grains, and the very low concentrations ( $\text{fg g}^{-1}$ ) (femtogram =  $10^{-15}$  gram) of REEs in seawater, incorporation is unlikely to occur directly. It almost certainly takes place during early diagenesis, following the release of REEs from dissolving carrier phases, particularly sub- $\mu\text{m}$  scale ferromanganese oxyhydroxides particles and grain-coatings and organic matter (cf. Jarvis 1984, 1992; Sholkovitz et al. 1989), which have previously scavenged large quantities of REEs from the water column. Reduced sedimentation and burial rates favour increased concentrations of REEs in francolite (Jarvis 1984; Elderfield & Pagett 1986; Wright et al. 1987), since the longer the mineral remains close to the oxic/suboxic redox boundary, the longer it is able to incorporate REEs being liberated from their carriers. Concentration levels also appear to be related to grain-size and possibly to kinetic effects. In most cases, additional sources of REEs such as siliciclastic and biogenic detritus and precursor carbonate grains, matrix and cements are unlikely to provide the quantities of REEs necessary to produce the enrichment levels observed in phosphorites. Nonetheless, biogenic silica has been advocated as a source of REEs in some deposits (Piper 1991).

McArthur & Walsh (1984) suggested that francolites precipitate with very low REE contents, but subsequently scavenge REEs from other sources throughout their history, so old onshore deposits contain the highest REE contents. This hypothesis is supported by data from fish remains (teeth, vertebrae), which initially contain negligible REEs (Bernat 1975) and rapidly acquire high concentrations *post-mortem* (Grandjean et al. 1987). However, the resulting patterns appear to mimic those of the overlying water column (Wright et al. 1987), and do not appear to be modified significantly by interaction with formation or groundwaters during burial (Grandjean & Albarède 1989).

Finally, it has been shown (Bonnot-Courtois & Flicoteaux 1989; Dar'in & Zanin 1990; Flicoteaux et al. 1990) that weathering leads to the leaching, fractionation and redistribution of REEs, and may substantially modify the primary REE signature of a deposit. For any particular phosphorite, therefore, several factors are likely to have influenced the final REE content and signature. The mechanisms of REE incorporation appear to be closely tied to the organic matter degradation and redox-driven processes which, on current evidence, are so fundamental in driving phosphogenesis. However, the probability that REE patterns reflect not only their depositional environment, but also the imprint of diagenesis and weathering, clearly must be taken into account when interpreting data.

## Uranium

Uranium is another of the elements generally showing strong enrichment in phosphorites ( $120 \mu\text{g g}^{-1}$  in “average” phosphorite, compared with  $3.7 \mu\text{g g}^{-1}$  in “average” shale; Altschuler 1980). Indeed, the enrichment of phosphorites in U is so strong, that mapping the radioactivity of the seafloor using a deep-tow scintillometer (Jones 1989) is an effective way of determining the distribution of phosphatic sediments in many parts of the oceans.

Much research on U in phosphorites (McKelvey 1956; Altschuler et al. 1958) has been driven by the wish to find alternative U sources for military and civilian nuclear programmes. However, more recently, U has also been studied because of its applications in dating (Kolodny & Kaplan 1970a; Baturin et al. 1972; Veeh et al. 1973, 1974; Bur-

nett et al. 1982, 1988; Kim & Burnett 1986; Barole et al. 1987; McArthur et al. 1988), as an indicator of diagenesis (Starinsky et al. 1982), and as a potential palaeo-redox marker (Burnett & Gomberg 1977). Although there is generally a strong positive correlation between U and P contents for individual deposits (e.g. Jeanmaire 1985), it is unclear whether U is incorporated into the francolite lattice, adsorbed onto crystallite surfaces and/or associated with enclosed accessory phases, particularly organic matter. Fission-track mapping (Burnett & Veeh 1977; Kress & Veeh 1980; Avital et al. 1983) convincingly demonstrates that U is intimately associated with francolite on a microscopic scale, and ESR studies (Gilinskaya 1991, 1993; Gilinskaya et al. 1993) confirm that both  $U^{4+}$  and  $Th^{4+}$  substitute for  $Ca^{2+}$  in the structures of many natural apatites. However, in some deposits (Lucas & Abbas 1989) U is dominantly complexed by organic matter and may not be present in the apatite lattice in significant quantities.

Uranium occurs as both U(IV) and U(VI) in varying proportions in different phosphorites (Altschuler et al. 1958; Kolodny & Kaplan 1970a; Veeh et al. 1973). East Australian ferruginous phosphorite nodules (O'Brien et al. 1987), for example, range from 11–86% U(IV) and, in general, it appears that samples with high U/ $P_2O_5$  ratios tend to contain higher proportions of U(IV). Crystallographic arguments indicate that U(IV) would be most easily accommodated by the apatite lattice, and indeed there is some evidence that U is incorporated into francolite almost entirely in that form (Altschuler et al. 1958; O'Brien et al. 1987), and is only subsequently converted to U(VI) by oxidation processes. It has also been suggested that phosphorites form initially with high U/ $P_2O_5$  ratios, and that precipitation of later francolite in nodules is accompanied by little further U-up-take. Possible causes for this differential uptake of U include: (1) kinetic effects causing rapidly growing francolites to be depleted in U; (2) the diffusional supply of U being reduced as porosity declined during lithification; (3) a varying supply of U(IV), possibly due to differing redox conditions and availability of labile organic matter during initial and later precipitation of francolite. Uranium occurs in seawater around  $3\text{ ng g}^{-1}$ , but is fixed and concentrated by organic complexes. Oxidation of organic matter leads to remobilisation of U as soluble U(VI) species (Colley & Thomson 1985), which may diffuse to be reprecipitated as U(IV) at the suboxic/oxic redox interface. Consequently, it is likely that redox-cycling also plays an important role in controlling the U supply to phosphorites.

Uranium contents vary enormously between different deposits (Cathcart 1978, 1992; Slansky 1980, 1986). For example, despite typical phosphorite concentrations of  $>100\text{ }\mu\text{g g}^{-1}$  U, many Cambrian examples in Siberia contain only  $4\text{ }\mu\text{g g}^{-1}$  (Ilyin pers. comm. 1993), and relatively low U concentrations ( $5\text{--}12\text{ }\mu\text{g g}^{-1}$ ) appear to be common in many other Proterozoic-Cambrian phosphorites (e.g. Mongolia,  $2\text{--}11\text{ }\mu\text{g g}^{-1}$  U in the Khubsugul phosphorites, Ilyin et al. 1986; Upper Volta,  $4\text{--}8\text{ }\mu\text{g g}^{-1}$  U in the Tapoa deposits, Lucas et al. 1986; Kazakhstan,  $4\text{--}8\text{ }\mu\text{g g}^{-1}$  U in Karatau phosphorites, Ilyin pers. comm. 1993). Low U concentrations of this order are also typical of Tertiary seamount phosphorites (Kolodny 1981; Roe & Burnett 1985; Cullen & Burnett 1986; Hein et al. 1993). At the other extreme, Eocene deposits near Bakouma (Central African Republic) contain  $1660\text{--}5600\text{ }\mu\text{g g}^{-1}$  U (Gony 1971), and some Middle Palaeozoic phosphorites in Siberia yield up to  $4000\text{ }\mu\text{g g}^{-1}$  U (Gavshin et al. 1974). Such extreme variation is almost certainly a consequence of post-depositional diagenetic, hydrothermal, metamorphic and/or weathering effects, all of which may cause significant U mobilisation (Altschuler et al. 1958; Cook 1972; Cathcart 1978; Slansky 1980, 1986; Jeanmaire 1985; McArthur & Herczeg 1990).

Much remains to be understood about the locations and mechanism of incorporation of U in phosphorites. It is now clear, however, that high U concentrations and high proportions of U(IV) in phosphorites do not necessarily indicate that francolite precipitation occurred under anoxic conditions, nor do they equate directly with deposition in organic-rich sediments. Indeed, nodules precipitated in association with organic-rich muds on the Peru margin commonly contain less U and more U(VI) than those accumulating in glauconitic foraminiferal sands off East Australia.

### Isotope geochemistry

The chemistry and relative stability of francolite makes it perhaps *the* most versatile sedimentary mineral for isotope studies. Many of the stable and radiogenic isotopes which are used as tracers or dating devices in sedimentary rocks occur in francolite (Fig. 5, Tab. 4). The three most commonly used elements for stable-isotopic work, O, C and S, are all major structural components of francolite, while three major radiogenic isotope tracers Sr, Nd and U, are constituents that replace major elements in the lattice.




Fig. 5. Schematic diagram showing the location of different isotopes in francolite. Note the stable and radiogenic isotopes of the light elements are constituents of major anions in francolite, whereas the radiogenic isotopes of the heavy elements are trace substituents for the Ca cation.

### Oxygen stable-isotopes

The use of oxygen-isotope ratios in marine carbonate  $\text{CO}_3^{2-}$  to estimate water palaeotemperatures (Urey et al. 1951) and to determine diagenetic conditions (Hudson 1977; Irwin et al. 1977) is well-established. For most purposes the  $\delta^{18}\text{O}/\delta^{16}\text{O}$  ratio is expressed in delta ( $\delta$ ) notation as a per mille ( $\text{\textperthousand}$ ) deviation from an ocean water (SMOW – Standard Mean Ocean Water) or calcite (PDB – Pee Dee Belemnite) standard. Francolite offers the rare opportunity of determining the oxygen-isotope ratios of two cogenetic pairs in a single mineral,  $\delta^{18}\text{O}_p$  of  $\text{PO}_4^{3-}$  and  $\delta^{18}\text{O}_c$  of  $\text{CO}_3^{2-}$ .

The application of variations in the  $\delta^{18}\text{O}_p$  of phosphatic fish debris as an indicator of modern and ancient water temperatures has been demonstrated (Longinelli & Nuti 1973a, b; Kolodny et al. 1983; Kolodny & Raab 1988; Kolodny & Luz 1992b). The proposed phosphate-water temperature equation (Longinelli & Nuti 1973a) is:

$$T \text{ } ^\circ\text{C} = 111.4 - 4.3 (\delta^{18}\text{O}_p - \delta^{18}\text{O}_w) \quad (2)$$

where  $T$  is the average environmental temperature,  $\delta^{18}\text{O}_p$  is francolite phosphate and  $\delta^{18}\text{O}_w$  is that of the water, both on the SMOW scale. Revised empirical equations have been published by Karhu & Epstein (1986), Shemesh et al. (1988) and Kastner et al. (1990).

Tab. 4. Isotopes in francolite and their applications

Element	Parameter measured	Genesis of phosphorites	About the World
Ca-sites			
Sr	$^{87}\text{Sr}/^{86}\text{Sr}$	age	land – sea distribution
Nd	$\epsilon\text{Nd}$		land – sea distribution palaeoceanography
U	$^{234}\text{U}/^{238}\text{U}$ $^{230}\text{Th}/^{234}\text{U}$ $^{231}\text{Pa}/^{235}\text{U}$	rate of growth, age	
PO <sub>4</sub> -site			
O	$\delta^{18}\text{O}_p$ , $\delta^{18}\text{O}_c$	temperature of formation, diagenetic environment	palaeoclimate
C	$^{14}\text{C}$ $\delta^{13}\text{C}$	age diagenetic environment	
S	$\delta^{34}\text{S}$	diagenetic environment	

Modified from Kolodny & Luz (1992)

The  $\delta^{18}\text{O}_c$  equation for apatite is assumed to be the same as  $\text{CaCO}_3$  (Epstein et al. 1953):

$$T \text{ } ^\circ\text{C} = 16.5 - 4.3 (\delta^{18}\text{O}_c - \delta^{18}\text{O}_w) + 0.14 (\delta^{18}\text{O}_c - \delta^{18}\text{O}_w)^2 \quad (3)$$

where  $\delta^{18}\text{O}_c$  is francolite carbonate.

In both cases, the  $\delta^{18}\text{O}$  of the water has been known or estimated to obtain a unique palaeotemperature. Most offshore Tertiary and Quaternary phosphorites yield  $\delta^{18}\text{O}_c$  values (Kolodny & Kaplan 1970b; McArthur et al. 1980, 1986) which are consistent with precipitation in equilibrium with marine waters at palaeoceanographically reasonable water temperatures (typically 6–20 °C). Namibian concretions, for example, yield  $\delta^{18}\text{O}_c$  temperatures of 6–13 °C, which are within the range of temperatures (6–16 °C) measured for modern upwelling water on the margin. However, many onshore deposits (e.g. Jordanian Upper Cretaceous Al-Hasa deposit, US Permian Phosphoria Formation, Australian Middle Cambrian Georgina Basin phosphorites) exhibit anomalously light compositions which have been attributed to weathering effects caused by interaction with isotopically light (i.e.  $^{16}\text{O}$ -enriched) groundwaters (Shemesh et al. 1983; McArthur et al. 1986). By contrast, Piper & Kolodny (1987) argued that  $\delta^{18}\text{O}_p$  values from their Phosphoria Formation samples are unaltered and indicate high Permian palaeowater temperatures of 33.7–40.4 °C. Stratigraphic variation was interpreted as reflecting glacial-interglacial cyclicity and the consequent change in the  $\delta^{18}\text{O}$  composition of seawater. High Oligo-Miocene palaeowater temperatures of 27.5–32.7 °C were also suggested by Piper (1991) based on  $\delta^{18}\text{O}_p$  data from deposits of Baja California Sur, Mexico.

It is generally agreed that although  $\delta^{18}\text{O}_c$  is more susceptible to alteration than  $\delta^{18}\text{O}_p$  (Shemesh et al., 1983, 1988; McArthur & Herczeg 1990), isotopic exchange may occur

from both sites, and  $\delta^{18}\text{O}_p$  does not provide the perfect palaeothermometer it was once held to be. When phosphorites of different ages are compared (Longinelli & Nuti 1968; Shemesh et al. 1983, 1988; Karhu & Epstein 1986),  $\delta^{18}\text{O}_c$  and  $\delta^{18}\text{O}_p$  display clear, roughly parallel, exponential trends of decreasing values with age, the greatest changes occurring during the last 100 Myr. This major shift must be due in part to falling global temperatures since the Late Cretaceous, but pre-Cretaceous values are almost certainly altered. Carbon isotopes show no such relationship.

In some ancient phosphorites coexisting fish material and francolite cements have comparable  $\delta^{18}\text{O}_p$  values, which yield geologically reasonable palaeotemperatures. Shemesh & Kolodny (1988), for example, interpreted stratigraphic variation of  $\delta^{18}\text{O}_p$  in Upper Cretaceous Mishash Formation phosphorites and fish remains, as indicating that a cool-water upwelling episode was associated with a major phosphogenic event in SE Tethys. In other deposits, such as the Middle Miocene Monterey Formation of California (Kastner et al. 1990),  $\delta^{18}\text{O}_p$  values of fish debris are considerably higher (by up to 6‰) and much less variable than associated phosphorites, suggesting that significant post-depositional isotopic exchange has occurred in the latter. Only fish-derived data consistently indicate viable palaeowater temperatures of 11–25 °C.

Variation in the isotopic composition of O in response to temperature for the carbonate and phosphate sites in francolite, is sufficiently different to offer the possibility of determining palaeotemperatures from  $\delta^{18}\text{O}_c$  –  $\delta^{18}\text{O}_p$  pairs, independently of water compositions. Unfortunately, although trends for the two ratios are collinear, late diagenetic interaction with formation (Shemesh et al. 1988; Kastner et al. 1990) or meteoric (McArthur & Herczeg 1990) waters commonly lead to diagenetic rather than depositional palaeotemperatures being recorded. Particularly in organic-rich sediments, reequilibration may occur even at relatively low water/rock ratios, and it remains unproved that precipitation invariably occurs under equilibrium conditions. Consequently, the interpretation of phosphorite oxygen-isotope data remains highly problematic.

It must be concluded that the use of  $\delta^{18}\text{O}$  data to estimate seawater temperatures or isotopic compositions should be undertaken with the greatest caution. Available evidence indicates that biogenic apatites are less prone to late diagenetic overprint than coexisting grain and nodule francolite cements, although even the former are liable to be affected under some circumstances. Nonetheless, deposits which have never been buried deeply and have experienced little movement of formation waters may still retain a depositional signature.

### Carbon stable-isotopes

In addition to measuring  $\delta^{18}\text{O}$  in francolite  $\text{CO}_3^{2-}$ , it is also possible to determine the  $^{13}\text{C}/^{12}\text{C}$  ratio ( $\delta^{13}\text{C}$ , reported as a ‰ deviation from the PDB standard). Used together,  $\delta^{18}\text{O}_c$  and  $\delta^{13}\text{C}$  data (Fig. 6) are highly instructive concerning the physicochemical conditions of francolite precipitation. It is necessary to make several assumptions concerning the timing and mechanism of incorporation of  $\text{CO}_2$  in the apatite lattice:

- (1) little or no carbon-isotope fractionation occurs during incorporation of carbonate, so values are representative of dissolved  $\text{CO}_2$  in the water from which the mineral precipitated;




Fig. 6. Variation of  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  in modern and ancient francolite carbonate (based on McArthur et al. 1986 with Senegal data from Girard et al. 1993).

- (2) carbonate enters the lattice at the time of precipitation;
- (3) isotopic re-equilibration has not occurred subsequently.

By analogy to  $\text{CaCO}_3$  precipitation, carbon isotope fractionation is expected to be minimal. This assumption is supported by a preponderance of seawater-like  $\delta^{13}\text{C}$  values reported for francolites from Tertiary and Quaternary seafloor phosphorites (Kolodny & Kaplan 1970b; McArthur et al. 1980, 1986; Benmore et al. 1983, 1984; Shemesh et al. 1988). However, the range of  $\text{CO}_2$  contents exhibited by individual phosphorites suggests that precipitation is not instantaneous and occurs within a 'window' of evolving pore-water compositions. Furthermore, in the case of biogenic phosphates,  $\text{CO}_2$  is added *post mortem* during the transition of the original hydroxyapatite to francolite, which must take place by a dissolution-reprecipitation process.

The use of  $\delta^{13}\text{C}$  data to constrain depositional and early diagenetic environments hinges on our understanding of the isotopic fractionation which occurs during the microbial degradation of organic matter in sediments (Irwin et al. 1977; Froelich et al. 1979). Marine organic matter is highly depleted in  $^{13}\text{C}$ , typically around  $-24\text{‰}$   $\delta^{13}\text{C}$  PDB, and when oxidised it produces dissolved inorganic carbon (DIC) species ( $\text{CO}_2$ ,  $\text{HCO}_3^-$ ) of the

same isotopic composition. The amount of bicarbonate in porewaters is small, so carbon stable-isotopes in DIC are very sensitive to organic degradation processes which affect the sediment during burial (in contrast, oxygen isotope compositions are dominated by the water).

The proportion of organically-derived  $\text{CO}_2$  (Fig. 7) varies with depth as organic matter is oxidised by a succession of bacterial populations. In the oxic and suboxic upper layers of sediment, where oxygen and then nitrate, iron and manganese compounds are used sequentially as oxygen sources, there is generally a modest increase in DIC with depth, causing decreasing porewater  $\delta^{13}\text{C}$  values to a minimum of around  $-6\text{\textperthousand}$  (or  $-3\text{\textperthousand}$  if calcite dissolution accompanies  $\text{CO}_2$  production). Below this, under anoxic conditions,




Fig. 7. Schematic diagram showing the relationships between porewater  $\text{CO}_3^{2-}$ ,  $\delta^{13}\text{C}$ ,  $\text{SO}_4$  and  $\delta^{34}\text{S}$  with increasing depth in the sediment. The  $\text{CO}_3^{2-}$  curve assumes that pH is constant and dissolved inorganic carbon (DIC) increases linearly with depth. Porewater data from modern sediments (Fig. 1, 2) and stable isotope data from modern and ancient phosphorites, indicate that phosphogenesis is confined largely to the suboxic and uppermost part of the anoxic zones (heavily stippled area). The depth and thickness of each zone varies considerably between sites in response to differences in sedimentation rate, organic carbon flux, bioturbation rate, seawater oxygen content and other factors; all three zones may not be distinguishable at all sites (modified from McArthur et al. 1986).

sulphate reduction leads to increased bicarbonate production and a pronounced decrease in  $\delta^{13}\text{C}$ , down to  $-18\text{\textperthousand}$  (Fig. 2, 7).

When sulphate is exhausted, organic matter is decomposed by methanogenic micro-organisms. Acetate fermentation produces isotopically very light  $\text{CH}_4$  ( $<-60\text{\textperthousand}$   $\delta^{13}\text{C}$ ) and, at least in the initial stages,  $\text{CO}_2$  ( $-20\text{\textperthousand}$ , rising to  $-5\text{\textperthousand}$  as the reaction progresses). At the same time, part of the total dissolved  $\text{CO}_2$  will undergo carbon dioxide reduction by reaction with bacterially-derived hydrogen, producing additional methane which is also isotopically light ( $\text{CH}_4$  is  $25\text{--}60\text{\textperthousand}$  lighter than its precursor  $\text{CO}_2$ ). Consequently, cements precipitated in association with the oxidation of  $\text{CH}_4$  have very light carbon stable-isotope signatures. However, the reservoir effect associated with carbon dioxide reduction causes the residual  $\text{CO}_2$  to become progressively heavier, ultimately attaining values as high as  $+15\text{\textperthousand}$   $\delta^{13}\text{C}$ . In practice, such extremes are quite rare (Emery & Robinson 1993), and cements precipitated in the methanogenic zone are typically between  $+2$  and  $-22\text{\textperthousand}$ .

The carbon stable-isotope composition of francolites in replaced limestones will be influenced by the marine  $\delta^{13}\text{C}$  values of their carbonate precursor, since porewater bicarbonate will generally be dominated by that source. Even in this situation, however, modification of the isotopic signature should occur. The replacement of  $\text{CaCO}_3$  requires its dissolution, which is most likely a result of  $\text{CO}_2$  production caused by the breakdown of organic matter in the oxic zone. In this situation, the reaction should buffer the  $\delta^{13}\text{C}$  to a point half way between the  $\delta^{13}\text{C}$  of the  $\text{CO}_2$  and the  $\text{CaCO}_3$ , and the francolite precipitated should display a negative isotope signature that is the mean of the two. The signature can only remain relatively unmodified in systems with low water/rock ratios and low organic matter contents, where carbonate dissolution and francolite precipitation are not directly coupled. In organic-rich water-dominated systems even phosphatised limestones should acquire negative  $\delta^{13}\text{C}$  values. Finally, although carbon isotope ratios are much less prone to alteration during deep burial and weathering than oxygen isotopes, the loss of structural- $\text{CO}_2$  from weathered francolites must be cause for concern. Certainly, interaction between groundwaters charged with soil-derived DIC and francolite during lateritic weathering consistently leads to light carbon ( $-15$  to  $-10\text{\textperthousand}$   $\delta^{13}\text{C}$ ) and oxygen ( $+22$  to  $+27\text{\textperthousand}$  SMOW  $\delta^{18}\text{O}_c$ ) isotope values in secondary apatites (Girard et al. 1993).

Current isotopic evidence indicates that most phosphorites formed under dominantly oxic to suboxic conditions. For example, the negative correlation between francolite  $\delta^{13}\text{C}$  and  $\text{CO}_2$  values in Peru margin Holocene phosphorites (Fig. 8), with maximum  $\text{CO}_2$  contents of  $6\%$  corresponding to minimum  $\delta^{13}\text{C}$  values of around  $-3\text{\textperthousand}$  (Glenn et al. 1988; Glenn 1990a), are consistent with precipitation within 10–20 cm of the sediment/water interface in dominantly suboxic conditions. Most other young (late Tertiary – Quaternary) seafloor phosphorites, including those from the eastern US Blake Plateau, Chatham Rise of New Zealand, East Australian margin, and off South Africa, also display  $\delta^{13}\text{C}$  signatures within a few per mille of seawater values (Fig. 6). Isotopically lighter compositions (down to  $-15\text{\textperthousand}$ ) characteristic of anoxic diagenesis have been recorded only from late Pleistocene – Holocene nodules off Namibia, for Tertiary phosphorites from the Ica Plateau and Sechura Desert of Peru, the Miocene of Baja California Sur (Piper 1991), and southern Florida (Compton et al. 1993), and some onshore Upper Cretaceous – Palaeogene Moroccan samples. One other notable exception has been reported (McArthur et al. 1980; Benmore et al. 1984) from the Namibian shelf, which has yielded a F-poor hydrated calcium magnesium phosphate-cemented nodule (?coprolite) containing

29.5%  $P_2O_5$  and a phosphate- $\delta^{13}C$  value of +15.6‰. Such a heavy isotopic signature suggests that this is a rare example of phosphate precipitation during  $CO_2$  reduction in a methanogenic environment. The unusual mineralogy of this nodule corresponds to a possible francolite-precursor (see above), yet its isotopic signature would indicate a highly evolved diagenetic history.

When plotted on a  $\delta^{13}C$  versus  $\delta^{18}O_c$  diagram, most ancient phosphorites fall on a broad array constrained at one end by seawater and at the other by secondary apatite (isotopically light with respect to both oxygen and carbon) compositions (Fig. 6), indicating that the former display varying amounts of alteration. Nonetheless, the broad range of  $\delta^{13}C$  values for young phosphorites (+0.2 to -9‰) is similar to that of ancient phosphorites and no clear time-related trends are evident (Shemesh et al. 1988). This suggests that under favourable circumstances  $\delta^{13}C$  values may be relatively well-preserved over geologic time, or if alteration occurs its end members lie within the range of original  $\delta^{13}C$  values of francolites.

Studies are not limited to  $\delta^{13}C$  of the apatite  $CO_2$ , important supporting data may be obtained from the  $\delta^{13}C$  of associated carbonate minerals. It is known that secular variation in marine carbonate  $\delta^{13}C$  has occurred through the Phanerozoic (e.g. Tucker & Wright 1990; Veizer 1992) with long-term fluctuations of -1 to +2‰ and short-term 'excursions' of between -2 and +5‰. Smaller-scale geographical and depth-related variation may also occur. Clearly, appropriate corrections need to be applied before interpreting diagenetic conditions during francolite or carbonate precipitation. Once this is accomplished, the isotopic signatures of carbonate cements and their relationship to francolite in a paragenetic sequence can provide valuable information about the conditions of phosphogenesis.

Francolite oxygen and carbon isotope ratios may be indicators of depositional or diagenetic palaeotemperatures and/or fluid compositions, so it is essential that oxygen and carbon isotopes constitute only part of a multi-element, and ideally multi-phase, isotopic and geochemical study. In such cases, it may be possible to correct for alteration effects (e.g. McArthur et al. 1987; McArthur & Herczeg 1990) and obtain initial compositions which indicate conditions at the time of formation.

### Sulphur stable-isotopes

The presence of  $SO_4^{2-}$  in the francolite lattice as an isomorphous substitution for  $PO_4^{3-}$  is now generally accepted. Sulphate levels in unaltered phosphorites are close to 2.7% (McArthur 1978a, 1980, 1985). The degree of substitution depends on the  $SO_4^{2-}$  content of the aqueous phase in equilibrium with francolite (Smirnov et al. 1962; Bliskovskiy et al. 1977; McArthur 1985). The relative lack of variation displayed by unweathered francolites must reflect precipitation under similar, largely non-sulphidic, conditions, since removal of  $SO_4^{2-}$  from porewaters during sulphate reduction (Fig. 2, 7) should be reflected by declining  $SO_4^{2-}$  contents in francolites. However, it should be noted that the high concentration of sulphate in seawater, coupled with the ability of diffusion to replace  $SO_4^{2-}$  lost by the precipitation of sulphides, mean that the system is insensitive to small amounts of reduction.

Substitution of  $SO_4^{2-}$  for  $PO_4^{3-}$  in the francolite lattice offers the possibility of using sulphur stable-isotopes ( $^{34}S/^{32}S$  ratios expressed as ‰  $\delta^{34}S$  CDT – Cañon Diablo Troilite)




Fig. 8. Diagram showing the negative relationship between  $\delta^{13}\text{C}$  and  $\text{CO}_2$  contents displayed by francolites in modern phosphorite grains and nodules from the Peruvian margin. Grains are bulk samples; nodules are from a single locality (Station 4, 12° S);  $\text{CO}_2$  contents calculated using the XRD method of Gulbrandsen (1970). Note the different slopes and relative positions of the two regression lines. High  $\text{CO}_2$  and low  $\delta^{13}\text{C}$  values in francolites indicate high alkalinities in associated porewaters, which ultimately limit phosphate precipitation (modified from Glenn et al. 1988, Glenn 1990a).

to constrain the origin of phosphorites (Bliskovskiy et al. 1977; Nathan & Nielsen 1980; Benmore et al. 1983; McArthur et al. 1986; Piper & Kolodny 1987; Compton et al. 1993). The residence time of  $\text{SO}_4^{2-}$  in the oceans is very long, in the order of 12 Myr, several orders of magnitude longer than oceanic mixing times of 1.5 kyr and the oceanic residence time of phosphorus (Ruttenberg 1993) of 16–38 kyr, so the isotopic composition of modern seawater sulphate is globally uniform, with values of around +23  $\delta^{34}\text{S}$ . Nonetheless, on a geological timescale it has varied between +10 and +30‰  $\delta^{34}\text{S}$  (Claypool et al. 1980), in response to changing proportions of oxidised and reduced sulphur being buried as evaporites or sulphides and organic compounds, respectively.

Sulphur retains a seawater isotope composition in the oxic and suboxic zones (Fig. 7), but sulphate reduction leads to strong fractionation, commonly between 25 and 50‰, with the reduced phase being enriched in  $^{32}\text{S}$  and residual sulphate becoming increasingly isotopically heavy. However, the degree of fractionation is inversely related to the rate of sulphate reduction (Kaplan 1983), so for sediments which are very rich in easily metabolisable organic matter the fractionation may approach zero or even reverse. In contrast, by analogy to evaporite sulphate precipitation, little fractionation (only about +1.5‰; Claypool et al. 1980) should occur during the incorporation of  $\text{SO}_4^{2-}$  by francolite. Consequently, francolite precipitating from porewaters will record the  $\delta^{34}\text{S}$  sulphate values at the time of precipitation, and after correction for seawater composition ( $\Delta^{34}\text{S} = \delta^{34}\text{S}_p - \delta^{34}\text{S}_w$ ) should differentiate between oxic and anoxic environments. Precipitation of francolite during sulphate reduction has been suggested (Nathan & Nielsen 1980; Piper & Kolodny 1987) for Phosphoria Formation phosphorites, in which a strong negative correlation between  $\text{SO}_4^{2-}$  and  $\delta^{34}\text{S}$  values has been observed, supporting dominantly Rayleigh distillation-type reduction of sulphate to sulphide in a closed porewater system. It is unlikely in this case that low  $\text{SO}_4^{2-}$  values are due to weathering, because they are associated with a shift towards isotopically heavy  $\delta^{34}\text{S}$ , rather than the light values which are typical of alteration by meteoric waters.

From the above arguments, one would generally expect a positive correlation between isotopically light  $\delta^{13}\text{C}$  and positive  $\Delta^{34}\text{S}$  values in francolites. Late Tertiary – Quaternary seafloor phosphorites from the Blake Plateau, Chatham Rise, and the South African, Californian, East Australian and Peruvian margins generally have both  $\delta^{13}\text{C}$  and  $\Delta^{34}\text{S}$  within 2‰ of seawater values (Fig. 9). However, late Pleistocene – Holocene Namibian concretions, Tertiary Sechura Desert and Ica Plateau deposits in Peru, Oligo-Miocene phosphorites from southern Florida, and Upper Cretaceous – Palaeogene Moroccan phosphate grains yield  $\delta^{13}\text{C}$  of close to -7‰ and  $\Delta^{34}\text{S}$  up to +10‰. The latter group, therefore, confirm that the  $\delta^{13}\text{C}$  of “anoxic” francolites is apparently relatively uniform, but  $\Delta^{34}\text{S}$  varies considerably between high positive and, in one case, low negative values (Fig. 9; Nathan & Nielsen 1980; McArthur et al. 1986; Piper & Kolodny 1987).

Large variation in  $\Delta^{34}\text{S}$  is not unexpected given the complexity of sulphur isotope systematics, and likely differences in the extent and rate of sulphate reduction and openness of diagenetic systems. Additionally, negative  $\Delta^{34}\text{S}$  values might in some cases be caused by  $^{34}\text{S}$ -depleted  $\text{H}_2\text{S}$  diffusing upwards from an Fe-poor sulphate reducing environment being oxidised to sulphate at the anoxic/ suboxic interface. Similarly, upwards diffusion of  $\text{CO}_2$  produced by high rates of organic degradation may promote lower  $\delta^{13}\text{C}$  values of dissolved inorganic carbon in overlying zones. Francolites precipitated in such condition might be expected to exhibit a light sulphur and carbon signature. A Namibian




Fig. 9. Variation of  $\delta^{34}\text{S}$  and  $\delta^{13}\text{C}$  in modern and ancient francolites. Sulphur isotopes are expressed as  $\Delta^{34}\text{S}$ , calculated as  $\delta^{34}\text{S}(\text{francolite}) - \delta^{34}\text{S}$  (coeval evaporite), to compensate for secular variation in seawater compositions (modified from McArthur et al. 1986).

concretion studied by McArthur et al. (1986) provides one possible example of precipitation under these conditions, although the negative  $\Delta^{34}\text{S}$  value might equally be a result of a very high rate of sulphate reduction caused by the abnormally high organic carbon content (up to 26%) of the host sediments.

Sulphur stable-isotopes, therefore, yield valuable information which together with co-existing oxygen and carbon data are able to provide strong constraints on phosphogenic environments. It is interesting to note that results obtained to date indicate that there is no clear relationship between phosphorite type (i.e. nodular versus granular) and diagenetic environment. In both facies, francolite cements precipitate under oxic to suboxic or, less commonly, anoxic conditions. Even in anoxic environments, however, the relatively constant composition of francolites suggests that major ions such as Ca, Mg and Sr are able to diffuse freely from seawater. The possibility that such ions are sourced from dissolving carbonate appears to be inconsistent with Sr-isotope data (see below), which indicate precipitation in equilibrium with contemporary ocean water rather than inheritance of carbonate-derived Sr. Furthermore, many 'anoxic' francolites such as those from Namibia and the Sechura are not associated with carbonate. It seems likely, therefore, that francolite precipitation occurs almost entirely in the uppermost few decimetres of the sediment column.

#### Strontium isotopes

The residence time of Sr in seawater is about 3 Myr (Richter & Turekian 1993), also several orders of magnitude longer than oceanic mixing times, so the oceans are well-mixed

with respect to Sr. The  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio of modern biogenic marine carbonates and seafloor cements is identical to that of seawater at  $0.709175 \pm 0.000004$  (McArthur et al. 1993a), but the ratio in marine carbonates has varied substantially through time, from values of  $<0.707$  during the Late Permian and Late Jurassic to  $>0.709$  during the Early Cambrian and Quaternary (Peterman et al. 1970; Burke et al. 1982; DePaulo & Ingram 1985; Koepnick et al. 1985; Palmer & Elderfield 1985; Hess et al. 1986; Capo & DePaulo 1990; Kaufman et al. 1993; McArthur et al. 1993a, b; Denison et al. 1994b). Recent reviews include those of Veizer (1989), Stille et al. (1992), McArthur (1994) and Smalley et al. (1994).

The  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio of seawater is controlled largely by two major fluxes: (1) riverine and groundwater input, currently around 0.7119; (2) hydrothermal fluids, resulting from the interaction of seawater with young oceanic crust, with compositions of 0.7030. A smaller diagenetic flux from ocean floor sediments has a ratio of 0.7084, and consequently has far less impact on the overall seawater ratio. Secular variation in strontium isotopes is caused, therefore, by factors such as changes in global climate and eustatic sea-level, mountain-building events, and variation in rates of seafloor spreading. Many authors have argued that high erosion rates associated with major phases of tectonic uplift provide the main forcing mechanism on the Phanerozoic Sr-isotope curve, other fluxes producing only second order variation.

The Sr-isotope curve is particularly well-documented for the Cenozoic. During the last 40 Myr,  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios have become systematically heavier from around 0.7078 during the Eocene to 0.7092 at the present day so, at least within this interval, by measuring the ratio of an unknown sample and referring back to a reference curve (Fig. 10), it is sometimes possible to date the sample to with an accuracy of a few hundred thousand years. The precision of such dates is inversely proportional to the slope of the curve and the curves of different authors also diverge at certain intervals, probably as a result of slight differences in stratigraphic assignments, so data tend to become less precise with increasing age. Nonetheless, strontium isotopes provide a means of dating marine sediments which is far beyond the range of more established methods such as U-series disequilibrium and  $^{14}\text{C}$  methods.

The high Sr content of francolite (typically around  $2700 \mu\text{g g}^{-1}$  when formed) makes the mineral well-suited to Sr-isotope studies. Strontium is incorporated into francolite during precipitation, so in the case of phosphatised limestones strontium-isotopes potentially provides a means of assessing the date of mineralisation as opposed to the age of the host limestone, which may be obtained independently via  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios in calcite (Fig. 10), or palaeontological data. Dating of phosphatised hardground surfaces or individual phosphorite clasts in polymictic conglomerates, such as those commonly recovered off the margins of South Africa, Morocco and Spain, should also be feasible.

The use of  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios to date phosphogenesis necessitates a number of assumptions: (1) early diagenetic porewaters retain seawater Sr-isotope ratios; (2) francolite does not inherit Sr from precursor carbonates; (3) ratios remain unaltered during burial diagenesis. These assumptions seem to be generally valid (McArthur et al. 1990; Compton et al. 1990; Hein et al. 1993; Stille et al. 1994). The  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios of modern francolites are within error of modern marine Sr, while older seafloor phosphorites (Fig. 10) yield independent calcite and francolite dates which are consistent with other geological data. For example, it appears (McArthur et al. 1990; McArthur pers. comm. 1994) that on the Chatham Rise off New Zealand, replacement of Oligocene carbonates (29.8–28.1 Ma)




Fig. 10.  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio plotted against age and corresponding dates derived for coexisting francolites (above) and calcites (below) from phosphorites. The stippled area encloses the 95% confidence limit for the Neogene isotope curve of Hodell et al. (1991). Sample data are from McArthur et al. (1990), recalculated for a modern seawater  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio of 0.709175; dates (in part after McArthur pers. comm. 1994) are derived from the regression equations of Hodell et al. (1991). Dating errors were calculated by adding errors on measured  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios to the spread of  $^{87}\text{Sr}/^{86}\text{Sr}$  data defining the isotope curve. Large errors for samples between 4.5–2.5 Ma are caused by the flat nature of the curve over that interval. Note that phosphatisation may post-date limestone deposition by many Myr.

occurred in the Early Pliocene at about 4.8 Ma, Namibian shelf peloidal phosphorites formed in the mid-Pliocene around 3.4 Ma and in the Pleistocene at 0.4 Ma, and the Sechura Desert deposits of Peru are Late Miocene, accumulating around 8.9 Ma. Calcite ages derived from limestones collected off Cape Province South Africa (Fig. 10) range widely from Miocene to Pliocene (18.5–3.4 Ma) but phosphatisation appears to have occurred very recently, during the Pleistocene to Holocene. Errors on these dates are

generally in the order of 0.5 Myr, providing a better resolution than any other suitable dating method currently available.

Similarly, Compton et al. (1993) dated Hawthorn Group phosphorites in southern Florida as early Late Miocene ( $9.2 \pm 1.4$  Ma) to Late Oligocene ( $25.6 \pm 0.7$  Ma), with a major phosphogenic episode occurring during the Early Miocene between 21.9 and 20.4 Ma. Strontium-derived ages for benthonic foraminifera in sediments surrounding nodules were consistently younger than those obtained from nodule francolites, supporting evidence of extensive reworking. Hein et al. (1993) recognised two major phosphogenic events in the equatorial Pacific at 39–34 Ma (Late Eocene – Early Oligocene) and 27–21 Ma (Late Oligocene – Early Miocene) plus two minor episodes based on Sr-isotope data from seamount phosphorites, although the validity of these particular data has been questioned (McMurtry et al. 1994).

Most recently, Stille et al. (1994) described the Sr-isotope stratigraphy of phosphorites on the North Carolina continental shelf of the US, obtaining Early Miocene ages of 19.3–17.8 Ma. These authors demonstrated that although peloids and brachiopod fragments gave identical ages which are totally consistent with biostratigraphic data, older dates were obtained from coexisting intraclasts (confirming reworking), while vertebrate material yielded younger ages which are consistent with diagenetic modification. Clearly, careful selection of material is essential to obtain a reliable high-resolution isotope stratigraphy from granular phosphorites. Pleistocene and Holocene sediments in the same area yielded peloids with slightly younger Miocene ages ( $17.24 \pm 0.74$  Ma), indicating that these grains were derived from a now-eroded part of the phosphorite sequence.

The Sr isotope compositions of ancient apatitic fossils such as conodonts, ichthyoliths, fish teeth and inarticulate brachiopods (Shaw & Wasserburg 1985; Staudigel et al. 1985; Grandjean et al. 1987; Schmitz et al. 1991; Ingram et al. 1994; Stille et al. 1994) are, for the least altered material, in general agreement with the  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio in contemporaneous seawater, although there is a bias towards anomalously high ratios. Trends towards heavier Sr-isotope ratios associated with Sr-loss from francolites during weathering was also observed by McArthur et al. (1987) in their study of meteoric alteration of Lower Pliocene phosphorites from Cape Province, South Africa, so post-depositional exchange can certainly occur.

Clearly, careful screening of samples using a combination of petrographic, trace-element and stable-isotope geochemical techniques (Denison et al. 1994a; McArthur 1994; Smalley et al. 1994) is essential before applying Sr isotopes as a stratigraphic tool. One further note of caution is that many authors have used strong mineral acids to dissolve their samples. Although in most cases relatively pure material was selected, minor contamination of  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios from the siliciclastic fraction remains possible, which is liable to induce slightly elevated values (McArthur et al. 1993b), and therefore, in the Tertiary, younger ages. The use of a less aggressive digestion procedure (e.g. McArthur et al. 1990) would clearly be advantageous.

It may be concluded that Sr isotopes provide an excellent method of dating phosphogenesis. Three reservations have still to be made: (1) the Cenozoic  $^{87}\text{Sr}/^{86}\text{Sr}$  seawater-curve is now fairly well-established, but for older sediments determinations are more problematic; even in the Tertiary, some intervals may be dated much less precisely than others due to the low gradient of the curve or to poorer biostratigraphic resolution; (2) multi-generation deposits in which reworking has redistributed phosphorite from several

sources and ages into a single bed are common, so it is important to select the same phosphate grain-type and grain generation to obtain reproducible and reliable ages; (3) different ages than the time of phosphogenesis may be obtained due to alteration or the use of inappropriate dissolution techniques.

### Neodymium isotopes

Unlike Sr, the residence time of Nd in the oceans is thought to be in the order of hundreds of years to perhaps 2 kyr (Piepgras et al. 1979; Piepgras & Wasserburg 1980; Jeandel 1993), shorter than or comparable to the ocean mixing time of around 1.5 kyr. Consequently, the distribution of Nd is heterogeneous, with each major ocean basin having a relatively limited and characteristic range of  $^{143}\text{Nd}/^{144}\text{Nd}$  ratios.

The  $^{143}\text{Nd}/^{144}\text{Nd}$  ratio of a sample is a product of the radioactive decay of  $^{147}\text{Sm}$  to  $^{143}\text{Nd}$  at a constant rate ( $\lambda_{\text{Sm}}$ ) of  $6.54 \times 10^{-12} \text{ yr}^{-1}$ , and so reflects both the age and initial  $^{147}\text{Sm}/^{144}\text{Nd}$  ratio. On a global scale,  $^{143}\text{Nd}/^{144}\text{Nd}$  ratios have been increasing steadily since the formation of the Earth at 4.6 Ga. Comparisons between  $^{143}\text{Nd}/^{144}\text{Nd}$  ratios of different samples is strictly only valid, therefore, if all are of the same age. To compensate for these changes, Nd isotopes are generally reported as epsilon ( $\epsilon_{\text{Nd}}$ ) values (De-Paulo & Wasserburg 1976), which incorporate a time factor:

$$\epsilon_{\text{Nd}}(T) = [[(^{143}\text{Nd}/^{144}\text{Nd})_{\text{rock}, T} / (^{143}\text{Nd}/^{144}\text{Nd})_{\text{CHUR}, T}] - 1] \times 10^4 \quad (4)$$

where:  $(^{143}\text{Nd}/^{144}\text{Nd})_{\text{rock}, T}$  is the ratio in the sample at time  $T$ , and CHUR is a chondritic uniform reservoir (i.e. the bulk earth).

The composition of (CHUR,  $T$ ) may be calculated (see Rollinson 1993 for a recent detailed exposition):

$$(^{143}\text{Nd}/^{144}\text{Nd})_{\text{CHUR}, T} = (^{143}\text{Nd}/^{144}\text{Nd})_{\text{CHUR, today}} - [(^{147}\text{Sm}/^{144}\text{Nd})_{\text{CHUR, today}} \cdot (\exp(\lambda_{\text{Sm}} T) - 1)] \quad (5)$$

Commonly used values for CHUR, *today* are:  $^{143}\text{Nd}/^{144}\text{Nd} = 0.512638$ ;  $^{147}\text{Sm}/^{144}\text{Nd} = 0.1967$  (Jacobsen & Wasserburg 1980). The initial ratio of the sample (rock,  $T$ ) may be calculated using the same general equation, but substituting the  $^{143}\text{Nd}/^{144}\text{Nd}$  and  $^{147}\text{Sm}/^{144}\text{Nd}$  ratios measured in the rock for those of CHUR.

Neodymium isotope ratios have been measured in phosphorites by Shaw & Wasserburg (1985), McArthur et al. (1987), Keto & Jacobsen (1988) and Stille et al. (1994), and in biogenic phosphate by a number of workers (Wright et al. 1984; Shaw & Wasserburg 1985; Staudigel et al. 1985; Grandjean et al. 1987; Keto & Jacobsen 1987, 1988). Unfortunately, some of the early work (including that by Shaw & Wasserburg 1985) was undertaken on weathered samples (McArthur pers. comm. 1994) making much of the data unreliable. The  $\epsilon_{\text{Nd}}$  values of modern phosphorites and biogenic apatites appear to reflect a time-averaged equivalent of those in seawater, which in the Atlantic (Stille et al. 1992; Bertram & Elderfield 1993) has low  $^{143}\text{Nd}/^{144}\text{Nd}$  ratios ( $\epsilon_{\text{Nd}}$  of  $-12 \pm 2$ ), while Pacific waters are higher, with  $\epsilon$  values of  $-3 \pm 2$ , and the Indian Ocean lies in between, around  $-8 \pm 2$ . These differences reflect the dominant sources of REEs in each ocean basin, from old continental crust in the Atlantic, compared with young crustal and active hydrothermal systems in the Pacific. Smaller differences occur within individual basins (e.g.

Jeandel 1993) and may be used to trace the origins, distribution and evolution of different water masses.

Geochemically distinct water masses seem to have existed in the Atlantic and Pacific for about the last 50 Myr (Fig. 11), prior to which a convergence of  $\epsilon_{\text{Nd}}$  values to between  $-10$  and  $-5$  indicates the existence of a single uniform ocean (Panthalassa; Keto & Jacobsen 1987, 1988). Earlier in the geological record, however, there is strong evidence of the existence of separate geochemical provinces. During the Cambrian and Ordovician (Keto & Jacobsen op. cit.), so-called "North Iapetus" rocks from North America have low  $^{143}\text{Nd}/^{144}\text{Nd}$  ratios and  $\epsilon_{\text{Nd}}$  values of  $-20$  to  $-10$ , while "South Iapetus" material from Europe and the south-eastern US exhibits values around  $-5$  to  $-9$ . Only with the onset of the Taconic Orogeny in eastern North America at the end of the Ordovician, did this geochemical separation of water masses in the Iapetus Ocean disappear.

On a smaller scale, an isolated basin in the South Atlantic during the Upper Cretaceous – Palaeocene (Grandjean et al. 1987) is indicated by low  $\epsilon_{\text{Nd}}$  values of around  $-15$  in biogenic phosphates (Fig. 11) from the rocks of West Africa (Togo, Guinea, Angola). In some cases samples have  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios of contemporaneous seawater, but  $\epsilon_{\text{Nd}}$  values of an isolated basin, indicating good connection with the world ocean but limited short-term mixing. Isolation appears to have continued until the mid-Eocene when a modern oceanic circulation pattern was established. More recently, Stille et al. (1994) demonstrated that francolite  $^{143}\text{Nd}/^{144}\text{Nd}$  ratios are constant in their Miocene phosphorite (peloid) samples and are thought to be representative of western mid-Atlantic (Gulf




Fig. 11.  $\epsilon_{\text{Nd}}$  versus age for Phanerozoic phosphatic fossils. Discrete Atlantic and Pacific provinces are identifiable back to 50 Ma, but data indicate the dominance of a single well-mixed oceanic reservoir (Panthalassa) between 50–450 Ma. A distinct West African Basin water mass existed during the U. Cretaceous – Palaeocene. Prior to 450 Ma, discrete South and North Iapetus water masses may be distinguished (modified from Kolodny & Luz 1992; data from: Shaw & Wasserburg 1985; Staudigel et al. 1985; Grandjean et al. 1987, 1988; Keto & Jacobsen 1987).

Stream) seawater at 19.3–17.8 Ma. However, marginally younger Miocene peloids reworked into Holocene sediments have substantially lower Nd-isotope ratios, suggesting that a major change occurred in oceanic current conditions along the Carolina continental margin around 17.2 Ma.

It is noteworthy that even though there is clear evidence that the REEs, including Nd, are lost from francolite during weathering, relatively intense weathering does not necessarily affect  $\epsilon_{\text{Nd}}$  values (McArthur et al. 1987; Stille et al. 1994). Clearly then, Nd isotopes in phosphorites and biogenic phosphates appear to offer a relatively robust and very powerful means of studying water-mass distributions and palaeoceanographic circulation, and may be used to estimate the average age of exposed continental crust bordering ancient ocean basins.

### Uranium isotopes

The basis of natural uranium-series dating of phosphorites is the assumption that U is incorporated during phosphogenesis and that  $^{230}\text{Th}$  and  $^{231}\text{Pa}$ , the radioactive daughters of U, develop in the sample as a function of time according to their half-lives. The residence time of U in the oceans is around 330 kyr (Richter & Turekian 1993), so the modern ocean are isotopically uniform with a  $^{234}\text{U}/^{238}\text{U}$  activity ratio of  $1.144 \pm 0.004$ . The use of radiogenic isotopes for dating requires that: (1) U, but little or no Th or Pa, was incorporated into the phosphorite during or very soon after its formation; (2) U is of marine origin with a  $^{234}\text{U}/^{238}\text{U}$  activity ratio close to seawater; (3) the phosphorite has remained closed with respect to U, Th and Pa isotopes. Extensive work over the last 25 years has shown that these assumptions appear to be broadly valid.

Uranium-series dating was fundamental in demonstrating the very limited nature of phosphogenesis in the modern oceans. Prior to the 1970s, it had been assumed (McKelvey et al. 1953; Sheldon 1964; McKelvey, 1967; see Cook et al. 1990 for a historical review) that the widespread occurrence of seafloor phosphorites along the western margins of most continents, provided definitive proof of the link between upwelling and phosphogenesis, as proposed many years earlier by Kazakov (1937). Uranium-series studies of phosphorites from off California, the Chatham Rise and Agulhas Bank by Kolodny (1969) and Kolodny & Kaplan (1970a), however, showed that none of these had an excess of  $^{234}\text{U}$  and must be older than 800 ka. It was only later, that studies of deposits off the Peru-Chile and Namibian margins (Baturin et al. 1972, 1974; Veeh et al. 1973, 1974; Burnett & Veeh 1977; Burnett et al. 1980) proved that these contained a  $^{234}\text{U}/^{238}\text{U}$  activity ratio very close to present seawater and must be forming at the present day. Subsequently, other modern phosphorites have been identified off western South Africa (Birch et al. 1983; McArthur et al. 1988) and East Australia (Kress & Veeh, 1980; O'Brien & Veeh 1980; O'Brien et al. 1986, 1990).

Dates obtained from  $^{234}\text{U}/^{238}\text{U}$  activity ratios may be confirmed by comparison with other components of the decay series. The use of “concordia” diagrams for  $^{234}\text{U}/^{238}\text{U}$  (Fig. 12) and  $^{231}\text{Pa}/^{235}\text{U}$  (Fig. 13) versus  $^{230}\text{Th}/^{234}\text{U}$ , may be used to test the coherence of the decay series data. Such work has confirmed that the initial  $^{234}\text{U}/^{238}\text{U}$  activity ratio in phosphorites is slightly higher than that in modern seawater, possibly due to the addition of small amounts of  $^{234}\text{U}$  to porewaters from the host sediment (Veeh & Burnett 1982). Alternatively, higher ratios might be caused by secular variation in the isotopic composi-




Fig. 12. Ideal development in time or "concordia" (heavy curved line) of  $^{234}\text{U}/^{238}\text{U}$  activity ratios versus  $^{230}\text{Th}/^{234}\text{U}$ , with data plotted for Peruvian margin nodules and grains. Note the occurrence of initial  $^{234}\text{U}/^{238}\text{U}$  activity ratios of  $>1.14$ , and the occurrence of old nodules with  $^{234}\text{U}/^{238}\text{U}$  ratios of  $<1.0$ . The dashed curved lines enclose a deviation of  $2\sigma$  from the concordia path (data from: Burnett et al. 1988; Burnett & Veeh 1977; Veeh & Burnett 1982).

tion of seawater (Richter & Turekian 1993), although there is no clear evidence that such variation has occurred during the last 200 kyr (Henderson et al. 1993).

The occurrence of old nodules with  $^{234}\text{U}/^{238}\text{U}$  activity ratios of  $<1$  and anomalously low  $^{234}\text{U}/^{238}\text{U}$  versus  $^{230}\text{Th}/^{234}\text{U}$  values (Fig. 12) indicate that decay occurs in a semi-closed system in which a small fraction of  $^{234}\text{U}$  is lost to porewaters. Limited isotopic exchange between phosphorites and porewaters in samples from the Peru-Chile Shelf is confirmed by  $^{230}\text{Th}$  and  $^{231}\text{Pa}$  data (Fig. 13; Veeh 1982; Burnett et al. 1988), but does not invalidate the dating method; samples from the South African (Roe et al. 1982; McArthur et al. 1988) and East Australian (Fig. 13; O'Brien et al. 1986) margins show no evidence of exchange.

Detailed geochronologies of Peru-Chile (Burnett & Veeh 1977) and western South African (McArthur et al. 1988) phosphorite nodules suggest that phosphogenesis is episodic, occurring preferentially during periods of high eustatic sealevel. On the East Australian margin, on the other hand, phosphogenesis appears to have been more or less continuous over the last 180 kyr (Kress & Veeh 1980; O'Brien & Veeh 1980; O'Brien et al. 1986). These differences must relate to the contrasting sedimentological and oceanographic settings of the two areas. It is interesting to note that phosphorite deposition also appears to be episodic on much longer time scales (Cook & McElhinny 1979; Cook 1984; Glenn et al. 1994b).




Fig. 13. Concordia diagram of the activity ratios of  $^{231}\text{Pa}/^{235}\text{U}$  versus  $^{230}\text{Th}/^{234}\text{U}$ , with data plotted for Peruvian margin nodules and grains, and East Australian margin nodules. Note the concordant ages of nodules from Australia, but the “excess Pa” in the Peruvian nodules and “excess Th” in Peruvian grains (data from: Veeh 1982; Burnett et al. 1982, 1988).

Detailed uranium-series studies of individual grains, nodules and crusts may be used to elucidate the modes and rates of growth of phosphorites. The  $^{230}\text{Th}/^{234}\text{U}$  age profiles of subsamples taken across nodules and crusts (Burnett et al. 1982, 1983; Kim & Burnett 1986) indicate that nodules located at the sediment/water interface grow *downwards* into the soft underlying mud, while buried crusts grow *upwards* within the sediment. This is not surprising given the subsurface nature of the porewater phosphate maximum (Fig. 1, 2). Measured nodule growth rates vary from  $<0.1\text{--}1\text{ cm kyr}^{-1}$ , considerably slower than the surrounding sediments which are accumulating at a rate of at least  $2\text{--}10\text{ cm kyr}^{-1}$ . The retention of nodules at the sediment/water interface, therefore, must be explained by the buoying action of the burrowing infauna, or perhaps by periods of non-deposition or erosion. Most recently, Burnett et al. (1988) employed a combination of high-precision U-disequilibrium dating with accelerator mass spectrometry (AMS)  $^{14}\text{C}$  dating of francolite $\text{CO}_2$  to show that Peru margin phosphorite grains grow more rapidly than nodules, probably taking  $<10$  years to become fully developed.

Kolodny & Kaplan (1970a) and later workers (Burnett & Veeh 1977; O'Brien & Veeh 1980; Veeh & Burnett 1982) demonstrated that the  $^{234}\text{U}/^{238}\text{U}$  ratio in phosphorites is affected by U oxidation states, with higher ratios in U(VI) than U(IV). It is suggested that these differences arise during the oxidation of U which causes a small leakage of  $^{234}\text{U}$  from U(IV) to U(VI), via the decay of both  $^{238}\text{U}(\text{IV})$  and  $^{238}\text{U}(\text{VI})$  to  $^{234}\text{U}(\text{VI})$ . Once in hexavalent state, a small fraction of these  $^{234}\text{U}$  atoms may form highly soluble carbonate complexes and diffuse from the phosphorite nodules, resulting in bulk  $^{234}\text{U}/^{238}\text{U}$  ratios which are less than 1.00 in very old samples. Leakage of  $^{234}\text{U}$  will introduce a small error into age calculations towards older values. Thus, the outer weathered rims of seafloor phosphorites have lower U(IV) and  $^{234}\text{U}/^{238}\text{U}(\text{IV})$  ratios than their unweathered cores (Kolodny & Kaplan 1970a, Baturin & Kochenov 1974).

Using the above principles, O'Brien et al. (1987) demonstrated that East Australian nodules underwent two stages: (1) incorporation of U(IV) under dominantly suboxic conditions in the sediment; (2) oxidation of U to U(VI) during exposure and reworking of the phosphorite on the seafloor. Differences between U(IV) contents and  $^{234}\text{U}/^{238}\text{U}$ (IV) ratios between buried and exposed nodules were related to the length of time nodules were exposed to oxidising conditions at the sediment/water interface. Uranium-series work, therefore, confirms petrographic, sedimentological and other geochemical studies which indicate the importance of reworking in the development of these (and most other) phosphorites.

### **Environmental issues**

On a world-wide basis, the overwhelming majority (>95%) of the 135 Mt of phosphate rock produced annually is used for fertiliser manufacture (Anon 1993). Consequently, long-term growth will ultimately be driven by human population growth, which is expected to remain at current rates for at least several more years (Brown et al. 1987).

In this section, we address some of the most serious problems which effect the environment during the exploitation of phosphate rock. These problems will include those associated with: (1) the mining process; (2) chemical treatments and procedures associated with conversion of phosphate rock to fertiliser; (3) release of toxic or radioactive elements to the environment as a result of mining or processing. In this review, we will emphasise those problems which relate directly to the chemical nature of phosphorites. Many of the current environmental issue centre on contamination by the impurities, especially radionuclides, contained in phosphogypsum, a by-product of fertiliser manufacture, so particular emphasis will be placed on this topic.

#### *Mining and processing*


Since most phosphate mining is of the "open-pit" variety, extremely large quantities of material are moved in the process. This type of mining significantly scars the landscape which may not be returned to its original contours following mining. Overburden, ore, ore concentrate, products and by-products are all moved in one form or the other, most often as a slurry which also consumes significant water resources. During the beneficiation process, the ore is concentrated by a variety of methods, usually including sizing and flotation techniques. This results in the production of extremely fine-grained phosphatic "waste clays", one of the most serious environmental problems of the mining/beneficiation part of the fertiliser cycle. This problem is minimised, of course, for deposits (e.g. the Upper Cretaceous – Eocene of the Tethyan province) where the clay content is volumetrically unimportant.

Waste clays are extremely fine grained, with tremendous surface area, so the settling characteristics are very poor. Without any treatment, these clays require decades to settle from about 5 to 20% solids. In Florida, the artificial ponds used for settling take up a substantial (40–60%) fraction of the mine area. Fortunately, progress has been made recently (McFarlin 1992) with rapid dewatering techniques involving the use of a flocculant and floc collector, which brings the solids content up to 25% within a few minutes. Hopefully, this type of technology will prevent the waste clay situation from getting worse in the fu-

ture, but the problem of the numerous clay-settling ponds which now exist must also be addressed.

### *The “phosphogypsum” problem*

Approximately 70% of phosphate ore mined is initially converted to phosphoric acid, an important intermediate product for the manufacture of soluble 'high analysis' phosphatic fertilisers. The commercial production of phosphoric acid may be achieved by thermal methods, involving the reduction of phosphate rock in an electric furnace, or more commonly, using the so-called "wet-process" where the rock is reacted with sulphuric acid. The combined capacity of existing phosphoric acid plants is so large that the phosphate fertiliser industry is the largest single consumer of sulphuric acid in the world. This process also results in the production of substantial quantities of by-product gypsum known as "phosphogypsum". A simplified chemical reaction may be shown as:



While the mole ratio between gypsum and phosphoric acid is 5 : 3, the mass ratio is about 3 : 1, *i.e.* about 3 tonnes of gypsum are produced for every tonne of phosphoric acid. As a consequence, the phosphate industry as it operates in most parts of the world could more accurately be termed a ‘phosphogypsum’ industry. Since the phosphate industry measures its phosphoric acid production in terms of many millions of tonnes per year, the amount of by-product gypsum produced is clearly substantial, on the order of 100 Mt yr<sup>-1</sup> on a world-wide basis.

During wet-process production,  $\text{CaSO}_4$  may be crystallised in either the dihydrate ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ) or hemihydrate ( $\text{CaSO}_4 \cdot \frac{1}{2}\text{H}_2\text{O}$ ) form, depending on the operating conditions. Phosphogypsum produced as a result of this reaction is filtered off and in many areas, including Florida, it is pumped as a slurry containing about 20% solids to a nearby storage area, the so-called “gyp-stacks”, where it is cooled and dewatered. The phosphoric acid produced is later reacted with ammonia or concentrated phosphate ore to produce ammonium phosphate or triple superphosphate, the two most common types of phosphate fertilisers.

Phosphogypsum has been used in Japan, Germany and some other countries for production of cement and other construction materials, but its commercial use in the US is currently limited to small quantities sold to farmers as a soil amendment. The US Environmental Protection Agency (EPA) has ruled that gypsum must be placed on stacks or mine cuts, and only limited quantities can be removed for agricultural or research purposes. Phosphogypsum cannot be used for construction or similar applications in the United States subsequent to the June 3, 1992 (57 Fed Reg 23305) EPA ruling. This ruling was predicated on: (1) risk assessments and environmental concerns associated with radionuclides in the  $^{238}\text{U}$  decay chain, especially radon (Rn); (2) the US already produces a substantial amount of natural gypsum by mining.

This situation is unfortunate for places such as Florida where the inventory of by-product gypsum has been growing tremendously within the past few decades. This problem does not exist in some countries, such as Morocco, where there are plans to pump

25 kt per day of phosphogypsum from one plant directly into the Atlantic Ocean (Becker 1989). In Florida, it is estimated that more than 600 Mt of phosphogypsum are currently stored at 17 sites throughout the state, with >30 Mt being added each year. At this pace, Florida will have close to 1 Gt of stockpiled phosphogypsum by the year 2000 (May & Sweeney 1982; Nifong 1988).

Besides the obvious waste of a potentially valuable by-product and the unsightly physical appearance of the stockpiles, the main problem associated with phosphogypsum storage is the potential effect on the surrounding environment, specifically the air and water resources in the vicinity of gypsum stacks (Kouloheris 1980). According to a review by Nifong (1988), the principal environmental concerns with phosphogypsum storage are fluoride emissions to the atmosphere and possible contamination of groundwater with radium, sulphate and fluoride, due to leaching by process water or rainwater. Other U-series nuclides associated with phosphogypsum, especially  $^{210}\text{Po}$ , have been found at high concentrations in shallow groundwater in the Florida mining area (Harada et al. 1989).

Studies which have considered the radiological aspects of the agricultural use of phosphogypsum have generally concluded that the increased risk relative to more traditional fertilisers is low, both from the increased radon flux from the soil and from uptake by plants (Lindeken & Coles 1978; Lindeken 1980; Roessler 1984, 1988; Mortvedt 1992). The most important pathway between phosphogypsum-derived radionuclides and humans, therefore, is probably through groundwater (Miller & Sutcliffe 1984). Whether the radioactive nuclides are an environmental hazard or not, may be a moot point in terms of use of phosphogypsum in the United States, since a recent EPA ruling has set a 10 pCi g<sup>-1</sup> limit of  $^{226}\text{Ra}$  allowable in phosphogypsum for agricultural purposes. At this stage, it is not at all clear whether radionuclides from phosphogypsum stacks are actually contaminating the surrounding groundwater. Although elevated levels of  $^{210}\text{Po}$  have been documented from areas in the vicinity of phosphogypsum stacks in Florida (Burnett et al. 1987; Harada et al. 1989), the source has not been determined. In some cases, it appears that the direct source may be from the natural phosphate rock of the area. In other cases, phosphogypsum appears to be the source.

Research currently in progress concerning the radiochemistry of phosphogypsum has shown that the radionuclides are apparently hosted in a separate phase from the calcium sulphate (Burnett et al. 1992). The radionuclide-bearing phase is insoluble in water and may be an aluminium phosphate with relatively high concentrations of Ba, Sr and REEs. This appears to be somewhat of a paradox. How can radionuclides be leached from phosphogypsum and enter the shallow aquifer when they are contained in a very insoluble phase? The answer may be in the action of bacteria. Cherrier et al. (1995) have shown that bacteria isolated from phosphogypsum seem to incorporate polonium and distribute it into their cellular components in a manner similar to sulphur. In a more recent set of experiments involving leaching of gypsum in columns, it was observed that Po was released when the gypsum was inoculated with sulphate-reducing bacteria (Burnett et al. 1993). These results suggest that sulphur bacteria may cycle Po inadvertently while they are processing sulphur. Both elements are in the same periodic group and have certain chemical similarities. Thus, the enzymatic action of bacteria could be the link which initiates mobilisation of radionuclides from phosphogypsum.

Is there a solution to this problem? Can phosphogypsum be economically purified or converted to a useful form without the associated impurities and put to some good use?

Natural gypsum is mined in several areas of the world and has a variety of uses, especially for construction activities. Unfortunately, phosphogypsum is less suitable for such purposes because of its high water content, fine particle size and generally inconsistent quality. At least two conversion processes are already known: (1) conversion and recovery of sulphur gases which may then be returned to the fertiliser cycle as sulphuric acid; (2) the Merseburg process or double decomposition method, where gypsum is reacted with ammonium carbonate and forms ammonium sulphate, an excellent fertiliser, and by-product calcium carbonate. Unfortunately, the present situation is such that neither process appears to be economically feasible. This is indeed unfortunate, as a potential resource is being wasted and at a great cost, not only in terms of harm to the environment, but in terms of actual expense. Schultz (1992) reported, for example, that if all the gypsum stacks in the United States had to be closed by covering with an impervious membrane and a soil cap to support vegetation, the cost would be in the neighbourhood of US\$ 1.3 billion or an average of about \$20 million per stack! Furthermore, this cost estimate does not include the on-going costs of environmental monitoring or maintenance. Clearly, it would be in the best interests of everyone involved to pursue options for an environmental-sound yet economically-attractive use of phosphogypsum.

The recent research on the radiochemistry of phosphogypsum mentioned above has shed some insights into this problem. The most important finding is that  $^{226}\text{Ra}$  and other radionuclides are not contained in the  $\text{CaSO}_4$  phase of the gypsum. Although the host has yet to be positively identified (an insoluble phosphate or sulphate appear to be the most likely), the observation that the radionuclide fraction resides in a discrete phase is significant. For example, this finding opens the promise of either physical or chemical separation of the radionuclide fraction from the gypsum component. But can this be done economically? Although the economic outlook does not appear favourable, the new findings provide additional incentive to search for a solution, especially one with a process that will also result in an economically attractive product.

#### *Release of toxic and radioactive elements*

Phosphorites may contain relatively high levels of certain elements that may be harmful to humans, either because of some inherent toxicity or because they are radioactive. Of the toxic elements which may be enriched in phosphorites, Cd, Se and As are probably the most important. It should be emphasised, however, that the concentration of these elements varies widely between and sometimes within deposits, so it is important to evaluate each case, on a deposit scale, rather than depend on average analyses. Cadmium in particular can, but does not necessarily (Anon 1989; Johnston & Jones 1992), accumulate in soils and plants through repeated fertiliser use. Japanese law, for example, stipulates a maximum concentration of  $1.5 \mu\text{g g}^{-1}$  Cd per 1%  $\text{P}_2\text{O}_5$  (Baechle & Wolstein 1984) in mineral fertilisers, while in Germany and some Scandinavian countries phosphorites containing  $>20 \mu\text{g g}^{-1}$  Cd are considered to be unsuitable for fertiliser production. Sales to Western Europe of phosphate rock from high-Cd deposits such as those of Senegal and Togo have suffered significantly in recent years because of environmental concerns.

Clearly, an understanding of the location and distribution of Cd and other toxic elements, is important in planning the development and beneficiation of a phosphorite deposit. It has been reported that Cd, Pb, Zn, Se, Hg and As may be reduced (volatilised)

considerably by high-temperature calcining of the ore. This is normally uneconomic except in cases where the content of organic matter is high (Isherwood 1992), and also leads to a marked decrease in the chemical reactivity of the phosphorite, reducing the efficiency of the phosphoric acid process. Furthermore, a transference of toxic metals from the fertiliser cycle to the atmosphere, and subsequently to the soil and groundwater in the vicinity of the processing plant, does not necessarily represent a significant improvement to the environmental problem.

During chemical processing, harmful gases may also be released. These include  $\text{SO}_2$ , associated with the huge quantities of sulphuric acid used by the industry, and fluoride emissions. The technology exists to limit these emissions to environmentally acceptable levels and thus the problem may be limited to ensuring that the regulatory restrictions are being followed. It should also be mentioned in this context that release of potentially harmful elements such as fluoride, may occur without any human intervention at all, i.e. "natural" contamination. This was documented in the case of an unmined area in Senegal where high fluoride levels in the groundwater, presumably leached out of the surrounding phosphate rock, has been implicated as the cause of health problems in the region (Travi 1988).

The one problem that is perceived as being the most important, at least in the United States, is that concerning the content of radioactive elements in phosphorites. Sedimentary phosphorites of most ages and areas (but with some notable exceptions) contain significant uranium concentrations, with typical values over  $100 \mu\text{g g}^{-1}$ . Since virtually all economic deposits of phosphorite are old enough for radioactive equilibrium to be established in the  $^{238}\text{U}$  decay chain (this requires approximately 1 Myr), the specific activity (number of radioactive disintegrations per unit time per unit mass) of all U-decay products will be approximately equal. Thus, a Miocene phosphorite from Florida with a U content of  $100 \mu\text{g g}^{-1}$  will not only have a  $^{238}\text{U}$  activity of approximately  $33 \text{ pCi g}^{-1}$ , but have individual activities of  $^{230}\text{Th}$ ,  $^{226}\text{Ra}$ ,  $^{222}\text{Rn}$  and other radionuclides of an equivalent amount as well. During mining, and especially during chemical processing of the ore, these radioelements can become separated and radioactive equilibrium is no longer established. This creates an entirely different situation, since each isolated daughter will behave according to its own chemistry and, if truly isolated, will decay away with its own half-life. For example, should  $^{210}\text{Po}$  be released into the environment without its radioactive predecessors, it will decay away with its own half-life (138.4 days), rather than being controlled by longer-lived parents.

The major concern in the United States involving the dispersal of radioactive elements associated with phosphate rock, is with the radioactivity of phosphogypsum. This concern, expressed in recent EPA rulings, is centred on the  $^{226}\text{Ra}$  content of phosphogypsum, the radioactive parent of gaseous  $^{222}\text{Rn}$ . In spite of the fact that emissions from gypsum stacks do not appear to be abnormally high, the concern over any possible contributor to the radon inventory is so great, that the US EPA has greatly restricted the use of phosphogypsum. As mentioned above, new research indicates that radium is apparently hosted in a discrete mineral phase in phosphogypsum, thus opening up the possibility that purification is possible. The problem, and the real challenge, lies in development of an economically feasible way to accomplish this task.

## Discussion and conclusions

As a broad generalisation, there is a correspondence between those trace elements which are enriched in seawater and/or marine organic matter (Tooms et al. 1969; Prévôt & Lucas 1979; Piper 1994), and those occurring at high levels in phosphorites. This relationship is not unexpected given the open-marine depositional setting of most deposits, and the inherent involvement of organic matter and bacterial processes in their formation. Levels of enrichments are controlled not only by the crystallographic properties of francolite and kinetic factors, but also by ambient geochemical conditions (Eh, pH, alkalinity) during precipitation, and may be modified substantially by late diagenetic, metamorphic and weathering effects. Moreover, phosphorites contain complex admixtures of other inorganic and organic constituents, and these may additionally host high proportions of trace elements. Consequently, there is commonly no simple correlation between the most enriched elements and P even within a single deposit, and different deposits will have different characteristics that relate to their unique geological histories.

Difficulties in apportioning trace-elements to particular phases are partly a consequence of sampling constraints. Phosphorites are by their very nature highly inhomogeneous at all scales, from the dissemination of phosphatic grains or nodules in a bed of siliciclastic and/or carbonate sediment, through the intimate association of  $\mu\text{m}$ -scale francolite crystallites with other authigenic and detrital particles and organic matter, down to geochemical variation on the surfaces of and within individual crystals. Most of our current knowledge concerning trace-element distributions in phosphorites originates from a relatively limited range of bulk-rock analyses. Only a few deposits, such as the Middle Cambrian Georgina Basin deposits of northern Australia (Cook 1972; Howard & Hough 1979), Permian Phosphoria Formation of the US (Gulbrandsen 1966, 1975; McKelvey et al. 1986; Piper & Medrano 1994), and Cretaceous-Eocene deposits of Morocco (Prévôt & Lucas 1985; Prévôt 1990) are well characterised. Additional high quality multi-element data sets for geologically and stratigraphically well-constrained sample sets are needed to complement these data, while the greater use of physical and chemical separation techniques would add significantly to our knowledge of trace-element partitioning in different sediment components.

Major advances require new approaches. The use of high resolution imaging techniques, including backscattered scanning electron microscopy, combined with sensitive microanalytical methods (see Potts et al. 1993a for a recent review) such as the ion microprobe, synchrotron X-ray fluorescence microprobe, laser ablation-inductively coupled plasma-mass spectrometer (LA-ICP-MS), particle-induced X-ray emission or LA-gas source-MS analysis, have enormous potential to determine inter-elemental and isotopic relationships on a single-crystal scale. Such work is in its infancy, but is already having an impact on the field (Grandjean & Albarède 1989; Jarvis & Jarvis 1991; Grandjean-Lécuyer et al. 1993). An ability to determine the geochemical composition of separate phases and to study compositional variation within individual grains and crystals is essential to improve our understanding of the processes which lead to the incorporation of trace elements in phosphorites. It should be stressed that this work must be integrated with established field, petrographic and geochemical techniques if interpretations of the information obtained are to be soundly based.

Experimental data concerning the kinetics and thermodynamics of trace-element incorporation by francolites is a prerequisite for quantitative modelling. Further experimental work, together with mathematical modelling of natural and artificial systems, are essential if the transition is to be made from empirical to theoretical studies of phosphogenesis. However, the complexities of natural phosphogenic environments should not be underestimated. It is also important to make the point that there remains a limit to which present-day phosphorites may be used as analogues for the genesis of ancient deposits. The most notable difference is that most major phosphorites are granular while modern phosphorites are dominantly nodular. Recent studies have demonstrated that granular phosphorites are forming in modern settings and that phosphorite hardgrounds are more widespread in both the modern and ancient than previously realised (Glenn et al. 1994a, b). Nonetheless, there remains no known exact analogue for the phosphorite 'giants' of the geological record. The basic physical-chemical processes which control the precipitation of francolite and the genesis of phosphorites must have been the same in the past as they are at the present day, and there is no doubt that we have learnt much about these from our studies of modern sediments. Nonetheless, there remains the fundamental questions as to why ancient deposits are so different and what are the environmental factors that these differences reflect?

Over the last two decades, geochemical and particularly isotopic studies have led to major advances in our understanding of the genesis and diagenesis of phosphorites, but many problems remain unsolved. What is the precursor mineral to francolite in modern environments? When and how are specific trace elements incorporated in phosphorites, where are they located and what was their source? Are humic compounds or other organic complexes instrumental in these processes? To what extent do kinetic and thermodynamic factors control trace-element contents? Are trace-elements good redox indicators? Are weathering and metamorphic effects kinetically controlled; is long term-burial at modest depths equivalent to short periods of deep burial?

To answer these questions, traditional geological analyses of ancient and modern phosphorites will have to be complemented by: (1) new analytical methods to empirically determine the distribution and behaviour of trace elements and isotopes in phosphorites; (2) better integration of inorganic and organic geochemical studies; (3) multi-element and multi-phase studies to examine the relationship between francolite geochemistry and those of associated mineral phases; (4) more extensive data on the porewater and solid-phase geochemistry of modern phosphogenic and associated non-phosphogenic environments; (5) experimental work to determine the kinetic and thermodynamic factors which control apatite precipitation and the incorporation of trace elements in francolite; (6) more refined modelling of natural and experimentally derived data.

Results from these geochemical studies will continue to provide one of the best means of refining models of phosphogenesis. An understanding of geochemical processes is essential to assessing and ameliorating the environmental impact of phosphate rock extraction and processing and the use of phosphate fertilisers.

### Acknowledgements

This paper is based partly on presentations made at the IGCP 325 International Symposium & Workshop, *Concepts and Controversies in Phosphogenesis*, held in Matten, Switzerland, 6–10 September 1993. The authors wish to thank Karl Föllmi (ETH Zurich), other members of the organising committee, and the sponsors, for their

enthusiasm and support during the meeting. Kingston University funded IJ during his attendance at the symposium. WB acknowledges the financial support of the Florida Institute of Phosphate Research (FIPR#90-05-035) for the phosphogypsum studies. The manuscript benefited greatly from critical reviews by Drs Peter Cook (British Geological Survey), Yoshua Kolodny (Hebrew University of Jerusalem), John McArthur (University College London) and Arthur Notholt (Mineral Resources Consultant, Uxbridge). This is a contribution to IGCP Project 325, Palaeogeography of Authigenic Minerals.

## REFERENCES

ABU MURRY, O.S. 1993: Distribution of rare earth elements in Jordanian phosphate. Concepts and Controversies in Phosphogenesis. Int. Symp. and Workshop, Matten, 6-10 September, 1993.

ALTSCHULER, Z.S. 1973: The weathering of phosphate deposits – geochemical and environmental aspects. In: Environmental Phosphates Handbook (Ed. by GRIFFITH, D., BEETON, A., SPENCER, J.M. & MITCHELL, D.T.). Wiley & Sons, New York, 33-96.

– 1980: The geochemistry of trace elements in marine phosphorites. Part I: Characteristic abundances and enrichment. In: Marine Phosphorites (Ed. by BENTOR, Y.K.). Soc. Econ. Paleont. Miner. Spec. Publ. 29, 19-30.

ALTSCHULER, Z.S., CISNEY, E.A. & BARLOW, I.H. 1952: X-ray evidence of the nature of carbonate apatite. Geol. Soc. Am. Bull. 63, 1230-1231.

ALTSCHULER, Z.S., CLARKE, R.S. & YOUNG, E.Y. 1958: Geochemistry of uranium in apatite and phosphorite. US Geol. Surv. Prof. Pap. 314-D, 45-90.

ALTSCHULER, Z.S., BERMAN, S. & CUTTITI, F. 1967: Rare earths in phosphorites – geochemistry and potential recovery. US Geol. Surv. Prof. Paper 575(B), B1-B9.

ANON 1989: Cadmium in phosphates: one part of a wider environmental problem. Phosphorus Potassium 162, 23-30.

– 1993: Phosphate Rock helps feed the world. Engineer. Min. Jour. 194, 2, 73.

ATLAS, E.L. 1975: Phosphate equilibria in seawater and interstitial waters. PhD Dissert. Oregon State University, Corvallis.

ATLAS, E.L. & PYTKOWICZ, R.M. 1977: Solubility behaviour of apatites in seawater. Limnol. Oceanogr. 22, 290-300.

AVITAL, Y., STARINSKY, A. & KOLODNY, Y. 1983: Uranium geochemistry and fission track mapping of phosphorites, Zefa Field, Israel. Econ. Geol. 78, 121-131.

BACHRA, B.N., TRAUTZ, O.R. & SIMON, S.L. 1965: The effect of magnesium and fluoride ions on the spontaneous precipitation of calcium carbonates and phosphates. Arch. Oral Biol 10, 731-738.

BACQUET, G., VO QUANG, T., BONEL, G. & VIGNOLES, M. 1980: Résonance paramagnétique électronique du centre F dans les fluorapatites carbonatées de type B. J. Solid State Chem. 33, 189-195.

BAECHLE, H.-T. & WOLSTEIN, F. 1984: Cadmium compounds in mineral fertilisers. Proc. Fertil. Soc. London 226, 1-18.

BAROLE, D.V., RAJAGOPALAN, G. & SOMAYAJULU, B.L.K. 1987: Radiometric ages of phosphorites from the west coast of India. Mar. Geol. 78, 161-165.

BATURIN, G.N. 1978: Phosphorites on the Ocean Floor (in Russian). Izdat. Nauka, Moscow.

BATURIN, K.N. & KOCHENOV, A.V. 1974: Uranium content of oceanic phosphorites (in Russian). Litologia I Poleznye Iskopaemye 1, 124-129.

BATURIN, G.N. & ORESHKIN, V.N. 1984: Behaviour of cadmium in ocean-floor bone phosphate. Geochem. Int. 35, 69-74.

BATURIN, G.N., KOCHENOV, A.V. & PETELIN, V.P. 1970: Phosphorite formation on the shelf of SW Africa. Lithol. Min. Res. 3, 266-276.

BATURIN, G.N., MERKULOVA, K.I. & CHALOV, P.I. 1972: Radiometric evidence for recent formation of phosphatic nodules in marine shelf sediments. Mar. Geol. 13, M37-M41.

– 1974: Absolute dating of oceanic phosphorites by disequilibrium uranium. Geochemistry 11, 568-574.

BECKER, P. 1989: Phosphogypsum: expensive waste or profitable by-product. Phosphates and Phosphoric Acid. Marcel Dekker Inc. 537-569.

BENMORE, R.A., COLEMAN, M.L. & MCARTHUR, J.M. 1983: Origin of sedimentary francolite from its sulphur and carbon isotope composition. Nature 302, 516-518.

BENMORE, R.A., MCARTHUR, J.M. & COLEMAN, M.L. 1984: Stable isotopic composition of structural carbonate in sedimentary francolite. Spec. Publ. Geol. Surv. India 17, 35-40.

BENTOR, Y.K. 1980: Phosphorites – the unsolved problems. In: *Marine Phosphorites* (Ed. by BENTOR, Y.K.). Soc. Econ. Paleont. Mineral. Spec. Publ. 29, 3–18.

BERNAT, M. 1975: Les isotopes de l'uranium et du thorium et les terres rares dans l'environnement marin. Cah. ORSTOM Série Géol. 7, 65–83.

BERNER, R.A., RUTTENBERG, K.C., INGALL, E.D. & RAO, J.-L. 1993: The nature of phosphorus burial in modern marine sediments. In: *Interactions of C, N, P and S Biogeochemical Cycles and Global Change* (Ed. by WOLLAST, R., MACKENZIE F.T. & CHOU, L.). Springer-Verlag, Berlin, 365–378.

BERTRAM, C.J. & ELDERFIELD, H. 1993: The geochemical balance of the rare earth elements and neodymium isotopes in the oceans. *Geochim. Cosmochim. Acta* 57, 1957–1986.

BERTRAND-SARFATI, J., MOUSSINE-POUCHKINE, A., FLICOTEAUX, R. & AIT KACI, A. 1993: Ministromatolites phosphatés liés à la transgression glacio-eustatique d'âge Cambrien Inférieur (Sahara, Algérie). Abstract, 4th. French Congress of Sedimentology, Villeneuve d'Asq, 16–18 November 1993.

BIRCH, G.F., THOMSON, J., MCARTHUR, J.M. & BURNETT, W.C. 1983: Pleistocene phosphorites off the west coast of South Africa. *Nature* 302, 601–603.

BLISKOVSKIY, V.Z. 1976: On kurkite and francolite. *Lithol. Miner. Resour.* 11, 332–341.

BLISKOVSKIY, V.Z., YEFIMOVA, V.A. & ROMANOVA, L.V. 1967: The strontium contents of phosphorites. *Geochem. Internat.* 4, 1186–1190.

BLISKOVSKIY, V.Z., GRINENKO, V.A., MIGDISOV, L.I. & SAVINA, L.I. 1977: Sulphur isotopic composition of the minerals of phosphorites. *Geochem. Internat.* 14, 148–155.

BONEL, G., LABARTHE, J.C. & VIGNOLES, C. 1973: Contribution à l'étude structurale des apatites carbonatées de type B. *Compt. Rend. Coll. Internat. Physico-Chimie et la Cristallographie des apatites d'intérêt biologique*, Paris. 230, 117–125.

BONNOT-COURTOIS, C. & FLICOTEAUX, R. 1989: Distribution of rare-earth and some trace elements in Tertiary phosphorites from the Senegal Basin and their weathering products. *Chem. Geol.* 75, 311–328.

BORNEMAN-STARINKEVICH, I.D. & BELOV, N.V. 1940: Isomorphic substitution in carbonate-apatite. *Compt. Rend. Dokl. Acad. Sci. USSR* XXVI, 18, 804–806.

– 1953: Carbonate-apatites. *Dokl. Akad. Nauk. SSSR* 90, 89–92.

BOYLE, E.A. 1981: Cadmium, zinc, copper and barium in foraminiferal tests. *Earth Planet. Sci. Lett.* 53, 11–35.

BOYLE, E.A., SCLATER, F. & EDMOND, J.M. 1976: On the marine geochemistry of cadmium. *Nature* 263, 42–44.

BROWN, L.R., CHANDLER, W.C., FLAVIN, C., JACOBSON J., POLLOCK, C., POSTEL S., STARKE, L. & WOLF, E.C. 1987: *State of the World*. New York, W.W. Norton.

BRULAND, K.W. 1983: Trace elements in seawater. In: *Chemical Oceanography Volume 8* (Ed. by RILEY, J.P. & CHESTER, R.), Academic Press, London, 157–220.

BURKE, W.H., DENISON, R.E., HETHERINGTON, E.A., KOEPNICK, R.B., NELSON, H.F. & OTTO, J.B. 1982: Variation of  $^{87}\text{Sr}/^{86}\text{Sr}$  throughout Phanerozoic time. *Geology* 10, 516–519.

BURNETT, W.C. 1977: Geochemistry and origin of phosphorite deposits from off Peru and Chile. *Geol. Soc. Amer. Bull.* 88, 813–823.

BURNETT, W.C. & GOMBERG, G.N. 1977: Uranium oxidation and probable subaerial weathering of phosphatized limestone from the Pouriates Terrace. *Sedimentology* 24, 291–302.

BURNETT, W.C. & VEEH, H.H. 1977: Uranium-series disequilibrium studies in phosphorite nodules from the West Coast of South America. *Geochim. Cosmochim. Acta* 41, 755–764.

BURNETT, W.C., VEEH, H.H. & SOUTAR, A. 1980: U-series, oceanographic and sedimentary evidence in support of recent formation of phosphate nodules off Peru. In: *Marine Phosphorites* (Ed. by BENTOR, Y.K.). Soc. Econ. Paleont. Mineral. Spec. Publ. 29, 61–71.

BURNETT, W.C., BEERS, M.J. & ROE, K.K. 1982: Growth rates of phosphate nodules from the continental margin off Peru. *Science* 215, 1616–1618.

BURNETT, W.C., ROE, K.K. & PIPER, D.Z. 1983: Upwelling and phosphorite formation in the ocean. In: *Coastal Upwelling – its Sedimentary Record, Part A: Responses of the Sedimentary Regime to Present Coastal Upwelling* (Ed. by SUÈSS, E. & THIEDE, J.). Plenum, New York, 377–397.

BURNETT, W.C., COWART, J.B. & CHIN, P.A. 1987: Polonium in the surficial aquifer of west central Florida. NWWA Conference, Somerset, NJ. (Ed. by GRAVES, B.). 251–269.

BURNETT, W.C., BAKER, K.B., CHIN, P.A., McCABE, W. & DITCHBURN, R. 1988: Uranium-series and AMS $^{14}\text{C}$  studies of modern phosphatic pellets from Peru Shelf muds. In: *The Origin of Marine Phosphorite. The Results of the R.V. Robert D. Conrad Cruise 23-06 to the Peru Shelf* (Ed. by BURNETT, W.C. & FROELICH, P.N.). *Marine Geol.* 80, 215–230.

BURNETT, W.C., LAROCK, P.A., COWART, J.B. & HULL, C.D. 1992: Microbiology and Radiochemistry of Phosphogypsum. Year 2 Report to the Florida Institute of Phosphate Research, Bartow, FL.

BURNETT, W.C., HULL, C.D., CABLE P., CHERRIER, J., LAROCK, P.A. & HYUN J. 1993: Radionuclides in phosphogypsum: migration and bacterial control. Proc. Fourth Int. Conf. on the Chemistry and Migration Behavior of Actinides, Abstract, Charleston, S.C.

CAPO, C. & DEPAULO, D.J. 1990: Seawater strontium isotope variation from 2.5 million years to the present. *Science* 249, 51–55.

CARSON, G.A. & CROWLEY, S.F. 1993: The glauconite-phosphate association in hardgrounds: examples from the Cenomanian of Devon, southwest England. *Cret. Res.* 14, 69–89.

CATHCART, J.B. 1978: Uranium in phosphate rock. US Geol. Surv. Prof. Pap. 988-A, 1–6.

— 1992: Uranium in phosphate rock with special reference to the Central Florida deposits. US Geol. Surv. Circ. 1069, 32–35.

CHERRIER, J., LAROCK P.A. & BURNETT, W.C. (in press): Uptake and cellular distribution patterns of polonium by a phosphogypsum bacterial isolate. *Jour. Appl. Environ. Microbiology*.

CHIEN, S.H. & BLACK, C.A. 1976: Free energy of formation of carbonate apatites in some phosphate rocks. *Soil Sci. Soc. Amer. Proc.* 40, 234–239.

CLAYPOOL, G.E., HOLZER, W.T., KAPLAN, I.R., SAKAI, H. & ZAK, I. 1980: The age curve of sulphur and oxygen isotopes in marine sulphates and their mutual interpretation. *Chem. Geol.* 28, 199–260.

COLLEY, S & THOMSON, J. 1985: Recurrent uranium relocations in distal turbidites emplaced in pelagic conditions. *Geochim. Cosmochim. Acta* 49, 2339–2348.

COMPTON, J.S., SNYDER, S.W. & HODELL, D.A. 1990: Phosphogenesis and weathering of shelf sediments from the south-eastern United States: Implications for Miocene  $\delta^{13}\text{C}$  excursions and global cooling. *Geology* 18, 1227–1230.

COMPTON, J.S., HODELL, D.A., GARRIDO, J.R. & MALLINSON, D.J. 1993: Origin and age of phosphorite from the south-central Florida Platform: Relation of phosphogenesis to sea-level fluctuations and  $\delta^{13}\text{C}$  excursions. *Geochim. Cosmochim. Acta* 57, 131–146.

COOK, P.J. 1972: Petrology and geochemistry of the phosphate deposits of Northwest Queensland, Australia. *Econ. Geol.* 67, 1193–1213.

— 1984: Spatial and temporal controls on the formation of phosphorite deposits – a review. In: *Phosphate Minerals* (Ed. By NRIAGU, J.O. & MOORE, P.B.). Springer, Heidelberg, 242–274.

COOK, P.J. & MCLEHINNY, M.W. 1979: A reevaluation of the spatial and temporal distribution of phosphate deposits in the light of plate tectonics. *Econ. Geol.* 74, 315–330.

COOK, P.J., SHERGOLD, J.H., BURNETT, W.C. & RIGGS, S.R., 1990: Phosphorite research: a historical overview. In: *Phosphorite Research and Development* (Ed. by NOTHOLT A.J.G. & JARVIS, I.). Geol. Soc. Spec. Publ. 52, 1–22.

COSSA, 1878: Sur la diffusion de cerium, du lanthane et du didyme, extract of a letter from Cossa to Sella, presented by Frény. *Compt. Rend. Acad. Sci. Paris* 87, 378–388.

CULLEN, D.J. & BURNETT, W.C. 1986: Phosphorite associations on seamounts in the tropical south-west Pacific. *Mar. Geol.* 71, 215–236.

DAR'IN, A.V. & ZANIN, YU. N. 1990: Rare-earth elements in a series of supergene transformations of phosphorites (in Russian). *Geologiya i Geofizika* 31, 89–97.

DA ROCHA ARAUJO, P.R., FLICOTEAUX, R., PARRON, C. & TROMPETTE, R., 1992: Phosphorites of Rocinha Mine – Patos de Minas (Minas Gerais, Brazil): Genesis and evolution of a Middle Proterozoic deposit tectonized by the Brasiliano Orogeny. *Econ. Geol.* 87, 332–351.

DENISON, R.E., KOEPNICK, R.B., FLETCHER, A., HOWELL, M.W. & CALLAWAY, W.S. 1994a: Criteria for the retention of original seawater  $^{87}\text{Sr}/^{86}\text{Sr}$  in ancient shelf limestones. *Chem. Geol.* 112, 131–143.

DENISON, R.E., KOEPNICK, R.B., BURKE, W.H., HETHERINGTON, E.A. & FLETCHER, A. 1994b: Construction of the Mississippian, Pennsylvanian and Permian seawater  $^{87}\text{Sr}/^{86}\text{Sr}$  curve. *Chem. Geol.* 112, 145–167.

DEPAULO, D.J. & INGRAM, B. 1985: High-resolution stratigraphy with strontium isotopes. *Science* 227, 938–941.

DEPAULO, D.J. & WASSERBURG, G.J. 1976: Nd isotopic variations and petrogenetic models. *Geophys. Res. Lett.* 3, 249–252.

ELDERFIELD, H. 1986: Strontium isotope stratigraphy. *Palaeogeog. Palaeoclimatol. Palaeoecol.* 57, 71–90.

ELDERFIELD, H. & GREAVES, M.J. 1982: The rare earth elements in seawater. *Nature* 296, 214–219.

ELDERFIELD, H. & PAGETT, R. 1986: REE in ichthyoliths: variations with redox conditions and depositional environment. *Sci. Total Environ.* 49, 175–197.

ELDERFIELD, H. & SHOLKOVITZ, E.R. 1987: Rare earth elements in the pore waters of reducing nearshore sediments. *Earth Planet. Sci. Lett.* 82, 280–288.

EL-KAMMAR, A.M., ZAYED, M.A. & AMER, S.A. 1979: Rare earths of the Nile Valley phosphorites, Upper Egypt. *Chem. Geol.* 24, 69–81.

EMERY, D. & ROBINSON, A. 1993: Inorganic Geochemistry: Applications to Petroleum Geology. Blackwell, Oxford.

EPSTEIN, S., BUCHSBAUM, R., LOWENSTAM, H.A. & UREY, H.C. 1953: Revised carbonate-water temperature scale. *Geol. Soc. Amer. Bull.* 62, 417–426.

FLICOTEAUX, R., 1982: Genèse des phosphates alumineux de Sénégal occidental étapes et guides de l'altération. *Sci. Géol. Mém.* 67.

FLICOTEAUX, R. & LUCAS, J., 1984: Weathering of phosphate minerals. In: *Phosphate Minerals* (Ed. by NRIAGU, J.O. & MOORE, P.B.). Springer-Verlag, Berlin, 292–317.

FLICOTEAUX, R., WALTER A.-V., BONNOT-COURTOIS, C. & TOLEDO-GROKE, M.-C., 1990: Transformation and precipitation of phosphates during weathering: characterization by REE distributions. *Geochemistry of the Earth's Surface and of Mineral Formation*, 2nd Int. Symp, Aix-en-Provence, 365–367.

FLICOTEAUX, R., PANNATIER, Y. & BONNOT-COURTOIS, C. 1993: Palaeoceanographic conditions of phosphate precipitation in the western part of the Palaeogene Senegal Basin. Contribution of geochemical data. *Concepts and Controversies in Phosphogenesis*. Int. Symp. and Workshop, Matten, 6–10 September, 1993.

FROELICH, P.N., KLINKHAMMER, G.P., BENDER, M.L., LUETKE, N.A., HEATH, G.R., CULLEN, D., DAUPHIN, P., HAMMOND, D., HARTMAN, B. & MYNARD, V. 1979: Early oxidation of organic matter in pelagic sediments of the eastern equatorial Atlantic: suboxic diagenesis. *Geochim. Cosmochim. Acta* 43, 1075–1090.

FROELICH, P.N., KIM, K.H., JAHNKE, R., BURNETT, W.C., SOUTAR, A. & DEAKIN, M. 1983: Pore water fluoride in Peru continental margin sediments: Uptake from seawater. *Geochim. Cosmochim. Acta* 47, 1605–1612.

FROELICH, P.N., ARTHUR, M., BURNETT, W.C., DEAKIN, M., HENSLEY, V., JAHNKE, R., KAUL, L., KIM, K., ROE, K., SOUTAR, A. & VATAKANON, C. 1988: Early diagenesis of organic matter in Peru continental margin sediments: Phosphate precipitation. In: *The Origin of Marine Phosphorite. The Results of the R.V. Robert D. Conrad Cruise 23–06 to the Peru Shelf* (Ed. by BURNETT, W.C. & FROELICH, P.N.). *Marine Geol.* 80, 309–343.

GAVSHIN, V.M., BOBROV, V.A. & ZORKINA, L.S. 1974: Quantitative relations between uranium and phosphorus in phosphorites and phosphatic sedimentary rocks. *Lithol. Miner. Res.* 9, 740–746.

GAUTHIER, M.J., CLEMENT, R.L., FLATAU, G.N. & AMIARD, J.C. 1986: Accumulation du cadmium par les bactéries marines à Gram négatif selon leur sensibilité au métal et leur type respiratoire. *Oceanol. Acta* 9, 333–337.

GILINSKAYA, L.G. 1991: A new type of  $\text{PO}_4^{3-}$  centre in apatite. *J. Struct. Chem.* 31, 892–898.

– 1993: Stable paramagnetic  $\text{Pb}^{3+}(^2\text{S}_{1/2})$  centers in natural  $\text{Ca}_5(\text{PO}_4)_3(\text{F},\text{Cl},\text{OH})$  apatites. *Phys. Solid State* 35, 35–37.

GILINSKAYA, L.G. & ZANIN, YU. N. 1983: EPR investigation of the isomorphic impurity in the form  $\text{VO}^{2+}$  in apatite from phosphorite (in Russian). *Dokl. Akad. Nauk. SSSR* 273, 1463–1467.

GILINSKAYA, L.G., ZANIN, YU. N., KNUBOVETS, R.G., KORNEVA, T.A. & FADEEVA, V.P. 1993: Organophosphorus radicals in natural apatites ( $\text{Ca}_5(\text{PO}_4)_3(\text{F},\text{OH})$ ). *J. Struct. Chem.* 33, 859–870.

GIRARD, J.-P., FLICOTEAUX, R., WALTER, A.-V., SAVIN, S.M. & NAHON, D. 1993: Oxygen and carbon stable isotope composition of structural carbonate in weathering apatites from laterites, southern Brazil and western Senegal. *Appl. Geochem.* 8, 617–632.

GLENN, C.R. 1990a: Pore water, petrologic and stable carbon isotopic data bearing on the origin of Modern Peru margin phosphorites and associated authigenic phases. In: *Phosphate Deposits of the World Volume 3: Neogene to Recent Phosphorites*. (Ed. by BURNETT, W.C. & RIGGS, S.R.). Cambridge University Press, Cambridge, 46–61.

– 1990b: Depositional sequences of the Duwi, Sibâya and Phosphate Formations, Egypt: phosphogenesis and glauconitization in a Late Cretaceous epeiric sea. In: *Phosphorite Research and Development* (Ed. by NOTHOLT A.J.G. & JARVIS, I.). *Geol. Soc. Spec. Publ.* 52, 205–222.

GLENN, C.R. & ARTHUR, M.A. 1988: Petrology and major element geochemistry of Peru margin phosphorites and associated diagenetic minerals: authigenesis in modern organic-rich sediments. In: *The Origin of Marine Phosphorite. The Results of the R.V. Robert D. Conrad Cruise 23–06 to the Peru Shelf* (Ed. by BURNETT, W.C. & FROELICH, P.N.). *Marine Geol.* 80, 231–267.

GLENN, C.R., ARTHUR, M.A., YEH, H.W. & BURNETT, W.C. 1988: Carbon isotopic composition and lattice-bound carbonate of Peru-Chile margin phosphorites. In: *The Origin of Marine Phosphorite. The Results of the R.V. Robert D. Conrad Cruise 23–06 to the Peru Shelf* (Ed. by BURNETT, W.C. & FROELICH, P.N.). *Marine Geol.* 80, 287–307.

GLENN, C.R., ARTHUR, M.A., RESIG, M.J., BURNETT, W.C., DEAN, W.E. & JAHNKE, R.A. 1994a: Are modern and ancient phosphorites really so different? In: *Siliceous, Phosphatic and Glauconitic Sediments of the Tertiary and Mesozoic* (Ed. by IJIMA, A., ABED, A.M. & GARRISON, R.E.). *Proc. 29th Int. Geol. Congr.*, Part C, VSP, Utrecht, 159–188.

GLENN, C.R., FÖLLMI, K.B., RIGGS, S.R., BATURIN, G.N., GRIMM, K.A., TRAPPE, J., ABED, A.M., GALLI-OLIVIER, C., GARRISON, R.E., ILYIN, A., JEHL, C., ROHRLICH, V., SADAQAH, R.M., SCHIDLOWSKI, M., SHELDON, R.E. & SIEGMUND, H. 1994b: Phosphorus and phosphorites: Sedimentology and environments of formation. *Eclogae geol. Helv.* 87, 747–788.

GOLDBERG, E.D., KOIDE, M., SCHMITT, R.A. & SMITH, H.V. 1963: Rare earth distribution in the marine environment. *J. Geophys. Res.* 68, 4209–4217.

GONY, J.N. 1971: Étude cristallochimique du phosphate uranifère de Bakouma (République centrafricaine). Commissariat à l'Energie Atomique, CEN Saclay, Report CEA-R-400E.

GRAHAM, D.W., BENDER, M.L., WILLIAMS, D.F. & KEIGWIN JR., L.D. 1982: Strontium-calcium ratios in Cenozoic planktonic foraminifera. *Geochim. Cosmochim. Acta* 46, 1281–1292.

GRANDJEAN, P. & ALBARÈDE, F. 1989: Ion probe measurement of rare earth element in biogenic phosphates. *Geochim. Cosmochim. Acta* 53, 3179–3183.

GRANDJEAN, P., CAPPETA, H., MICHARD, A. & ALBARÈDE, F. 1987: The assessment of REE patterns and  $^{143}\text{Nd}/^{144}\text{Nd}$  ratios in fish remains. *Earth Planet. Sci. Lett.* 84, 181–196.

GRANDJEAN, P., CAPPETA, H. & ALBARÈDE, F. 1988: The REE and  $\epsilon\text{Nd}$  of 40–70 Ma old fish debris from the West-African platform. *Geophys. Res. Lett.* 15, 389–392.

GRANDJEAN-LÉCUYER P., FEIST, R. & ALBARÈDE, F. 1993: Rare earth elements in old biogenic apatites. *Geochim Cosmochim. Acta* 57, 2507–2514.

GRUNER, J.W. & McCONNELL, D. 1937: The problem of carbonate-apatites. The structure of francolite. *Z. Krist. Abt. A* 97, 208–215.

GULBRANDSEN, R.A. 1966: Chemical composition of the phosphorites of the Phosphoria Formation. *Geochim. Cosmochim. Acta* 30, 769–778.

– 1970: Relation of carbon dioxide content of apatite of the Phosphoria Formation to regional facies. US Geol. Surv. Prof. Pap. 700-B, 9–13.

– 1975: Analytical data on the Phosphoria Formation, Western United States. US Geol. Surv. Open-File Rep. 75–554.

GULBRANDSEN, R.A., KRAMER, J.R., BESTY, L.B. & MAYS, R.E. 1966: Carbonate-bearing apatite from Faraday Township, Ontario, Canada. *Amer. Mineral.* 51, 819–824.

GULBRANDSEN, R.A., ROBERSON, C.E. & NEIL, S.T. 1984: Time and the crystallisation of apatite in seawater. *Geochim. Cosmochim. Acta* 48, 213–218.

GUSEV, G.M., ZANIN, YU.N., KRIVOPUTSKAYA, L.M., LEMINA, N.M. & IUSUPOV, T.S. 1976: Transformation of apatite composition under conditions of weathering and leaching. *Dokl. Acad. Nauk. SSSR* 229, 971–973.

HARADA, K., BURNETT, W.C., LAROCK, P.A. & COWART, J.B. 1989: Polonium in Florida groundwater and its relationship to the sulfur cycle and bacteria. *Geochim. Cosmochim. Acta* 53, 143–150.

HEGGIE, D.T., SKYRING, G.W., O'BRIEN, G.W., REIMERS, C., HERCZEG, A., MORIARTY, D.J.W., BURNETT, W.C. & MILNES, A.R. 1990: Organic carbon cycling and modern phosphorite formation on the East Australian continental margin: an overview. In: *Phosphorite Research and Development* (Ed. by NOTHOLT, A.J.G. & JARVIS, I.). *Geol. Soc. Spec. Publ.* 52, 87–117.

HEIN, J.R., YEH, H.-W., GUNN, S.H., SLITER, W.V., BENNINGER, L.M. & WANG, C.-H. 1993: Two major Cenozoic episodes of phosphogenesis recorded in Equatorial Pacific seamount deposits. *Paleoceanogr.* 8, 293–311.

HENDERSON, G.M., COHEN, A.S. & O'NIIONS, R.K. 1993:  $^{234}\text{U}/^{238}\text{U}$  ratios and  $^{230}\text{Th}$  ages for Hateruma Atoll corals: implications for coral diagenesis and seawater  $^{234}\text{U}/^{238}\text{U}$  ratios. *Earth Planet. Sci. Lett.* 15, 65–73.

HESS, J., BENDER, M.L. & SCHILLING, J.-G. 1986: Evolution of the ratio of strontium-87 to strontium-86 in seawater from Cretaceous to Present. *Science* 231, 979–984.

HODELL, D.A., MUELLER, P.A. & GARRIDO, J.R. 1991: Variations in the strontium isotopic composition of seawater during the Neogene. *Geology* 19, 24–27.

HOWARD, P.F. & HOUGH, M.J. 1979: On the geochemistry and origin of the D-Tree, Wonarah and Sherrin Creek phosphorite deposits of the Georgina Basin, Northern Australia. *Econ. Geol.* 74, 260–284.

HOYLE, J., ELDERFIELD, H., GLEDHILL, A. & GREAVES, M. 1984: The behaviour of rare earth elements during mixing of river and seawaters. *Geochim. Cosmochim. Acta* 48, 143–149.

HUDSON, J.D. 1977: Stable isotopes and limestone lithification. *J. Geol. Soc.* 133, 637–660.

ILYIN, A.V. & RATNIKOVA, G.I. 1976: Rare-earth element distribution in the Hobso Gol phosphorites (Mongolia). *Geochem. Int.* 13, 53–56.

ILYIN, A.V., ZAITSEV, N.S. & BJAMBA, Z. 1986: Proterozoic and Cambrian phosphorites deposits: Khubsugul, Mongolian People's Republic. In: *Phosphate Deposits of the World Vol. 1, Proterozoic and Cambrian Phosphorites* (Ed. by COOK, P.J. & SHERGOLD, J.H.). 162–174.

INGRAM, B.L., COCCIONI, R., MONTANARI, A. & RICHTER, F.M. 1994: Strontium isotopic composition of mid-Cretaceous seawater. *Science* 264, 546–550.

IRWIN, H., CURTIS, C. & COLEMAN, M. 1977: Isotopic evidence for source of diagenetic carbonates formed during burial of organic rich sediments. *Nature* 269, 209–213.

ISHERWOOD, K. F. 1992: Phosphate Industry and the Environment. In: *Phosphate Fertilizers and the Environment* (Ed. by SCHULTZ, J. J.). Tampa FL, 115–124.

JACOB, K.D., HILL, W.L., MARSHALL, M.R. & REYNOLDS, D.S. 1933: The composition and distribution of phosphate rock with special reference to the United States. US Dept. Agric. Techn. Bull. 364.

JACOBSEN, S.B. & WASSERBURG, G.J. 1980: Sm-Nd isotopic evolution of chondrites. *Earth Planet. Sci. Lett.* 50, 139–155.

JAHNKE, R.A. 1984: The synthesis and solubility of carbonate fluorapatite. *Am. J. Sci.* 284, 58–78.

JAHNKE, R.A., EMERSON, S.R., ROE, K.K. & BURNETT, W.C. 1983: The present day formation of apatite in Mexican continental margin sediments. *Geochim. Cosmochim. Acta* 47, 259–266.

JALLAD, I.S., ABU MURREY, O.S. & SADAQAH, R.M. 1989: Upper Cretaceous phosphorites of Jordan. In: *Phosphate Deposits of the World Volume 2: Phosphate Rock Resources* (Ed. by NOTHOLT, A.J.G., SHELDON, R.P. & DAVIDSON, D.F.). Cambridge University Press, Cambridge, 344–351.

JARVIS, I. 1984: Rare-earth element geochemistry of late Cretaceous chalks and phosphorites from northern France. *Spec. Publ. Geol. Surv. India* 17, 179–190.

– 1992: Sedimentology, geochemistry and origin of phosphatic chalks, the Upper Cretaceous deposits of NW Europe. *Sedimentology* 39, 55–97.

JARVIS, I. & JARVIS, K.E. 1991: Geochemical variation in phosphorite nodules: a study using laser ablation inductively coupled plasma-mass spectrometry. In: *First Gentner Symposium on Geoscience. Phosphorites and Black Shales*. Jerusalem, September 1991, Program with Abstracts, 22.

JEANDEL, C. 1993: Concentration and isotopic composition of Nd in the South Atlantic Ocean. *Earth Planet. Sci. Lett.* 117, 581–591.

JEANMAIRE, J.-P. 1985: Répartition de l'uranium dans les niveaux phosphatés Maestrichtien supérieur – Éocène inférieur du secteur de Benguerir (Bassin des Ganntour, Maroc occidental). *Sci. Géol. Mém.* 77, 53–68.

JOHNSON, A.K.C. 1987: Le bassin côtier à phosphates du Togo (Maastrichtien-Eocène moyen). Thèse Univ. Dijon.

JOHNSTON, A.E. & JONES, K.C. 1992: The cadmium issue – long-term changes in the cadmium content of soils and the crops grown on them. In: *Phosphate Fertilizers and the Environment*, Proc. Int. Workshop, IFDC, March 1992 Tampa, Florida, 255–269.

JONES, E.J.W. 1989: Radioactivity of the ocean floor and marine phosphorite deposits: observations with a new deep-towed scintillometer. *Geophys. Res. Lett.* 16, 123–126.

KAPLAN, I.R. 1983: Stable isotopes of sulphur, nitrogen and deuterium in Recent marine environments. In: *Stable Isotopes in Sedimentary Geology*, SEPM Short Course 10, 2.1–2.108.

KARHU, J. & EPSTEIN, S. 1986: The implication of the oxygen isotope records in coexisting cherts and phosphates. *Geochim. Cosmochim. Acta* 1745–1757.

KASTNER, M., GARRISON, R.E., KOLODNY, Y., REIMERS, C.E. & SHEMESH, A. 1990: Coupled changes of oxygen isotopes in  $\text{PO}_4^{3-}$  and  $\text{CO}_3^{2-}$  in apatite, with emphasis on the Monterey Formation, California. In: *Phosphate Deposits of the World Volume 3: Neogene to Modern Phosphorites* (Ed. by BURNETT, W.C. & RIGGS, S.R.). Cambridge University Press, Cambridge, 312–324.

KAUFMAN, A.J., JACOBSEN, S.B. & KNOLL, A.H. 1993: The Vendian record of Sr and C isotopic variations in seawater: implications for tectonics and paleoclimate. *Earth Planet. Sci. Lett.* 120, 409–430.

KAZAKOV, A.V. 1937: The phosphorite facies and the genesis of phosphorites. In: *Geological Investigations of Agricultural Ores*. Trans. Sci. Inst. Fertil. Insecto-Fungicides 142, 93–113.

KETO, L.S. & JACOBSEN, S.B. 1987: Nd and Sr isotopic variations in Early Paleozoic oceans. *Earth Planet. Sci. Lett.* 84, 27–41.

– 1988: Nd isotopic variations of Phanerozoic paleoceans. *Earth Planet. Sci. Lett.* 90, 395–410.

KIM, K.H. & BURNETT, W.C. 1986: Uranium-series growth history of a Quaternary phosphatic crust from the Peruvian continental margin. *Chem. Geol.* 58, 227–244.

KOEPNICK, R.B., BURKE, W.H., DENISON, R.E., HETHERINGTON, E.A., NELSON, H.F., OTTO, J.B. & WAITE, L.E. 1985: Construction of the seawater  $^{87}\text{Sr}/^{86}\text{Sr}$  wave for the Cenozoic and Cretaceous: supporting data. *Chem. Geol. (Iso. Geosci. Sect.)* 58, 55–81.

KOLODNY, Y. 1969: Are marine phosphorites forming today? *Nature* 244, 1017–1019.

– 1981: Phosphorites In: *The Sea Volume 7* (Ed. by EMILIANI, C.). Wiley Interscience, New York, 981–1023.

KOLODNY, Y. & KAPLAN, I.R. 1970a: Uranium isotopes in sea floor phosphorites. *Geochim. Cosmochim. Acta* 34, 3-24.

- 1970b: Carbon and oxygen isotopes in apatite  $\text{CO}_2$  and co-existing calcite from sedimentary phosphorite. *J. Sediment. Petrol.* 40, 954-959.

KOLODNY, Y. & LUZ, B. 1992a: Isotope signatures in phosphate deposits; Formation and diagenetic history. In: *Isotopic Signatures and Sedimentary Records*, Lecture Notes in Earth Sciences No. 43 (Ed. by CLAUSER, N. & CHAUDHURI, S.). Springer-Verlag, Berlin, Heidelberg, 69-121.

- 1992b: Oxygen isotopes in phosphates of fossil fish – Devonian to Recent. In: *Stable Isotope Geochemistry: A Tribute to Samuel Epstein* (Ed. by TAYLOR, H.P., O-NEIL, J.R. & KAPLAN, I.R.). *Geochem. Soc. Spec. Publ.* 3, 105-119.

KOLODNY, Y. & RAAB, M. 1988: Oxygen isotopes in phosphatic fish remains from Israel: Paleothermometry of tropical Cretaceous and Tertiary shelf waters. *Palaeogeog. Palaeoclimatol. Palaeoecol.* 64, 59-67.

KOLODNY, Y., LUZ, B. & NAVON, O. 1983: Oxygen isotope variations in phosphate of biogenic apatite. I: Fish bone apatite – rechecking the rules of the game. *Earth Planet. Sci. Lett.* 64, 398-404.

KOULOHERIS, A. P. 1980: Chemical nature of phosphogypsum as produced by various wet process phosphoric acid processes. *Proc. Int. Symp. Phosphogypsum*. (Ed. by BORRIS, D. P. & BODY, P. W.). Florida Institute of Phosphate Research 8-35.

KRAJEWSKI, K.P., VAN CAPPELLEN, P., TRICHET, J., KUHN O., LUCAS, J., MARTIN-ALGARRA, A., PRÉVÔT, L., TEWARI, V.C., GASPAR, L., KNIGHT, R.I. & LAMBOY, M. 1994: Biological processes and apatite formation in sedimentary environments. *Eclogae geol. Helv.* 87, 701-745.

KRESS, A.G. & VEEH, H.H. 1980: Geochemistry and radiometric ages of phosphatic nodules from the continental margin of northern New South Wales, Australia. *Mar. Geol.* 36, 143-157.

LEHR, J.R., MCCLELLAN, G.H., SMITH, J.P. & FRAZIER, A.W. 1968: Characterisation of apatites in commercial phosphate rock. *Coll. int. Phosphates Minéraux Solides*, Toulouse 1967 2. *Soc. Chim. Fr.*, Paris 29-44.

LINDEKEN C.L. 1980: Radiological considerations of phosphogypsum utilization in agriculture. *Proc. Second Int. Symp. on Phosphogypsum*. (Ed by BORRIS, D.P. & BODY, P.W.). Florida Institute of Phosphate Research, 459-80.

LINDEKEN, C.L. & COLES, D.G. 1978: The radium-226 content of agricultural gypsums. *Radioactivity in Consumer Products*. (Ed. by MOGHISSI, A., PARAS, P., CARTER, M. & BARKER, R.). 369-75.

LONGINELLI, A. & NUTI, S. 1968: Oxygen isotopic composition of phosphorites from marine formations. *Earth Planet. Sci. Lett.* 5, 13-16.

- 1973a: Revised phosphate-water isotopic temperature scale. *Earth Planet. Sci. Lett.* 19, 373-376.
- 1973b: Oxygen isotope measurements of phosphate from fish teeth and bones. *Earth Planet. Sci. Lett.* 20, 337-340.

LUCAS, J. & ABBAS, M. 1989: Uranium in natural phosphorites: the Syrian example. *Sci. Géol. Bull.* 42, 223-236.

LUCAS, J. & PRÉVÔT, L. 1981: Synthèse d'apatite à partir de matière organique phosphate (ARN) et de calcite par voie bactérienne. *Compt. Rend. Acad. Sci. Paris* 292, 1203-1208

- 1984: Synthèse d'apatite par voie bactérienne à partir de matière organique phosphatée et de divers-carbonates de calcium dans des eaux douce et marine naturelles. *Chem. Geol.* 42, 101-118.
- 1985: The synthesis of apatite by bacterial activity: mechanism. In: *Phosphorites* (Ed. by LUCAS, J. & PRÉVÔT, L.). *Sci. Géol. Mém.* 77, 83-92.

LUCAS, J., PRÉVÔT, J. & LAMBOY, M. 1978: Les phosphorites de la marge nord de l'Espagne. *Chimie, minéralogie, genèse. Oceanol. Acta* 1, 55-72.

LUCAS, J., FLICOTEAUX, R., NATHAN, Y., PRÉVÔT, L. & SHAHAR, Y. 1980: Different aspects of phosphorite weathering. In: *Marine Phosphorites – Geochemistry, Occurrence, Genesis*. (Ed. by BENTOR, Y.K.). *Soc. Econ. Paleontol. Mineral.* 29, 41-51.

LUCAS, J., ILYIN, A.V. & KUHN, A. 1986: Proterozoic and Cambrian phosphorites deposits: Volta basin, West Africa. In: *Phosphate Deposits of the World Vol. 1, Proterozoic and Cambrian Phosphorites* (Ed. by COOK, P.J. & SHERGOLD, J.H.). 235-243.

LUCAS, J., EL FALEH, E. M. & PRÉVÔT, L. 1990: Experimental study of the substitution of Ca by Sr and Ba in synthetic apatites. In: *Phosphorite Research and Development* (Ed. by NOTHOLT, A.J.G. & JARVIS, I.). *Geol. Soc. Spec. Publ.* 52, 33-47.

MALONE, P.H.G. & TOWE, K.M. 1970: Microbial carbonate and phosphate precipitates from seawater cultures. *Mar. Geol.* 9, 301-309.

MANHEIM, F.T., ROWE, G.T. & JIPPA, D. 1975: Marine phosphorite formation off Peru. *J. Sed. Petrol.* 45, 243-251.

MARSHALL, J.F. & COOK, P.J. 1980: Petrology of iron- and phosphorus-rich nodules from the Eastern Australian continental shelf. *J. Geol. Soc.* 137, 765-771.

MARTENS, C.S. & HARRIS, R.C. 1970: Inhibition of apatite precipitation in the marine environment by magnesium ions. *Geochim. Cosmochim. Acta* 34, 621–625.

MARTIN, J.H. & KNAUER, G.A. 1973: The elemental composition of plankton. *Geochim. Cosmochim. Acta* 37, 1639–1653.

MARTIN-ALGARRA, A., SANCHEZ-NAVAS, A. & VERA, J.A. 1993: Phosphate stromatolites in the pelagic Mesozoic record of the Betic Cordillera (southern Spain). Concepts and Controversies in Phosphogenesis. Int. Symp. and Workshop, Matten, 6–10 September, 1993.

MATTHEWS, A. & NATHAN, Y. 1977: The decarbonation of carbonate fluorapatite (francolite). *Amer. Mineral.* 62, 565–573.

MAY, A. & SWEENEY, J. W. 1982: Assessment of environmental impacts associated with phosphogypsum in Florida. US Bureau of Mines Report 8639, 1–19.

MCARTHUR, J.M. 1978a: Systematic variations in the contents of Na, Sr, CO<sub>2</sub> and SO<sub>4</sub> in marine carbonate fluorapatite and their relation to weathering. *Chem. Geol.* 21, 41–52.

- 1978b: Element partitioning in ferruginous and pyritic phosphorite from the continental margin off Morocco. *Mineral. Mag.* 42, 221–228.
- 1980: Post-depositional alteration of the carbonate-fluorapatite phase of Moroccan Phosphates. In: *Marine Phosphorites – Geochemistry, Occurrence, Genesis.* (Ed. by BENTOR, Y.K.). Soc. Econ. Paleontol. Mineral. 29, 53–60.
- 1985: Francolite geochemistry – compositional controls during formation, diagenesis, metamorphism and weathering. *Geochim. Cosmochim. Acta* 49, 23–35.
- 1990: Fluorine-deficient apatite. *Min. Mag.* 54, 508–510.
- 1994: Recent trends in strontium isotope stratigraphy. *Terra Nova* 6, 331–358.

MCARTHUR, J.M. & HERCZEG, A. 1990: Diagenetic stability of the isotopic composition of phosphate oxygen: paleoenvironmental implications. In: *Phosphorite Research and Development* (Ed. by NOTHOLT, A.J.G. & JARVIS, I.). *Geol. Soc. Spec. Publ.* 52, 119–124.

MCARTHUR, J.M. & WALSH, J.N. 1984: Rare-earth geochemistry of phosphorites. *Chem. Geol.* 47, 191–220.

MCARTHUR, J.M., COLEMAN, M.L. & BREMNER, J.M. 1980: Carbon and oxygen isotopic composition of structural carbonate in sedimentary francolite. *J. Geol. Soc.* 137, 669–673.

MCARTHUR, J.M., BENMORE, R.A., COLEMAN, M.L., SOLDI, C., YEH, H.-W. & O'BRIEN, G.W. 1986: Stable isotope characterisation of francolite formation. *Earth Planet. Sci. Lett.* 77, 20–34.

MCARTHUR, J.M., HAMILTON, P.J., GREENSMITH, J.T., BOYCE, A.J., FALICK, A.E., BIRCH, G., WALSH, J.N., BENMORE, R.A. & COLEMAN, M.L. 1987: Phosphorite geochemistry: isotopic evidence for meteoric alteration of francolite on a local scale. *Chem. Geol. (Isotope Geosci. Sect.)* 65, 415–425.

MCARTHUR, J.M., THOMSON, J., JARVIS, I., FALICK, A.E. & BIRCH, G.F. 1988: Eocene to Pleistocene phosphogenesis off western South Africa. *Mar. Geol.* 85, 41–63.

MCARTHUR, J.M., SAHAMI, A.R., THIRLWALL, M., HAMILTON, P.J. & OSBORN, A.O. 1990: Dating phosphogenesis with strontium isotopes. *Geochim. Cosmochim. Acta* 54, 1343–1351.

MCARTHUR, J.M., THIRLWALL, M.F., CHEN, M., GALE, A.S. & KENNEDY, W.J. 1993a: Strontium isotope stratigraphy in the Late Cretaceous: numerical calibration of the Sr isotope curve and intercontinental correlation for the Campanian. *Paleoceanography* 8, 859–873.

MCARTHUR, J.M., THIRLWALL, M.F., GALE, A.S., KENNEDY, W.J., BURNETT, J.A., MATTEY, D. & LORD, A.R. 1993b: Strontium isotope stratigraphy for the Late Cretaceous: a new curve based on the English Chalk. In: *High Resolution Stratigraphy* (Ed. by HAILWOOD, E.A. & KIDD, R.B.). *Geol. Soc. Spec. Publ.* 70, 195–209.

MCCONNELL, D. 1973: Apatite – Its Crystal Chemistry, Mineralogy, Utilization, and Geologic and Biologic Occurrences. Springer Verlag, Vienna–Heidelberg–New York.

MCCLELLAN, G.H. 1980: Mineralogy of carbonate fluorapatites. *J. Geol. Soc. London* 137, 675–681.

MCCLELLAN, G.H. & LEHR, J.R. 1969: Crystal chemical investigations of natural apatites. *Amer. Mineral.* 54, 1374–1391.

MCCLELLAN, G.H. & SAAVEDRA, F.N. 1986: Chemical and mineral characteristics of some Cambrian and Precambrian phosphorites. In: *Phosphorite Deposits of the World Volume 1, Precambrian and Cambrian Phosphorites* (Ed. by COOK, P.J. & SHERGOLD, J.H.). Cambridge University Press, Cambridge 244–267.

MCCLELLAN, G.H. & VAN KAUWENBERGH, S.J. 1990: Mineralogy of sedimentary apatites. In: *Phosphorite Research and Development* (Ed. by NOTHOLT A.J.G. & JARVIS, I.). *Geol. Soc. Spec. Publ.* 52, 23–31.

MFARLIN, R. F. 1992: Current and anticipated research priorities for the utilization and management of phosphate process by-products and wastes. *Phosphate Fertilizers and the Environment* (Ed. by SCHULTZ, J. J.). Tampa, FL., 179–184.

McKELVEY, V.E. 1950: Rare earths in western phosphate rocks. US Geol. Surv. Trace Element Mem. Rept 194.

— 1956: Uranium in phosphate rock. US Geol. Surv. Prof. Pap. 300, 477–481.

— 1967: Phosphate deposits. US Geol. Surv. Bull. 1252-D, 1–21.

McKELVEY, V.E., SWANSON, R.W. & SHELDON, R.P. 1953: The Permian phosphorite deposits of the Western United States. In: Compt. Rend. Congr. Geol. Int. 19th Algiers 1952, 21, 45–64.

McKELVEY, V.E., STROBELL, J.D. JR. & SLAUGHTER, A.L. 1986: The vanadiferous zone of the Phosphoria Formation in western Wyoming and southeastern Idaho. US Geol. Surv. Prof. Pap. 1465.

McMURTRY, G.M., VON DER HAAR, D.L., YEH, H.-W., MAHONEY, J.J. & EISENHAUER, A. 1994: Cenozoic accumulation history of a Pacific ferromanganese crust. Earth Planet. Sci. Lett. 125, 105–118.

MENOR, E. 1975: Pétrographie, minéralogie et géochimie des gisements de phosphates de Taïba (Sénégal) et d'Olinda (Brésil). Thèse Doct. Ing. Strasbourg.

MIDDELBURG, J.J. & COMANS, R.N.J. 1991: Sorption of cadmium on hydroxyapatite. Chem. Geol. 90, 45–53.

MILLER, R. L. & SUTCLIFFE, H. 1984: Effects of three phosphate industrial sites on ground-water quality in central Florida, 1979 to 1980. US Geol. Surv. Water Resour. Investi. Report 83-4256, 160–84.

MINSTER, T. & FLEXER, A. 1991: Cadmium-bearing sphalerite in the Senonian section, northern Negev – remarks on its formation and technological implications. In: First Gentner Symp. on Geoscience. Phosphorites and Black Shales. Jerusalem, September 1991, Program with Abstracts, 35.

MORTVEDT, J.J. 1992: The radioactivity issue – effects on crops grown on mined phosphate lands, P-fertilized soils, and phosphogypsum-treated soils. In: Phosphate Fertilizers and the Environment, Proc. Int. Workshop, IFDC, March 1992 Tampa, Florida, 271–278.

NATHAN, Y. 1984: The mineralogy and geochemistry of phosphorites. In: Phosphate Minerals (Ed. by NRIAGU, J.O. & MOORE, P.B.). Springer Verlag, Berlin, 275–291.

NATHAN, Y. & LUCAS, J. 1976: Expériences sur la précipitation directe de l'apatite dans l'eau de mer: implications dans la genèse des phosphorites. Chem. Geol. 18, 181–186.

NATHAN, Y. & NIELSEN, H. 1980: Sulfur isotopes in phosphorites. In: Marine Phosphorites (Ed. by BENTOR, Y.K.). Soc. Econ. Paleont. Miner. Spec. Publ. 29, 73–78.

NATHAN, Y., SHILONI, Y., RODED, R., GAL, I. & DEUTSCH, Y. 1979: The geochemistry of the northern and central Negev phosphorites (southern Israel). Geol. Surv. Israel Bull. 73, 1–41.

NATHAN, Y., SOUDRY, D. & AVIGOUR, A. 1990: Geological significance of carbonate substitution in apatites: Israeli phosphorites as an example. In: Phosphorite Research and Development (Ed. by NOTHOLT, A.J.G. & JARVIS, I.) Geol. Soc. Spec. Publ. 52, 179–191.

NATHAN, Y., SOUDRY, D., DORFMAN, E., LEVY, Y. & SHITRIT, D. 1991: The geochemistry of cadmium in the Negev phosphorites. In: First Gentner Symp. on Geoscience. Phosphorites and Black Shales. Jerusalem, September 1991, Program with Abstracts, 35.

NATHAN, Y., WATKINS, R.T. & BREMNER, J.M. 1993: Rare earth elements from phosphatic-rich sediments off the South African continental shelf. Concepts and Controversies in Phosphogenesis. Int. Symp. and Workshop, Matten, 6–10 September, 1993.

NIFONG, G. D. 1988: Environmental aspects of phosphogypsum. Unpublished Florida Institute of Phosphate Research (FIPR) Manuscript.

NOTHOLT, A.J.G., SHELDON, R.P. & DAVIDSON, D.F. 1989: Phosphate Deposits of the World Volume 2: Phosphate Rock Resources. Cambridge University Press, Cambridge.

O'BRIEN, G.W. & VEEH, H.H. 1980: Holocene phosphorite on the east Australian continental margin. Nature 288, 690–692.

O'BRIEN, G.W., VEEH, H.H., CULLEN, D.J., & MILNES, A.R. 1986: Uranium-series isotopic studies of marine phosphorites and associated sediments from the East Australian continental margin. Earth Planet. Sci. Lett. 80, 19–35.

O'BRIEN, G.W., VEEH, H.H., MILNES, A.R. & CULLEN, D.J. 1987: Sea-floor weathering of phosphorite nodules off East Australia: Its effect on uranium oxidation state and isotopic composition. Geochim. Cosmochim. Acta 51, 2051–2064.

O'BRIEN, G.W., MILNES, A.R., VEEH, H.H., HEGGIE, D.T., RIGGS, S.R., CULLEN, D.J., MARSHALL, J.F. & COOK, P.J. 1990: Sedimentation dynamics and redox iron-cycling: controlling factors for the apatite-glauconite association on the East Australian continental margin. In: Phosphorite Research and Development (Ed. by NOTHOLT, A.J.G. & JARVIS, I.) Geol. Soc. Spec. Publ. 52, 61–86.

PACEY, N.R. 1985: The mineralogy, geochemistry and origin of pellet phosphates in the English Chalk. Chem. Geol. 48, 243–256.

PALMER, M.R. & ELDERFIELD, H. 1985: Sr isotope composition of sea water over the past 75 Myr. Nature 314, 526–528.

PANCZER, G., NATHAN, Y. & SHILONI, Y. 1989: Trace element characterisation of phosphate nodules in Israel. *Sci. Geol. Bull.* 42, 173–184.

PARKER, R.J., 1975: The petrology and origin of some glauconitic and glauco-conglomeratic phosphorites from the South African continental margin. *J. Sediment. Petrol.* 45, 230–242.

PARKER, R.J. & SIESSER, W.G. 1972: Petrology and origin of some phosphorites from the South African continental margin. *J. Sediment. Petrol.* 42, 434–440.

PETERMAN, Z.E., HEDGE, C.E. & TOURTELOT, H.A. 1970: Isotopic composition of strontium in seawater throughout Phanerozoic time. *Geochim. Cosmochim Acta* 34, 105–120.

PIEGRAS, D.J. & WASSERBURG, G.J. 1980: Neodymium isotopic variation in seawater. *Earth Planet. Sci. Lett.* 50, 128–138.

PIEGRAS, D.J., WASSERBURG, G.J. & DASCH, E.J. 1979: The isotopic variations in seawater. *Earth Planet. Sci. Lett.* 45, 223–236.

PIPER, D.Z. 1991: Geochemistry of a Tertiary sedimentary phosphate deposit: Baja California Sur, Mexico. *Chem. Geol.* 92, 283–316.

— 1994: Seawater as a source of minor elements in black shales, phosphorites and other sedimentary rocks. *Chem. Geol.* 114, 95–114.

PIPER, D.Z. & ISAACS, C.M. 1994: Geochemistry of minor elements in the Monterey Formation, CA: seawater chemistry of deposition. In: *Analogy of a Source Rock: The Cooperative Monterey Organic Geochemistry Study* (Ed. by ISAACS, C.M. & RULLKÖTTER, J.). Am. Assoc. Pet. Geol. Stud. Geol. (in press).

PIPER, D.Z. & KOLODNY, Y. 1987: The stable isotopic composition of a phosphate deposit:  $\delta^{13}\text{C}$ ,  $\delta^{34}\text{S}$  and  $\delta^{18}\text{O}$ . *Deep Sea Res.* 34, 897–911.

PIPER, D.Z. & MEDRANO, M.D. 1994: The geochemistry of the Phosphoria Formation at Montpelier Canyon, Idaho: The environment of deposition. *US Geol. Surv. Bull.* 2023B (in press).

PIPER, D.Z., BAEDECKER, P.A., CROCK, J.G., BURNETT, W.C. & LOEBNER, B.J. 1988: Rare earth elements in the phosphatic-enriched sediments of the Peru Shelf. In: *The Origin of Marine Phosphorite. The Results of the R.V. Robert D. Conrad Cruise 23–06 to the Peru Shelf* (Ed. by BURNETT, W.C. & FROELICH, P.N.). *Marine Geol.* 80, 269–285.

POTTS, P.J., HAWKESWORTH, C.J., VAN CALSTEREN, P. & WRIGHT, I.P. 1993a: Advances in analytical technology and its influence on the development of modern inorganic geochemistry: a historical perspective. In: *Magmatic Processes and Plate Tectonics* (Ed. by PRICHARD, H.M., ALABASTER, T., HARRIS N.B.W. & NEARY, C.R.). *Geol. Soc. Spec. Publ.* 76, 501–520.

POTTS, P.J., TINDLE, A.G. & WEBB, P.C. 1993b: *Geochemical Reference Material Compositions*. CRC Press, Boca Raton FL.

PRÉVÔT, L. 1990: Geochemistry, petrography, genesis of Cretaceous – Eocene phosphorites. *Mém. Soc. Géol. Fr.* 158.

PRÉVÔT, L. & LUCAS, J. 1979: Comportement de quelques éléments traces dans les phosphorites. *Sci. Géol. Bull.* 32, 91–105.

— 1980: Behaviour of some trace elements in phosphatic sedimentary formations. In: *Marine Phosphorites* (Ed. by BENTOR, Y.K.). *Soc. Econ. Paleont. Miner. Spec. Publ.* 29, 31–39.

— 1985: Utilization of geochemistry to explain the setting of the phosphatic series of the Ganntour Basin (Morocco). *Sci. Géol. Mém.* 77, 45–51.

— 1986: Microstructure of apatite – replacing carbonate in synthetized and natural samples. *J. Sed. Petrol.* 56, 153–159.

PRÉVÔT, L., EL FALEH, E.M. & LUCAS, J. 1989: Details on synthetic apatites formed through bacterial mediation. *Mineralogy and chemistry of the products*. *Sci. Geol. Bull.* 42, 237–254.

PRICE, N.B. & CALVERT, S.E. 1978: The geochemistry of phosphorites from the Namibian Shelf. *Chem. Geol.* 23, 151–170.

PRICE, N.M. & MOREL, F.M.M. 1990: Cadmium and cobalt substitution for zinc in a marine diatom. *Nature* 344, 658–660.

RICHTER, F.M. & TUREKIAN, K.K. 1993: Simple models for the geochemical response of the ocean to climatic and tectonic forcing. *Earth Planet. Sci. Lett.* 119, 121–131.

ROE, K.K. & BURNETT, W.C. 1985: Uranium geochemistry and dating of Pacific island apatite. *Geochim. Cosmochim. Acta* 49, 1581–1592.

ROE, K.K., BURNETT, W.C., KIM, K.H. & BEERS, M.J. 1982: Excess protactinium in phosphate nodules from the coastal upwelling zone. *Earth Planet. Sci. Lett.* 60, 39–46.

ROESSLER, C. E. 1984: The radiological aspects of phosphogypsum. *Proc. Seminar on Phosphogypsum. Florida Institute of Phosphate Research* 11–36.

- 1988: Radiological assessment of the application of phosphogypsum to agricultural land. *Proc. Second Int. Symp. on Phosphogypsum*. (Ed. by LLOYD, G. M.). Florida Institute of Phosphate Research 5–23.
- ROLLINSON, H. 1993: *Using Geochemical Data: Evaluation, Presentation, Interpretation*. Longman, Burnt Mill, Harlow.
- ROMANKEVICH, Y.A. & BATURIN, G.N. 1972: Composition of the organic matter in phosphorites from the continental margin off southwest Africa. *Geochem. Int.* 9, 464–470.
- RUTTENBERG, K.C. 1993: Reassessment of the oceanic residence time of phosphorus. *Chem. Geol.* 107, 405–409.
- RUTTENBERG, K.C. & BERNER, R.A. 1993: Authigenic apatite formation and burial in sediments from non-upwelling, continental margin environments. *Geochim. Cosmochim. Acta* 57, 991–1007.
- SANDELL, E.B., HEY, M.H. & MCCONNELL, D. 1939: The composition of francolite. *Mineral. Mag.* 25, 395–401.
- SAVENKO, V.S. 1978: Experimental studies on the conditions of chemical precipitation of calcium phosphates from seawater. *Dokl. Akad. Nauk. SSSR* 243, 1302–1305.
- SCHMITZ, B., ABERG, G., WERDELIN, L., FORY, P. & BENDIX-ALMGREEN, S.E. 1991:  $^{87}\text{Sr}/^{86}\text{Sr}$ , Na, F, Sr and La in skeletal fish debris as a measure of the paleosalinity of fossil-fish habitats. *Geol. Soc. Am. Bull.* 103, 786–794.
- SCHULTZ, J. J. 1992: An examination of the environmental issues facing the phosphate fertilizer production sector – indicated cost of environmental compliance. *Phosphate Fertilizers and the Environment*, Tampa, FL, (Ed. by SCHULTZ, J.J.). 283–309.
- SCHUFFERT, J.D., KASTNER, M., EMANUELE, G. & JAHNKE, R.A. 1990: Carbonate-ion substitution in francolite: A new equation. *Geochim. Cosmochim. Acta* 54, 2323–2328.
- SCHWAB, R.G. & OLIVEIRA, N.P. DA 1981: The influence of phosphorus during lateritic weathering. *Zentralbl. Geol. Paläont. Teil 1 3/4*, 419–436.
- SEMELEV, E.I., KHOLODOV, V.N. & BARINSKII, R.L. 1962: Rare earths in phosphorites. *Geochemistry* 5, 501–507.
- SHAW, H.F. & WASSERBURG, G.J. 1985: Sm-Nd in marine carbonates and phosphorites: Implications for Nd isotopes in seawater and crustal ages. *Geochim. Cosmochim. Acta* 49, 503–518.
- SHELDON, R.P. 1964: Paleolatitudinal and paleogeographic distribution of phosphorite. *US Geol. Surv. Prof. Pap.* 501-c, 106–113.
- SHEMESH, A., KOLODNY, Y & LUZ, B. 1983: Oxygen isotope variations in phosphate of biogenic apatites. II. Phosphorite rocks. *Earth Planet. Sci. Lett.* 64, 405–416.
- SHEMESH, A., KOLODNY, Y & LUZ, B. 1988: Isotope geochemistry of oxygen and carbon in phosphate and carbonate of phosphorite francolite. *Geochim. Cosmochim. Acta* 52, 2565–2572.
- SHOLKOVITZ, E.R., PIEGRAS, D.J. & JACOBSEN, S.B. 1989: The pore water chemistry of rare earth elements in Buzzard Bay sediments. *Geochim. Cosmochim. Acta* 53, 2847–2856.
- SLANSKY, M. 1979: Nomenclature et classification des phosphates sédimentaires. *Bull. BRGM*, IV, 2, 153–156.
  - 1980: *Géologie des Phosphates Sédimentaires*. Mém. BRGM 114.
  - 1986: *Geology of Sedimentary Phosphates*. Elsevier, New York.
  - 1989: Nomenclature descriptive des phosphorites. *Sci. Géol. Bull.* 42, 255–266.
- SMALLEY, P.C., HIGGINS, A.C., HOWARTH, R.J., NICHOLSON, H., JONES, C.E., SWINBURNE, N.H.M. & BESSA, J. 1994: Seawater Sr isotope variations through time: a procedure for constructing a reference curve to date and correlate marine sedimentary rocks. *Geology* 22, 431–434.
- SMIRNOV, A.I., IVNITZKAYA, R.B. & ZALAVINA, T.P. 1962: Experimental data on the possibility of the chemical precipitation of phosphate from seawater. In: *Geology of Phosphorite Deposits*. State Sci. Res. Inst. Chem. Raw Mater. 7, 289–302.
- SMITH, J.P. & LEHR, J.R. 1966: An X-ray investigation of carbonate apatite. *J. Agric. Food Chem.* 14, 342–349.
- STARINSKY, A., KATZ, A. & KOLODNY, Y. 1982: The incorporation of uranium into diagenetic phosphorite. *Geochim. Cosmochim. Acta* 46, 1365–1374.
- STAUDIGEL, H., DOYLE, P. & ZINDLER, A. 1985: Sr and Nd isotope systematics in fish teeth. *Earth Planet. Sci. Lett.* 76, 45–56.
- STILLE, P., CHAUDHURI, S., KHARAKA, Y.K. & CLAUER, N. 1992: Neodymium, strontium, oxygen and hydrogen isotope compositions of waters in present and past oceans: a review. In: *Isotopic Signatures and Sedimentary Records, Lecture Notes in Earth Sciences No. 43* (Ed. by CLAUER, N. & CHAUDHURI, S.). Springer-Verlag, Berlin, Heidelberg, 389–410.
- STILLE, P., RIGGS, S., CLAUER, N., AMES, D., CROWSON, R. & SYNDER, S.W. 1994: Sr and Nd isotopic analysis of phosphorite sedimentation through one Miocene high-frequency depositional cycle on the North Carolina continental shelf. *Marine Geol.* 117, 253–273.

SUMMERHAYES, C.P. & MCARTHUR, J.M. 1990: Moroccan offshore phosphorite deposits. In: Phosphate Deposits of the World Volume 3: Genesis of Neogene to Recent Phosphorites. (Ed. by BURNETT, W.C. & RIGGS, S.R.). Cambridge University Press, Cambridge 159–166.

TLIG, S., SASSI, A., BELAYOUNI, H. & MACHEL, D. 1987: Distribution de l'uranium, du thorium, du zirconium, du hafnium et des terres rares (TR) dans les grains de phosphates sédimentaires. *Chem. Geol.* 62, 209–221.

TOOMS, J.S., SUMMERHAYES, C.P. & CRONAN, D. D.S. 1969: Geochemistry of marine phosphate and manganese deposits. *Oceanogr. Marine Biol. Ann. Rev.* 1, 49–100.

TRAVI, Y. 1988: Hydrogéochimie et hydrologie isotopique des aquifères fluorures du bassin du Sénégal – origine et conditions de transport du fluor dans les eaux souterraines. Thèse d'etat, Université de Paris-Sud, France.

TUCKER, M.E. & WRIGHT, V.P. 1990: Carbonate Sedimentology. Blackwell, Oxford.

TUREKIAN, K.K. & WEDEPOHL, K.H. 1961: Distribution of some major elements of the Earth's crust. *Geol. Soc. Am. Bull.* 72, 175–192.

UREY, H.C., LOWENSTAM, H.A., EPSTEIN, S. & MCKINNEY, U. 1951: Measurements of paleotemperatures and temperatures of the Upper Cretaceous of England, Denmark and the south-eastern US. *Geol. Soc. Amer. Bull.* 62, 399–416.

VAN CAPELLEN, P. & BERNER, R. A. 1988: A mathematical model for the early diagenesis of phosphorus and fluorine in marine sediments: apatite precipitation. *Amer. J. Sci.* 288, 289–333.

– 1991: Fluorapatite crystal growth from modified seawater solutions. *Geochim. Cosmochim. Acta* 55, 1219–1234.

VEEH, H.H. 1982: Concordant  $^{230}\text{Th}$  and  $^{231}\text{Pa}$  ages of marine phosphorites. *Earth Planet. Sci. Lett.* 57, 278–284.

VEEH, H.H. & BURNETT, W.C. 1982: Carbonate and phosphate sediments. In : Uranium-Series Disequilibrium: Application to Environmental Problems (Ed. by IVANOVICH, M. & HARMON, R.S.). Clarendon Press, Oxford, 459–480.

VEEH, H.H., BURNETT, W.C. & SOUTAR, A. 1973: Contemporary phosphorites on the continental margin off Peru. *Science* 181, 844–845.

VEEH, H.H., CALVERT, S.E. & PRICE, N.B. 1974: Accumulations of uranium in sediments and phosphorites on the south-west African shelf. *Mar. Chem.* 2, 188–202.

VEIZER, J. 1989: Strontium isotopes in seawater through time. *Ann. Rev. Earth Planet. Sci.* 17, 141–167.

– 1992: Depositional and diagenetic history of limestones: stable and radiogenic isotopes. In: Isotopic Signatures and Sedimentary Records, Lecture Notes in Earth Sciences No. 43 (Ed. by CLAUER, N. & CHAUDHURI, S.). Springer-Verlag, Berlin, Heidelberg, 13–48.

VIGNOLES, M., BONEL, G. & BACQUET, G. 1982: Étude physico-chimique des apatites carbonatées phospho-calciques semblables à la francolite. *Bull. Minéral.* 105, 307–311.

WRIGHT, J., SEYMOUR, R.S. & SHAW, H.F. 1984: REE and Nd isotopes of conodont apatite: variations with geological age and depositional environment. *Geol. Soc. Am. Spec. Pap.* 196, 325–340.

WRIGHT, J., SCHRADER, H. & HOLSER, W.T. 1987: Paleoredox variations in ancient oceans recorded by rare earths in fossil apatite. *Geochim. Cosmochim Acta* 51, 631–644.

ZANIN, YU. N. 1968: Zones of lateritic weathering of secondary phosphorites of Altay-Sayan region. *Int. Geol. Rev.* 10, 1119–1127.

ZANIN, YU. N. & KRIVOPUTSKAYA, L.M. 1976: Transformation of apatite material of phosphorite in catagenesis. *Dokl. Acad. Sci. USSR Earth Sci. Sect.* 230, 200–202.

ZANIN, YU. N., GILINSKAYA, L.G., KRASIL'NIKOVA, N.A., KRIVOPUTSKAYA, L.M., MIRTOV, YU. V. & STOLPOVSKAYA, V.N. 1985: Calcium phosphates in phosphorites of different types. *Int. Geol. Rev.* 27, 1212–1229.

Manuscript received April 4, 1994

Revision accepted July 15, 1994