

Zeitschrift: Eclogae Geologicae Helvetiae
Herausgeber: Schweizerische Geologische Gesellschaft
Band: 88 (1995)
Heft: 3

Artikel: The Prosanto Formation, a marine Middle Triassic Fossil-Lagerstätte near Davos (Canton Graubünden, eastern Swiss Alps)
Autor: Furrer, Heinz
DOI: <https://doi.org/10.5169/seals-167694>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The Prosanto Formation, a marine Middle Triassic Fossil-Lagerstätte near Davos (Canton Graubünden, Eastern Swiss Alps)

HEINZ FURRER¹

Key words: Taphonomy, Fossil-Lagerstätte, Middle Triassic, Prosanto Formation, Graubünden, Eastern Swiss Alps, Austroalpine Silvretta Nappe

Introduction

During the last 15 years, more than 1000 small but well-preserved actinopterygian fishes, a few sauropterygian reptiles and some crustaceans, bivalves, gastropods, dasycladacean algae and terrestrial plants have been collected from the Ladinian (Middle Triassic) Prosanto Formation in the Ducan and Landwasser region near Davos (Canton Graubünden, Eastern Swiss Alps). The sequence of dark limestone, shale and dolomite of the Prosanto Formation occurs as a 100 to 200 m thick and more than 20 km wide lenticular intercalation in shallow water dolomites (Vallatscha and Altein Formations) of the strongly deformed sediments of the Austroalpine Silvretta Nappe. A joint paper on the geology, stratigraphy, paleoecology and the fossils was published by Bürgin et al. (1991). As most of these fossils were collected in the scree, small excavations have been organized in 1992 and 1993 in an alpine locality at 2600 m altitude near Davos to obtain more information about the taphonomy of the fossils and their distribution within the layers. Fossiliferous beds of 2.5 m thickness were studied in detail on a surface of 20 square meters, yielding about 500 complete or fragmentary fossils.

Taphonomy, paleoecology and paleogeography

Articulated skeletons of actinopterygian fishes, aquatic reptiles, crustaceans and twigs of conifers have been preserved only in finely laminated limestone and shale beds without any bioturbation. There are no signs of current activity in these background sediments, such as orientation of skeletons or unidirectional drifting of finer skeletal elements. Some fragmentary fish skeletons and many isolated skulls, jaws, teeth and scales suggest a slow settling of the carcasses through the water column. Slumps, breccias and frequent calcareous turbidites, often rich in siliceous sponge spicules, ostracods and benthic foraminifers, document the instability of the basin margins. A few beds with trace fossils (*Thalassinoides* and *Rhizocorallium*) document short events of bottom population by deposit

¹ Paläontologisches Institut und Museum der Universität Zürich, Karl Schmid-Strasse 4, CH-8006 Zürich

(*Araucarioxylon*) and the terrestrial protorosaur *Macrocnemus*. The rarity of stenohaline organisms, such as cephalopods and echinoderms, indicate a restricted environment. The only connection to the open sea probably existed over areas of the shallow carbonate platform. Intense evaporation possibly led to increased salinity and formation of a pycnocline by concentration of the dense brine in the deeper parts of the restricted basin. Stagnant hypersaline bottom water and depletion of oxygen by decaying organic matter could explain the abiotic environment.

The Prosanto Formation of Graubünden can be compared to the Lower Meride Limestone of the Monte San Giorgio area in Canton Ticino (Southern Switzerland, Furrer 1995, this volume) and the Perledo-Varenna Formation (Northern Italy, Gaetani et al. 1992), both time equivalent restricted basinal facies in the Southern Alps. These formations were deposited in similar intraplateau basins separated by shallow carbonate platforms from the open ocean of the Triassic Tethys.

REFERENCES

- BÜRGIN, T., EICHENBERGER, U., FURRER, H. & TSCHANZ, K. 1991: Die Prosanto-Formation – eine fischreiche Fossil-Lagerstätte in der Mitteltrias der Silvretta-Decke (Kanton Graubünden, Schweiz). *Eclogae geol. Helv.* 84/3, 921–990.
- FURRER, H. 1995: The Kalkschieferzone (Upper Meride Limestone; Ladinian) near Meride (Canton Ticino, Southern Switzerland) and the evolution of a Middle Triassic intraplateau basin. *Eclogae geol. Helv.* 88/3, 827–852.
- GAETANI, M., GNACCOLINI, M., POLIANI, G., GRIGNANI, D., GORZA, M. & MARTELLINI, L. 1992: An Anoxic Intraplateau Basin in the Middle Triassic of Lombardy (Southern Alps, Italy): Anatomy of a Hydrocarbon Source. *Riv. ital. Paleont. Strat.* 97/3–4, 329–354.

