

Zeitschrift: Eclogae Geologicae Helvetiae
Band: 88 (1995)
Heft: 3

Artikel: Doomed-pioneers bioturbation events related to gravity-flow deposits in marine dysaerobic environments
Autor: Föllmi, Karl B. / Grimm, Kurt A.
DOI: <https://doi.org/10.5169/seals-167696>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 15.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Doomed-pioneers bioturbation events related to gravity-flow deposits in marine dysaerobic environments

KARL B. FÖLLMI¹ & KURT A. GRIMM²

Key words: Trace fossils, *Thalassinoides*, sediment-gravity flows, oxygen, doomed pioneers

In a number of localities we have observed discrete horizons with *Thalassinoides* and *Gyrolithes* burrows occurring exclusively at the base of gravity-flow deposits which are intercalated in otherwise non- to poorly bioturbated successions (e.g., Santonian-Maastrich-




Fig. 1. Genesis of doomed-pioneers burrow assemblages: A. Disruption of original habitat of thalassinidean crustacea by, e.g., a storm event. B. Turbulent transport of thalassinidean crustacea within gravity flows. C. Arrival in oxygen-depleted environment and feeding on organic-rich sediments until dysaerobic bottom-water conditions prevent further burrowing. D. Infill of burrows by active backfilling and passive collapse infill. Return to deposition of laminated sediments (from Föllmi & Grimm, 1990; reproduced with permission from the Geological Society of America).

¹ Geological Institute, ETH Centre, CH-8092 Zürich

² Department of Geological Sciences, Univ. of British Columbia, Vancouver, BC V6T 1Z4, Canada

tian Guadalupe Formation, Columbia; Campanian Duwi Formation, Egypt; Campanian Mishash Formation, Israel; Oligocene San Gregorio Formation, Baja California; Miocene Monterey Formation, California). These horizons lack burrows of presumably soft-bodied organisms such as *Zoophycus*, *Planolites*, and *Chondrites*. We attribute a causal relationship between gravity-flow deposition and the presence of *Thalassinoides* and *Gyrolithes* and suggest that these gravity flows have entrained thalassinidean crustacea. Upon deposition in oxygen-deficient environments, the surviving crustacea reworked substantial quantities of laminated, mostly organic-rich sediments in an environment from which they were previously excluded. The persistence of or rapid return to oxygen-depleted conditions limited the survival time and ecological complexity of the imported infaunal dwellers and rendered them doomed pioneers. Ecological and physiological data support this hypothesis: thalassinidean crustacea have the capability to endure turbulent transport and survive upto 5–7 days of anoxia without being severely limited in their biological activities.

For these reasons, isolated horizons with *Thalassinoides* and *Gyrolithes* burrows in otherwise laminated successions are not necessarily indicative of phases of basin-wide oxygenation. Instead, they may represent doomed-pioneer trace-fossil assemblages; i.e., ecologically short-termed phenomena related to the transport of thalassinidean crustacea in gravity flows (Föllmi & Grimm, 1990; Föllmi, 1995; Grimm & Föllmi, 1994).

REFERENCES

- FÖLLMI, K.B. & GRIMM, K.A. 1990: Doomed pioneers: Gravity-flow deposition and bioturbation in marine dysaerobic environments. *Geology*, 18, 1069–1072.
- GRIMM, K.A. & FÖLLMI, K.B. 1994: Doomed pioneers: allochthonous crustacean tracemakers in anaerobic basinal strata, Oligo-Miocene San Gregorio Formation, Baja California Sur, Mexico. *Palaios*, 9, 313–334.
- FÖLLMI, K.B. 1995: The significance of *Thalassinoides* in phosphate-rich beds of the Upper Campanian Duwi Formation, Egypt. *Egypt. J. Geol.* 39/1, 415–430.