

Zeitschrift: Eclogae Geologicae Helvetiae
Herausgeber: Schweizerische Geologische Gesellschaft
Band: 89 (1996)
Heft: 3

Artikel: The larger Foraminifera of Trinidad (West Indies)
Autor: Caudri, Bramine C.M.

Inhaltsverzeichnis

DOI: <https://doi.org/10.5169/seals-167940>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The larger Foraminifera of Trinidad (West Indies)

C.M. BRAMINE CAUDRI¹

Key words: Larger Foraminifera, Trinidad, Early Cretaceous to Miocene, biostratigraphy, taxonomy

ABSTRACT

The present investigations are a follow up to Vaughan & Cole's Preliminary Report on the Cretaceous and Tertiary Larger Foraminifera of Trinidad, British West Indies. They are based on considerable additional faunal material and more advanced stratigraphic information for dating the investigated Cretaceous to Miocene sediments that contain larger foraminifera. Special attention is paid to the Late Eocene transgression in the San Fernando area where it took place in two phases, and to a transgressive interval (Marabella marl) between the Late Eocene and the Oligocene. So far this interval could not be recognized by means of planktic foraminifera but stands out clearly for its larger foraminifera. As such it is correlatable over great distances in the Caribbean region. 24 localities that contain larger foraminifera are described. Their locations are shown on Figure 1. Special attention in the study of the Trinidad larger foraminifera was given to the San Fernando area (Fig. 2–4) which is of particular interest for the abundance of larger foraminifera and their detailed stratigraphic record.

120 taxa were investigated with their ranges shown on distribution charts (Fig. 5–7). They are illustrated on Plates 1–30.

The following taxa are described as new: *Operculina bontourensis*, *O. bontourensis* var. *ornata*, *Operculinoides ocalanus* var. *decoratus*, *O. suteri*, *Heterostegina indicata*, *Lepidocyclina (Polylepidina) nitida*, *L. yurnagunensis* var. *inflata*, *L. sanfernandensis* var. *depressata*, *L. asterocolumnata*, *Eoconuloides senni* var. *conicus*, *Helicolepidinoides intermedius*, *Amphistegina farallonensis*.

ZUSAMMENFASSUNG

Die vorliegenden Untersuchungen sind eine Ergänzung zu Vaughan & Cole's Preliminary Report on the Cretaceous and Tertiary Larger Foraminifera of Trinidad, British West Indies (1941). Sie stützen sich auf bedeutendes zusätzliches Faunenmaterial und fortgeschrittenere stratigraphische Information zur Datierung der untersuchten Kreide- bis Miozänsedimente, die Grossforaminiferen enthalten. Figur 1 zeigt die besprochenen 24 Lokalitäten. Wegen der Häufigkeit von Grossforaminiferen und ihrer stratigraphischen Bedeutung wurde besonders Gewicht auf die obereozäne Transgression im Gebiet von San Fernando gelegt (Fig. 2–4). Diese erfolgte hier in zwei Phasen sowie einem transgressiven Abschnitt (Marabella Mergel) zwischen Obereozän und Oligozän. Dieser konnte bisher mittels planktonischer Foraminiferen nicht festgestellt werden, während er mit Grossforaminiferen klar erkennbar ist. Als solcher kann er in der karibischen Region über grosse Distanzen verfolgt werden.

Die stratigraphische Verbreitung der 120 untersuchten Taxa ist auf den Figuren 5–7 dargestellt. Sie sind auf den Tafeln 1–30 illustriert.

Die folgenden Taxa werden als neu beschrieben: *Operculina bontourensis*, *O. bontourensis* var. *ornata*, *Operculinoides ocalanus* var. *decoratus*, *O. suteri*, *Heterostegina indicata*, *Lepidocyclina (Polylepidina) nitida*, *L. yurnagunensis* var. *inflata*, *L. sanfernandensis* var. *depressata*, *L. asterocolumnata*, *Eoconuloides senni* var. *conicus*, *Helicolepidinoides intermedius*, *Amphistegina farallonensis*.

¹ † 2nd of February 1991

CONTENTS

Note	1139
Introduction	1139
Localities and Biostratigraphy	1140
Description of the larger foraminifera localities	1141
Cretaceous	1143
Paleocene	1143
Eocene and transition Eocene-Oligocene	1145
The San Fernando area	1145
Other localities	1163
Oligo-Miocene	1169
Southern Basin	1169
Oligocene	1169
Miocene	1174
Central Range	1176
Conclusions on the Trinidad larger foraminifera localities	1179
Taxonomy	1181
Family Orbitolinidae	1181
Genus <i>Orbitolina</i> D'ORBIGNY	1181
Family Nummulitidae	1181
Genus <i>Ranikothalia</i> CAUDRI	1181
Genus <i>Operculina</i> D'ORBIGNY	1185
Genus <i>Operculinoides</i> HANZAWA	1187
Genus <i>Nummulites</i> LAMARCK	1192
Genus <i>Heterostegina</i> D'ORBIGNY	1193
Genus <i>Spiroclypeus</i> H. DOUVILLÉ	1193
Group Discocycliniformes	1194
Familiy Discocyclinidae	1199
Genus <i>Athecocyclina</i> VAUGHAN & COLE	1199
Genus <i>Proporocyclina</i> VAUGHAN & COLE	1199
Family Orbitoclypeidae	1200
Genus <i>Hexagonocyclina</i> CAUDRI	1200
Genus <i>Neodiscocyclina</i> CAUDRI	1203
Genus <i>Stenocyclina</i> CAUDRI	1205
Genus <i>Pseudodiphragmina</i> DOUVILLÉ	1206
Genus <i>Asterocyclus</i> GÜMBEL	1206
Genus <i>Actinosiphon</i> VAUGHAN	1210
Genus <i>Lepidocyclus</i> GÜMBEL	1210
Subgenus <i>Polylepidina</i> VAUGHAN	1212
The Lepidocylinas of Grimsdale's (1959) Lineage X	1217
Group of <i>Lepidocyclus yurnagunensis</i>	1220
Group of <i>Lepidocyclus undosa</i>	1223
Group of <i>Lepidocyclus canellei</i>	1227
Genus <i>Econuloides</i> COLE & BERMUDEZ	1229
Genus <i>Helicolepidinoides</i> TAN	1231
Genus <i>Helicolepidina</i> TOBLER	1233
Family Amphisteginidae	1234
Genus <i>Amphistegina</i> D'ORBIGNY	1234
Genus <i>Helicosteginopsis</i> CAUDRI	1235
Genus <i>Helicocyclus</i> TAN	1236
Family Miogypsinidae	1236
Genus <i>Miogypsinia</i> SACCO	1236
Family Planorbulinidae	1237
Genus <i>Planorbulinella</i> CUSHMAN	1237
Family Soritidae	1237

Genus <i>Sorites</i> EHRENBURG	1237
Annotations to Figure 7	1238
Acknowledgments	1239
Index	1240
References	1243
Plates 1–30	1250

NOTE

Bramine Caudri completed her manuscript on the larger foraminifera of Trinidad consisting of 501 typewritten pages, 30 plates and 7 figures and range charts in 1985. Together with the illustrated specimens the manuscript was deposited for eventual publication at the Museum of Natural History, Basel. Because of its size, in particular the systematic part, it proved difficult to fund its publication. It was therefore decided to condense its size without however loosing pertinent data including illustrations and charts. Basically the reductions affected the systematic part. While the newly proposed taxa remain fully documented, synonymy lists, detailed descriptions and discussions of already published forms were curtailed. In numerous instances reductions concerned taxa that were already treated by Caudri in her 1975 Soldado Rock publication. For additional information references to this paper is therefore recommended, and also to the Preliminary Report on the Cretaceous and Tertiary Larger Foraminifera of Trinidad British West Indies by Vaughan & Cole (1941). Together with these two publications Caudri's present contribution substantially complements the documentation on the systematics and the stratigraphic distribution of the larger foraminifera of the main island of Trinidad and nearby Soldado Rock. An original unabridged copy of Caudri's manuscript is deposited in the geological library of the Museum of Natural History, Basel, where it is available for consultation. The manuscript as published here was edited by H.M. Bolli in collaboration with J.P. Beckmann.

1. Introduction

Vaughan & Cole's comprehensive work of 1941 on Trinidad larger foraminifera was presented as «preliminary» because the material at their disposal consisted of rather haphazardly collected samples without a sufficient stratigraphic background. Today, we have more advanced information on the stratigraphy and moreover are now more alert to the important factor of reworking. Consequently we are in a better position to report on the occurrence of the larger foraminifera and their place in the stratigraphic sequence of Trinidad.

Our material consists of a great number of samples collected over many years by geologists of the former Trinidad Leaseholds Ltd. (T.L.L.), later taken over by Texaco Trinidad, Inc. and now nationalized. The collection was supplemented by the Shell subsidiary United British Oilfields of Trinidad (U.B.O.T.), a number of other oil companies and by the Government Geological Survey.

Our collection did not comprise the specimens described by Vaughan & Cole (1941) which are deposited at the U.S. National Museum, Washington D.C. However, from many of the localities mentioned by these authors duplicate material was left in the laboratory at Pointe-a-Pierre, where the greater part of the technical work for this paper was carried out. Most of the examined material is now stored at the Museum of Natural History in Basel where it is catalogued under the numbers C 31049–31259 and C 35911–36130. Also stored there are detailed fossil lists for each locality.