

Zeitschrift: Eclogae Geologicae Helvetiae
Herausgeber: Schweizerische Geologische Gesellschaft
Band: 90 (1997)
Heft: 3

Artikel: Eomyops hebeiseni n.sp., a new large Eomyidae (Rodentia, Mammalia) of the Upper Freshwater Molasse of Switzerland
Autor: Kälin, Daniel
DOI: <https://doi.org/10.5169/seals-168202>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 21.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Eomyops hebeiseni n. sp., a new large Eomyidae (Rodentia, Mammalia) of the Upper Freshwater Molasse of Switzerland

DANIEL KÄLIN¹

Key words: New species, Eomyidae, Rodentia, Mammalia, Middle Miocene

ZUSAMMENFASSUNG

Beschrieben wird eine neue, extrem grosse Art von *Eomyops* aus der Oberen Süsswassermolasse (Mittelmiozän, MN 5/MN 6) der Schweiz. *Eomyops hebeiseni* n. sp. der Fundstelle Chatzloch zeichnet sich gegenüber den drei bisher bekannten Arten *Eomyops catalaunicus* (HARTENBERGER 1966), *Eomyops bodvanus* (JANOSSY 1972) und *Eomyops oppligeri* (ENGESSER 1990) im wesentlichen durch deutlich grössere Dimensionen aus. Morphologisch steht die neue Art *Eomyops catalaunicus* am nächsten.

ABSTRACT

In this paper a new, extremely large species of *Eomyops* is described. *Eomyops hebeiseni* n. sp. of the locality Chatzloch (Upper Freshwater Molasse of Switzerland, Middle Miocene, MN 5/MN 6) is distinctly larger than the three hitherto known species *Eomyops catalaunicus* (HARTENBERGER 1966), *Eomyops bodvanus* (JANOSSY 1972) and *Eomyops oppligeri* (ENGESSER 1990). Morphologically, *Eomyops hebeiseni* n. sp. is closest to *Eomyops catalaunicus*.

Introduction/Historical review

In 1966 Hartenberger described a new European small bunodont eomyid with a lingual anteroloph on the M¹ and M² from the Young Miocene locality Can Llobateres (Spain, MN 9) which he then named *Eomys catalaunicus*.

Later, in 1968 Huguency & Mein described similar teeth from the French localities La Grive and Lissieu (MN 7 + 8 and MN 13) which they incorporated in the North American genus *Leptodontomys* SHOTWELL 1956. They also described another new bunodont eomyid from the fissure filling Vieux Collonges (MN 4 and MN 5, Dép. Rhône, France) as *Eomys? rhodanicus*. This eomyid was also characterized by the possession of a lingual anteroloph on the M¹ and M².

In his discussion of the relationships of the North American and European eomyids Fahlbusch 1973 followed the argumentation of Huguency & Mein and agreed on the morphological affinities between the North American and European forms of *Leptodontomys*.

In 1974 Fejfar described new findings of *Leptodontomys* from the localities Franzensbad (MN 5), Neudorf Spalte 1 and Strakonice (both MN 6). Fejfar (1974: 106) also postulated an

immigration from east for *Leptodontomys*, together with the immigration wave of *Megacricetodon*, *Democricetodon* and *Eumyarion*.

In 1979 Engesser denied the unity of the North American genus *Leptodontomys* with the above mentioned European forms and created for the latter ones the genus *Eomyops*. Since then, various discussions have evolved concerning the generic membership of these small Old World bunodont eomyids.

In 1990 Engesser created in his monograph on eomyids a new genus for the other European bunodont eomyid (*Eomys? rhodanicus*). Because of the morphological differences to the oligocene genus *Eomys* he named the new genus *Pentabuneomys* with the hitherto only known species *P. rhodanicus*.

Recent investigations by Qiu (1994) proved that Chinese *Leptodontomys*-forms are morphologically intermediate between European (*Eomyops*) and North American (*Leptodontomys*) forms. Therefore, Qiu concluded that it is unlikely that the differences between the European and American forms are large enough to justify two different genera.

¹ Naturhistorisches Museum Basel, Augustinergasse 2, CH-4001 Basel

Family Eomyidae DEPÉRET & DOUXAMI 1902

Genus *Eomyops* ENGESSER 1979

Eomyops hebeiseni n. sp.

Fig. 1

Diagnosis

Largest known species of the genus *Eomyops* ENGESSER 1979 with bunodont cheek teeth and an antique *Eomys*-like tooth pattern. It is characterized by the possession of a lingual anteroloph on the M^1 and M^2 . In contrast to *Pentabuneomys*, the upper cheek teeth are with mesoloph and the lower with mesolophid.

Derivatio nominis. – Honouring Mr. Markus Hebeisen, preparator at the Paleontological Institute and Museum of Zurich, Switzerland, for his prospecting of micromammal sites in the Upper Marine and Upper Freshwater Molasse of the Napf alluvial fan, Switzerland. He also discovered the type-locality Chatzloch.

Holotype. – left $M^{1/2}$, Caz 2. Natural History Museum of Basle. Fig. 1.

Dimensions. – 1.20 x 1.39 mm

Type locality. – Chatzloch, Canton of Berne, Switzerland. National grid reference 631'300/203'400/1100 m. Locality-No. 1168–7 in the Register of the Tertiary Mammal-bearing localities of Switzerland at NMB.

Type formation. – Napfschichten, see Kaufmann (1872), Matter (1964), Salis (1967), Habicht (1987).

Stratum type. – 10 to 25 cm dark grey to green-greyish marls with small coal layers. Freshwater molluscs abundant (*Helicidae*, *Clausilia*, *Testacella*).

Referred material of the type locality. – 31 isolated teeth, 8 fragments.

Age. – Middle Miocene, intermediate in age between the Froberg and Rümikon levels in the Biozonation of the Basis of the Upper Freshwater Molasse of Switzerland (Tab. 3, see Kälin 1997, Kempf et al. 1997). In the European MN-Unit system, it corresponds either to the youngest MN 5 or the eldest MN 6

Stratigraphic range. – *Eomyops hebeiseni* is known only from the youngest MN 5 to the eldest MN 6.

Other localities with Eomyops hebeiseni. – Dürrenäsch-Höhenrain (youngest MN 5 with *Megacricetodon lappi*), Uzwil-Nutzenbuech (youngest MN 5 or eldest MN 6).

Differential diagnosis

Eomyops hebeiseni differs from *Eomyops catalaunicus* (HARTENBERGER 1966) in its:

- considerably larger dimensions
- sometimes missing lingual part of the anteroloph on the $M^{1/2}$
- weaker developed 4th syncline on $M_{1/2}$
- sometimes weaker developed anterolophid on $M_{1/2}$

Eomyops hebeiseni differs from *Eomyops bodvanus* (JÁNOSSY 1972) in its:

- considerably larger dimensions
- longer mesolophs on the $M^{1/2}$
- shorter mesolophids on the $M_{1/2}$
- sometimes missing lingual part of the anteroloph on the $M^{1/2}$
- weaker anterolophid on $M_{1/2}$
- non-right-angled junction of posterolophid and hypoconid on the $M_{1/2}$
- commonly backwards directed mesolophids on the $M_{1/2}$
- weaker developed labial part of the anterolophid on the $M_{1/2}$
- weaker developed 4th syncline on $M_{1/2}$.

Eomyops hebeiseni differs from *Eomyops opligeri* ENGESSER 1990 in its:

- considerably larger dimensions
- weaker developed or even missing lingual part of the anteroloph on the $M^{1/2}$
- lack of a bifurcation of the mesoloph(id)s in upper and lower $M^{1/2}$
- commonly backwards directed mesolophids on the $M_{1/2}$
- non-right-angled junction of posterolophid and hypoconid on the $M_{1/2}$
- deeper and narrower exterior syncline on the $M^{1/2}$

Description of the type. – The nearly unworn left $M^{1/2}$ is slightly wider than it is long. It shows a lingual part of the anteroloph that is less developed than the labial part. The longitudinal crest joins the protoconus nearly transversely. In consequence, the lingual syncline is asymmetric and directed slightly forwards. The mesoloph is directed forwards and is short. The exterior syncline is deep and narrow.

Description of the dentition

The P^4 is moderately smaller than $M^{1/2}$ and is wider than it is long. The lingual part is shorter than the labial part. The mesoloph is short and directed slightly forwards. The longitudinal crest is situated more lingually.

$M^{1/2}$ show a nearly quadratic shape or are wider than they are long. Four $M^{1/2}$ show a strong lingual part of the anteroloph, which is weakly developed in three $M^{1/2}$ and completely missing in one $M^{1/2}$ (Fig. 1). On the $M^{1/2}$ with a lingual part of the anteroloph, the lingual part is always less developed than the labial part. The longitudinal crest joins the protoconus nearly transversely. In consequence, the 2nd syncline is longer than the 3rd syncline and the lingual syncline is asymmetric and directed slightly forwards. The mesoloph is short or moderately long and directed forwards. The exterior syncline is narrow.

The M^3 is distinctly smaller than $M^{1/2}$ and shows a strong proto- and paraconus. An anteroloph is present in all four specimens. The mesoloph is moderately long.




Fig. 1. *Eomyops hebeiseni* n. sp. in comparison to a normal-sized representative of the genus *Eomyops*.
 a: *Eomyops hebeiseni* n. sp. from Chatzloch. P⁴-M³. P⁴ dext. (invers), Caz 1, M^{1/2} sin. (Holotyp), Caz 2, M^{1/2} dext. (invers), Caz 3, M³ dext. (invers), Caz 4.
 b: *Eomyops hebeiseni* n. sp. from Chatzloch. P₄-M₂. P₄ sin., Caz 5, M_{1/2} sin., Caz 6, M_{1/2} dext. (invers), Caz 7, M₃ dext. (invers), Caz 8.
 c: *Eomyops* aff. *opligeri* from Nebelbergweg. P²-M³. P⁴ sin., Nbw 41, M¹ dext. (invers), Nbw 42, M² dext. (invers), Nbw 43, M³ sin., Nbw 44.
 d: *Eomyops* aff. *opligeri* from Nebelbergweg. P₄-M₃. P₄ sin., Nbw 48, M₁ dext. (invers), Nbw 49, M₂ dext. (invers), Nbw 50.
 all figures 25x




Fig. 2. Scatter diagramme of the molars and premolars (P4, M1/2) of *Eomyops catalaunicus* (HARTENBERGER 1966), *Eomyops bodvanus* (JÁNOSSY 1972), *Eomyops oppligeri* ENGESSER 1990, and *Eomyops hebeiseni* n. sp. from different localities. Values after citations in Table 2 and Kälin (1993).

Tab. 1. Key features for the morphological characterization of the four hitherto described species of *Eomyops*.

	<i>Eomyops catalaunicus</i> (HARTENBERGER 1966)	<i>Eomyops bodvanus</i> (JÁNOSSY 1972)	<i>Eomyops oppligeri</i> ENGESSER 1990	<i>Eomyops hebeiseni</i>
lingual Anteroloph	present	present	present, pronounced	present
Mesoloph	moderately long always single	short always single	moderately long or short sometimes split	moderately long always single
direction of Mesoloph	forwards	forwards	forwards	forwards
exterior Syncline	deep and narrow	deep and narrow	wide	deep and narrow
Mesolophid	moderately long	long	moderately long	moderately long
direction of Mesolophid	backwards or transversal	transversal	backwards or transversal	backwards
Anterolophid	present, strong	present, strong	present	present
junction of Posterolophid	acute-angled	right-angled	right-angled	acute-angled
4th Syncline	strong	strong	weak	weak

The P₄ is longer than it is wide. One of the two specimens shows a small residue of an anterolophid. The mesolophid is short in one specimen and missing in the other one. The posterolophid is short, but strong and joins the hypoconid directly.

The M_{1/2} are longer than they are wide. They show a strong and well developed anterolophid. In six of nine specimens the anterolophid is connected with the metalophid. The mesolophid is usually short, though rarely moderately long. The mesolophid is directed backwards in five of nine specimens and transversal in four specimens. The posterolophid does not join the hypolophid at a right angle.

The M₃ are relatively large and slightly elongated. The anteroloph is lingually and labially well developed, the 4th syncline is strongly reduced. The mesolophid is moderately long and directed transversal (3) or backwards (1).

An important diagnostic feature of *Eomyops* is the crenulated lower incisor. Unfortunately, among the few lower incisors of the locality Chatzloch there is not one with a crenula-

tion which could be attributed to *Eomyops* as described in Figure 8 in Engesser (1979).

Discussion

The present discovery of a new large species of *Eomyops* of MN 5/MN 6 age does not lessen the above-mentioned quandary. Morphologically, *Eomyops hebeiseni* n. sp. is closest to *Eomyops catalaunicus* (Tab. 1). But *Eomyops catalaunicus* is much smaller and very typical for localities of younger Miocene age (Can Llobateres, Montredon, Nebelbergweg, Petersbuch 14; see Tab. 2). All other contemporaneous findings of MN 5-age (Massendorf, Franzensbad) are considerably smaller and morphologically closer to *Eomyops oppligeri*. Neither can *Eomyops hebeiseni* be incorporated into the genus *Pentabuneomys*. Indeed, there is a size increase within the evolutionary lineage of *Pentabuneomys rhodanicus* (hitherto unpublished material of the Swiss Molasse and also Engesser

Tab. 2. List of sites yielding *Eomyops* in Western and Central Europe (excluding Swiss localities).

Locality	Age	Literature	
Schernfeld	MN 17	Dehm 1962 Fahlbusch 1973	<i>Eomyops</i> sp.
Ivanovce	MN 15	Fejfar 1989	<i>Eomyops</i> sp.
Podlesice	MN 14	Fahlbusch 1978 Kowalski 1989	<i>Eomyops</i> cf. <i>catalaunicus</i>
Osztramos 1	MN 14	Jánossy 1972	<i>Eomyops bodvanus</i> type locality
Osztramos 9	MN 14	Jánossy 1975	<i>Eomyops</i> cf. <i>bodvanus</i>
Osztramos 10	?	Jánossy & Kordos 1977	<i>Eomyops</i> cf. <i>bodvanus</i>
Lissieu	MN 13	Hugueney & Mein 1968	<i>Eomyops</i> aff. <i>catalaunicus</i>
Ambérieu 3	MN 11	Mein 1984	<i>Eomyops</i> sp.
Dorn-Dürkheim	MN 11	Franzen 1981	<i>Eomyops</i> sp.
Suchomasty	MN 10	Fejfar 1989	<i>Eomyops</i> cf. <i>catalaunicus</i>
Ambérieu 1+2	MN 10	Farjanel & Mein 1984	<i>Eomyops catalaunicus</i>
Kohfidisch	MN 10	Bachmayer & Wilson 1980	<i>Eomyops</i> sp.
Richardhof	MN 10	Daxner-Höck 1996	<i>Eomyops</i> sp.
Montredon	MN 10	Aguilar 1982	<i>Eomyops catalaunicus</i>
Soblay	MN 10	Guérin & Mein 1971	<i>Eomyops catalaunicus</i>
Can Llobateres	MN 9	Hartenberger 1966	<i>Eomyops catalaunicus</i> type locality
Jujurieux	MN 9	Mein 1985	<i>Eomyops</i> sp.
Rudabanya	MN 9	Rabeder 1985	<i>Eomyops catalaunicus</i>
Götzendorf	MN 9	Rögl et al. 1993	<i>Eomyopscatalaunicus</i>
Stixneusiedl	MN 9	Rögl et al. 1993	<i>Eomyops</i> sp.
Priay II	MN 9	Welcomme et al. 1991	<i>Eomyops catalaunicus</i>
Hammerschmiede	MN 9	Fahlbusch 1975	<i>Eomyops</i> sp.
Markt	MN 9	Fahlbusch 1973	<i>Eomyops</i> sp.
Petersbuch 14	MN 9	Bolliger & Rummel 1994	<i>Eomyops catalaunicus</i>
Petersbuch 10	MN 8/MN 9	Bolliger & Rummel 1994	<i>Eomyops</i> cf. <i>oppligeri</i>
Giggenhausen	MN 8	Fahlbusch 1973 Fahlbusch 1975	<i>Eomyops oppligeri</i>
Felsőtárkány	MN 7/8	Kretzoi 1982	<i>Eomyops</i> sp.
La Grive	MN 7/8	Hugueney & Mein 1968	<i>Eomyops oppligeri</i>
Las Planas SK	MN 6	Daams & Freudenthal 1988	<i>Eomyops</i> sp.
Manchones	MN 6	Bruijn 1967 Fahlbusch 1973	? <i>Eomyops oppligeri</i>
Arroyo del Val	MN 6	Fahlbusch 1973 Fahlbusch 1975	? <i>Eomyops oppligeri</i>
Neudorf	MN 6	Fejfar 1974	? <i>Eomyops oppligeri</i>
Massendorf	MN 5	Schötz 1979	<i>Eomyops</i> sp.
Strakonice	MN 5	Fejfar 1974	<i>Eomyops</i> sp.
Franzensbad	MN 5	Fejfar 1974	<i>Eomyops</i> sp.

Tab. 3. Distribution of findings of *Eomyops* and *Pentabuneomys* in the Swiss Molasse Basin.

	International Reference Faunas (Mein 1989)	MN-Unit	Swiss Reference Fauna	Swiss localities with <i>Pentabuneomys</i> or <i>Eomyops</i>	
Upper Freshwater Molasse	Can Lobateres	MN 9	unnamed	Nebelbergweg	
	La Grive M	Anwil	MN 8	Anwil	Anwil Ottenberg 3
		Steinheim	MN 7	unnamed	
	Sansan	MN 6	unnamed		
			Rümikon	Uzwil-Nutzenbuech Chatzloch Dürrenäsch-Höhenrain	
	Faluns Pont-Levoy	MN 5	Frohberg		
			Tobel Hombrechtikon		
			Vermes 1		
	Upper Marine Molasse	La Romieu	MN 4	Tägeraustasse	
		Wintershof-West	MN 3b	Trub-Sältenbach	
Goldinger Tobel 8				Hasenbach 1 Waltwil-Dorfbachtobel 1, 2, 5 Goldinger Tobel 8 Goldinger Tobel 2	
MN 3a			Bierkeller	Muhen-Rütisgraben 1+2 Bierkeller	
Lower Freshwater Molasse		Laugnac	MN 2b	Vully 1	
			MN 2a	La Mèbre 698	
				La Chauz 7	
			Les Berglières		

Eomyops

Pentabuneomys

1990: 118) but the genus *Pentabuneomys* is characterized by the possession of a mesoconus, respectively a mesoconid on M1/2 and has hitherto never been found in localities younger than MN 4¹. Further, in the Swiss Molasse basin *Pentabuneomys* is frequent in the assemblage zones of Bierkeller and Goldinger Tobel 8 (Tab. 3, younger MN 3a and older MN 3b, Kälin 1997) and has not been recorded in younger layers.

¹ Except *Pentabuneomys rhodanicus* of Vieux Collonges which is however a fissure filling of both MN 4 and MN 5 age.

Hünemann (1981) described an isolated P₄ from the locality Rodenberg (probably MN 5) as *Pentabuneomys rhodanicus*, but the homogeneity of this locality is uncertain until a possible confirmatory reworking is carried out.

So the phylogenetic relationship of *Eomyops hebeiseni* to the three other known species of *Eomyops* is not clear. For the moment, *Eomyops hebeiseni* has to be regarded as an isolated form of *Eomyops* with hitherto unknown ancestor and descendant.

There is thus no further advance on the question of whether or not *Eomyops* and *Leptodontomys* are independent genera as originally suggested by Engesser (1979). Until this problem is resolved, we use the nomen *Eomyops* for middle and late Miocene European eomyids with an *Eomys*-like tooth pattern as proposed by Fahlbusch & Bolliger (1996).

Generally *Eomyops* seems to be a rare eomyid. There are only a few faunas that have yielded rare teeth of *Eomyops*. Teeth of *Eomyops* have been recorded from the following localities of the Swiss Molasse:

Vue-des-Alpes	MN 15	Bolliger et al. 1993 <i>Eomyops</i> cf. <i>bodvanus</i>
Nebelbergweg	MN 9	Kälin 1993 <i>Eomyops</i> aff. <i>catalaunicus</i> + <i>Eomyops</i> aff. <i>oppligeri</i>
Anwil	MN 8	Engesser 1972 <i>Eomyops oppligeri</i> , type locality
Ottenberg 3	MN 8	Bolliger 1996 <i>Eomyops oppligeri</i>
Uzwil-Nutzenbuech	MN 5/MN 6	<i>Eomyops hebeiseni</i> n. sp.
Chatzloch	MN 5/MN 6	this paper <i>Eomyops hebeiseni</i> n. sp., type locality
Dürrenäsch-Höhenrain	MN 5	<i>Eomyops hebeiseni</i> n. sp.

Localities outside the Swiss Molasse are listed in Table 2.

In the Chatzloch fauna, *Eomyops hebeiseni* n. sp. is an abundant element (9%). The faunal list of Chatzloch is:

	n =	% =
<i>Amphiperatherium</i> sp.	1	0.3
<i>Galerix</i> sp.	4	1
<i>Dinosorex</i> cf. <i>zapfei</i>	7	2
Soricidae indet.	3	0.9
<i>Plesiodimylus</i> sp.	11	3
<i>Lanthanotherium</i> sp.	6	2
Chiroptera indet.	2	0.5
<i>Megacricetodon</i> cf. <i>minor</i>	23	6.5
<i>Democricetodon</i> aff. <i>gracilis</i>	33	9.4
<i>Eumyarion</i> sp.	83	23.6
<i>Lartetomys</i> sp.	3	0.9
<i>Anomalomys</i> aff. <i>gaudryi</i>	5	1.4

Tab. 4. Measurements of *Eomyops hebeiseni* n. sp. in mm.
M1sL = Length of an upper first molar, M2iB = Width of a lower second molar.
P = premolar, D = milk-tooth

	D4sL	D4sB	P4sL	P4sB	M1/2sL	M1/2sB	M3sL	M3sB	D4iL	D4iB	P4iL	P4iB	M1/2iL	M1/2iB	M3iL	M3iB
1	1.04	1.26	1.10	1.27	1.20	1.39	1.00	1.23	1.18	0.84	1.13	1.06	1.28	1.31	1.28	1.24
2			1.06	1.20	1.17	1.42	1.00	1.23			1.24	1.04	1.25	1.26	1.20	1.28
3					1.12	1.38	0.88	1.10					1.22	1.31	1.20	1.26
4					1.22	1.42	0.96	1.18					1.16	1.33	1.08	1.14
5					1.12	1.34							1.23	1.16		
6					1.24	1.44							1.30	1.30		
7					1.12	1.42							1.32	1.26		
8					1.12	1.36							1.28	1.26		
9													1.20	1.28		
Ø	1.04	1.26	1.08	1.24	1.16	1.40	0.96	1.19	1.18	0.84	1.19	1.05	1.25	1.27	1.19	1.23

<i>Eomyops hebeiseni</i> n. sp.	31	8.8
<i>Keramidomys carpathicus</i>	65	18.5
<i>Spermophilinus aff. bredai</i>	7	2
<i>Miopetaurista</i> sp.	5	1.4
<i>Palaeosciurus</i> sp.	7	2
<i>Blackia miocaenica</i>	1	0.3
<i>Paraglitulus werenfelsi</i>	30	8.5
<i>Miodromys aegercii</i>	5	1.4
<i>Microdromys koenigswaldi</i>	9	2.6
<i>Amphilagus</i> sp.	3	0.9
<i>Prolagus oeningensis</i>	4	1.1
Suidae indet.	1	0.3
Cervidae indet.	1	0.3
Carnivora indet.	1	0.3
	$\Sigma = 351$	

The age of the Chatzloch fauna can be easily calibrated. In the Swiss Molasse biozonation, it is intermediate in age between the Frohberg and Rümikon levels (Tab. 3, Kálin 1997). This calibration also correlates well with the litho- and magnetostratigraphy (Kempf et al. 1997). In the European MN-Unit system, it corresponds either to the youngest MN 5 or the eldest MN 6. In addition to this, the evolutionary level of selected taxa shows more similarities with the reference locality of MN 5 (Faluns Pont Levoy – Thenay) than with Sansan, the reference locality of MN 6.

Furthermore, there are strong similarities with the Slovakian Devínska Nová Ves (Neudorf) locality (Schaub & Zapfe 1953).

One of the two other localities containing *Eomyops hebeiseni* n. sp. (Dürrenäsch-Höhenrain) yielded very large specimens of *Megacricetodon lappi* and certainly belongs to the youngest MN 5, whereas the other locality (Uzwil-Nutzenbuech) is, in its composition, almost identical with the Chatzloch locality although slightly younger. Additionally, the locality Uzwil-Nutzenbuech is probably situated slightly above the “Blockhorizont” of the Upper Freshwater Molasse of Eastern Switzerland, a marker bed which is correlated with the Ries impact of Southern Germany (Hofmann 1978). The age of the Ries event is dated at 15.1 Ma (Staudacher et al. 1982) or 14.6 ± 0.6 Ma (Gentner & Wagner 1969).

It can therefore be concluded that *Eomyops hebeiseni* n. sp. seems to be restricted to a very short time interval around the MN 5/MN 6 – “boundary”.

Acknowledgments

This work was supported by the Swiss National Science Foundation (project No. 20-43150.95). I am indebted to many persons for their help. M. Hebeisen (Zurich) provided me with information on vertebrate localities in the Napf alluvial fan. Dr. B. Engesser (Basle) contributed stimulating discussions. Dr. J.H. Russel (Zollikon) is kindly acknowledged for a critical review of the english manuscript; and also the reviewers Drs. O. Fejfar and T. Bolliger. U. Leuthold, D. Oppliger and M. Weick (Basle) supported me in the field and H. Gredig (Basle) washed and picked out large quantities of sediments. SEM photographs were made by U. Jauch (Univ. of Zurich).

REFERENCES

- AGUILAR, J.-P. 1982: Contributions à l'étude des micromammifères du gisement miocène supérieur de Montredon (Hérault). 2 – Les rongeurs. *Palaeovertebrata* 12 (3), 81–117.
- BACHMAYER, F. & WILSON, R.W. 1980: A Third Contribution to the Fossil Small Mammal Fauna of Kohfidisch (Burgenland), Austria. *Ann. Naturhist. Mus. Wien* 83, 351–386.
- BOLLIGER, T. 1996: Drei neue Kleinsäugerfaunen aus der miozänen Molasse der Ostschweiz (MN 3 und MN 7/8). *N. Jb. Geol. Paläont. Abh.* 202, 95–110.
- BOLLIGER, T., ENGESSER, B. & WEIDMANN, M. 1993: Première découverte de mammifères pliocènes dans le Jura neuchâtelois. *Eclogae geol. Helv.* 86/3, 1031–1068.
- BOLLIGER, T. & RUMMEL, M. 1994: Säugetierfunde aus Karstspalten – Die komplexe Genese am Beispiel eines Steinbruches bei Petersbuch, Südliche Frankenalb (Bayern). *Mitt. bayer. Staatsslg. Paläont. hist. Geol.* 34, 239–264.
- BRUJN, H. DE. 1967: Gliridae, Sciuridae y Eomyidae (Rodentia, Mammalia) miocenos de Calatayud (provincia de Zaragoza, España) y su relación con la biostratigraphía del area. *Bol. Inst. Geol., Min., España* 78, 187–373.
- DAAMS, R. & FREUDENTHAL, M. 1988: Synopsis of the Dutch-Spanish collaboration program in the Aragonian type area, 1975–1986. *Scripta Geologica, Spec. Issue* 1, 3–18.
- DAXNER-HÖCK, G. 1996: Faunenwandel im Obermiozän und Korrelation der MN-“Zonen” mit den Biozonen des Pannons der Zentralen Paratethys. *Beitr. Paläont.* 21/1, 1–9.
- DEHM, R. 1962: Altpleistozäne Säuger von Schernfeld bei Eichstätt in Bayern. *Mitt. bayer. Staatsslg. Paläont. hist. Geol.* 2, 17–61.
- ENGESSER, B. 1972: Die obermiozäne Säugetierfauna von Anwil (Baselland). *Tätigkeitsber. Natf. Ges. Basell.* 28, 37–363.

- 1979: Relationships of some insectivores and rodents from the Miocene of North America and Europe. *Bull. Carnegie Mus. nat. Hist.* 14, 1–68.
- 1990: Die Eomyidae (Rodentia, Mammalia) der Molasse der Schweiz und Savoyens. *Schweiz. Paläont. Abh.* 112, 1–144.
- FAHLBUSCH, V. 1973: Die stammesgeschichtlichen Beziehungen zwischen den Eomyiden (Mammalia, Rodentia) Nordamerikas und Europas. *Mitt. Bayer. Staatslg. Paläont. hist. Geol.* 13, 141–175.
- 1975: Die Eomyiden (Rodentia, Mammalia) der Oberen Süßwasser-Molasse Bayerns. *Mitt. Bayer. Staatslg. Paläontol. hist. Geol.* 15, 63–90.
- 1978: Pliozäne und Pleistozäne Eomyidae (Rodentia, Mammalia) aus Polen. *Acta Zoologica Cracoviensia*, Tome 23, No. 2, 13–27.
- FAHLBUSCH, V. & BOLLIGER, T. 1996: Eomyids and Zapodids (Rodentia, Mammalia) in the Middle and Upper Miocene of Central and Southeastern Europe and the Eastern Mediterranean. In: BERNOR, R.L., FAHLBUSCH, V. & MITTMANN, H.-W. (eds): *The evolution of western Eurasian Neogene mammal faunas*. Columbia University Press, 208–212.
- FARJANEL, G. & MEIN, P. 1984: Une association de mammifères et de pollens dans la formation continentale des “Marnes de Bresse” d’âge Miocène supérieure, à Ambérieu (Ain). *Géologie de la France* 1984, 1–2, 131–148.
- FEJFAR, O. 1974: Die Eomyiden und Cricetiden (Rodentia, Mammalia) des Miozäns der Tschechoslowakei. *Paläontographica A* 146, 100–180.
- 1989: The Neogene VP Sites of Czechoslovakia: A Contribution to the Neogene Terrestrial Biostratigraphy of Europe based on Rodents. In: LINDSAY, E.H. et al. (eds): *European Neogene Mammal Chronology*. NATO ASI Ser. A. 180 (Plenum Press New York), 211–236.
- FRANZEN, J.L. 1981: Eine turoloische (obermiozäne) Waldfauna aus Rheinhessen. *Alzeier Geschichtsblätter* 16, 192–202.
- GENTNER, W. & WAGNER, G.A. 1969: Altersbestimmungen an Riesgläsern und Moldaviten. *Geologica Bavarica* 61, 296–303.
- GUERIN, C. & MEIN, P. 1971: Les principaux gisements de mammifères miocènes et pliocènes du domaine rhodanien. *Docum. Lab. Geol. Univ. Lyon, H.S.* 131–170.
- HABICHT, J.K.A. 1987: Schweizerisches Mittelland (Molasse). *Lex. stratigr. int. Europe* 1/7b, 1–528.
- HARTENBERGER, J.-L. 1966: Les rongeurs du Vallésien (Miocène supérieur) de Can Llobateres (Sabadell, Espagne): Gliridae et Eomyidae. *Bull. Soc. géol. France* 8, 596–604.
- HOFMANN, F. 1978: Spuren eines Meteoriteneinschlages in der Molasse der Ostschweiz und deren Beziehungen zum Riesereignis. *Bull. Ver. Schweiz. Petroleum-Geol. u. Ing.* 44/107, 17–27.
- HUGUENEY, M. & MEIN, P. 1968: Les Eomyidés (Mammalia, Rodentia) néogènes de la Région lyonnaise. *Geobios* 1, 187–204.
- HUNERMANN, K.A. 1981: Die Glimmersandgrube am Rodenberg bei Schlatingen (Kt. Thurgau) als paläontologisches Studienobjekt in der Oberen Süßwassermolasse. *Mitt. thurg. naturf. Ges.* 44, 7–32.
- JANOSSY, D. 1972: Middle Pliocene Microvertebrate Fauna from the Osztramos Loc. 1 (Northern Hungary). *Ann. Hist. Natur. Mus Nat. Hungar.* 64, 27–52.
- 1975: New “Middle Pliocene” Microvertebrata fauna from Northern Hungary (Osztramos Loc. 9. and 13.). *Trabajos sobre Neogeno-Cuaternario* 4, 93–97. Madrid.
- JANOSSY, D. & KORDOS, L. 1977: Az Osztramos gerinces lelöhelyeinek faunisztikai és karsztmorfológiai áttekintése (1975-ig). *Fragm. Min. et Pal.* 8, 39–72.
- KALIN, D. 1993: Stratigraphie und Säugetierfaunen der Oberen Süßwassermolasse der Nordwestschweiz. *Ph. D. Thesis ETH Zürich No.* 10'152, 1–238.
- 1997 (in press): The Mammal Zonation of the Upper Marine Molasse of Switzerland reconsidered – a local Biozonation of MN 2–MN 5. In: AGUILAR, J.-P., LEGENDRE, S. & MICHAUX, J. (Eds.), *Actes du Congrès Biochrom'97. Mémoires et Travaux de l'E.P.H.E., Institut de Montpellier*, 21.
- KAUFMANN, F.J. 1872: Rigi- und Molassegebiet der Mittelschweiz. *Beitr. geol. Karte Schweiz* 11, 534 p.
- KEMPF, O., BOLLIGER, T., KALIN, D., ENGESSER, B. & MATTER, A. 1997 (in press): Revised magnetostratigraphic calibration of Oligocene to Miocene mammal biozones of the North Alpine foreland basin. In: AGUILAR, J.-P., LEGENDRE, S. & MICHAUX, J. (Eds.), *Actes du Congrès Biochrom'97. Mémoires et Travaux de l'E.P.H.E., Institut de Montpellier*, 21.
- KOWALSKI, K. 1989: Stratigraphy of Neogene Mammals of Poland. In: LINDSAY, E.H. et al. (eds): *European Neogene Mammal Chronology*. NATO ASI Ser. A. 180 (Plenum Press New York), 193–209.
- KRETZOI, M. 1982: Wichtigere Streufunde aus der wirbeltierpaläontologischen Sammlung der Ungarischen Geologischen Anstalt. 7. Mitteilung. *Földt. Int. Évi Jel.* 1980–röf., 385–394. Budapest.
- MATTER, A. 1964: Sedimentologische Untersuchungen im östlichen Napfgebiet (Entlebuch – Tal der grossen Fontanne, Kt. Luzern). *Eclogae geol. Helv.* 57/2, 315–428.
- MEIN, P. 1984: Composition quantitative des faunes de mammifères du Miocène moyen et supérieur de la région Lyonnaise. *Paléobiol. continent.* 14, 339–346.
- 1985: A new direct correlation between marine and continental scales in rhodanian Miocene. *Abstracts VIIIth Congress of the R.C.M.N.S. Budapest*, 377–379.
- 1989: Updating of MN Zones. – In: LINDSAY, E.H. et al. (eds): *European Neogene Mammal Chronology*. NATO ASI Ser. A. 180 (Plenum Press New York): 73–90.
- QIU, Z. 1994: Eomyidae in China. *Nat. Sc. Mus. Tokyo Monogr.* 8, 49–55.
- RABEDER, G. 1985: Die Säugetiere des Pannonien. In: PAPP, A., JAMBOR, A. & STEININGER, F.F. (eds): *Chronostratigraphie und Neostratotypen. Miozän der zentralen Paratethys, M6 Pannonien*. Akademia Kiado, Budapest, 440–463.
- RÖGL, F., ZAPFE, H., BERNOR, R.L., BRZOBHATY, R.L., DAXNER-HÖCK, G., DRAXLER, I., FEJFAR, O., HERRMANN, P., RABEDER, G., SCHULTZ, O. & ZETTER, R. 1993: Die Primatenfundstelle Götzendorf an der Leitha (Obermiozän des Wiener Becken, Niederösterreich). *Jb. Geol. B.-A.* 136/2, 503–526.
- SALIS, K. VON 1967: Geologische und sedimentologische Untersuchungen in Molasse und Quartär südlich Wolhusen (Entlebuch, Kt. Luzern). *Mitt. Natf. Ges. Luzern* 21, 1–107.
- SCHAUB, S. & ZAPFE, H. 1953: Die Fauna der mittelmiozänen Spaltenfüllung von Neudorf an der March (CSSR). *Simplicidentata. Sitzber. österr. Akad. Wiss., math.-natwiss. Kl. Abt. 1.* 162, 181–215.
- SCHÖTZ, M. 1979: Neue Funde von Eomyiden aus dem Jungtertiär Niederbayerns. *Aufschluss* 30, 465–473.
- STAUDACHER, T., JESSBERGER, E.K., DOMINIK, B., KIRSTEN, T. & SCHAEFFER, O.A. 1982: 40Ar–39Ar Ages of Rocks and Glasses from the Nördlinger Ries Krater and the Temperature History of Impact Breccias. *Journ. Geophys.* 51, 1–11.
- WELCOME, J.-L., AGUILAR, J.-P. & GINSBURG, L. 1991: Découverte d’un nouveau Pliopithèque (Primates, Mammalia) associé à des rongeurs dans les sables du Miocène supérieur de Priay (Ain, France) et remarques sur la paléogéographie de la Bresse au Vallésien. *C.R. Acad. Sci. Paris*, t. 313/2, 723–729.

Manuscript received May 6, 1997

Revision accepted September 25, 1997

