

**Zeitschrift:** Eclogae Geologicae Helvetiae  
**Herausgeber:** Schweizerische Geologische Gesellschaft  
**Band:** 93 (2000)  
**Heft:** 3

**Artikel:** A Holocene fossil coleopteran (beetle) assemblage from the Central Swiss Alps : climate and ecology  
**Autor:** Jost-Stauffer, Monika  
**DOI:** <https://doi.org/10.5169/seals-168835>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 25.11.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# A Holocene fossil coleopteran (beetle) assemblage from the Central Swiss Alps: Climate and ecology

MONIKA JOST-STAUFFER

*Key words:* Fossil beetles, fen peat, palaeoenvironment, timberline, Central Swiss Alps, Holocene

## ABSTRACT

The beetle fauna of a fen peat melted out in pieces from below the retreating Unteraar Glacier was examined. The sediment (12 kg of peat) yielded more than 5000 insect fragments. Identification of the fossils provided a detailed picture about palaeoenvironment in the Gletschervorfeld 3600–3800 yr BP ago. Coleopterans living in marshy alpine areas near lakes and running water associated with reeds and moss dominate the assemblage. Species indicating persisting snow patches even during summer months are *Helophorus glacialis* and *Nebria castanea*. Many of the specimens reported live in flowering meadows and shrubs (often *Rhododendron*, *Alnus*, *Salix*). There are several species in the assemblage which are not known to live in the alpine zone (e.g. *Stenus bifoveolatus*, *Gymnusa brevicollis*). These observations and the total absence of tree-dependent species suggest that the ancient Cyperaceae fen peat was growing in the subalpine dwarf-shrub heath - above the forest limit but below the treeline. From the finding of wood fragments in the same deposits, the presence of isolated trees can be inferred, probably on the slopes where the conditions were locally more favourable. Reconstructed July temperatures based on the beetle assemblage yielded a mean value similar to today's temperature measured 5.5 km east from the site at Grimsel Hospiz.

## ZUSAMMENFASSUNG

Im Zusammenhang mit dem aktuellen Gletscherrückzug wurde die Käferfauna eines Flachmooses untersucht, das in Stücken aus dem Unteraargletscher geschmolzen wurde. Die Untersuchung von 12 kg Torf lieferte mehr als 5000 Insektenreste. Anhand der Fossilien war es möglich, ein detailliertes Bild der Umgebung des Gletschervorfeldes vor 3600–3800 yr BP zu erhalten. Die Fauna wird dominiert von Arten, die sumpfiges Habitat bevorzugen, in der Nähe von Seen und fließenden Gewässern leben und mit Seggen und Moos assoziiert sind. Zwei Spezies (*Helophorus glacialis* und *Nebria castanea*) weisen auf die Anwesenheit von Schneeflecken, auch während der Sommermonate, hin. Viele der bestimmten Arten leben auf blühenden Wiesen und Büschen (bevorzugt *Rhododendron*, *Alnus* und *Salix*). Es gibt mehrere nicht-alpine Arten in der Faunenvergesellschaftung (z.B. *Stenus bifoveolatus*, *Gymnusa brevicollis*). Diese Beobachtungen sowie die vollständige Abwesenheit von Coleopterenarten, die von Bäumen abhängig sind, führen zum Schluss, dass das Cyperaceen-Flachmoor zur Zeit seiner Entstehung in der subalpinen Zwergstrauchstufe lag – oberhalb der Wald-, aber unterhalb der Baumgrenze. Einzelne Holzfragmente in den torfigen Ablagerungen lassen auf die Existenz einzelner Bäume in der Nähe schließen. Wahrscheinlich wuchsen diese an den Hängen, ähnlich der heutigen Situation, wo das Klima lokal vorteilhafter ist. Die Käferfauna wurde zur Rekonstruktion einer mittleren Julitemperatur verwendet. Sie ergab einen ähnlichen Wert dem heutigen, wie er 5.5 km weiter östlich am Talausgang beim Grimsel Hospiz gemessen wird.

## Introduction

In the summer of 1995, tree stems and pieces of a Cyperaceae fen peat were found in the Vorfeld of the Unteraar Glacier (Central Alps of Switzerland) (Fig.1). Due to the actual retreat, they were eroded from the glacier's bed, transported by meltwater outburst and deposited just in front of the glacier at an altitude of 1920 m a.s.l. It could be seen with the naked eye that the cake-shaped organic sediment clasts were full of insect fragments.

The analysis of coleopteran assemblages to reconstruct palaeoclimate and palaeoenvironments is well established (Atkinson et al. 1986, 1987; Coope et al. 1998). In the region of the Alps, only a few investigations about fossil beetles have been made so far. Ponel (1992, in: Tessier et al. 1993) used the

method to reconstruct timberline fluctuations in the French Alps. A study of a high altitude site in the Pyrénées has recently been carried out by the same author (Ponel et al. 1999).

During the last decades, different records have been included to register Holocene climatic oscillations in the Alps. Gamper & Suter (1982) made a compilation of data from different authors. "25 000 Jahre Gletschergeschichte" is the title of a publication where Furrer (1990) summarised twenty years of research of Swiss glaciers at the University in Zurich. Several articles about treeline fluctuations in the Swiss Alps during the Holocene are included in "Oscillations of the Alpine and Polar Tree Limits in the Holocene" (Frenzel, ed. 1993). Beside dendrochronology, glacial morphology, fossil soils, pollen

and plant macrofossils, insect analysis is able to furnish important data to complete the picture about Holocene palaeoenvironment.

### Locality and History

The Unteraar Glacier is situated north of the Grimsel Pass in the Bernese Oberland. It is the fourth largest ice-stream of the Alps. Zumbühl (in: Zumbühl & Holzhauser 1988) describes the Unteraar Glacier as the cradle of glacial field research; the first cartographic depiction dates back to the beginning of the eighteenth century. Famous naturalists like Scheuchzer, Escher, Agassiz and others visited and described the impressive glacier, which is 6 km long today. Since 1876, the Unteraar Glacier has been retreating at an average of 16.9 m per year (<sup>2</sup>Data from World Glacier Monitoring Service). Investigations did not stop in this century. Frey (1921, 1922) made detailed descriptions about the vegetation before inundation. In 1931 the Vorfeld Aarboden was dammed to gain hydroelectric power. Glaciological studies were carried out by Haefeli (1970), Gudmunsson (1994) and others. The area of the Grimsel Pass attracted entomologists too (C. Huber, personal communication 1999). As far as the author knows, however, there is no literature which deals specifically with the Unteraar region. The finds made by different collectors were recorded in general overviews of distribution of beetle species in Switzerland (e.g. Marggi 1992, Puthz 1981, Comellini 1974).

### Radiocarbon dating

To date the peat, six radiocarbon measurements were carried out by Hormes et al. (1998). Five twigs of *Salix* sp. extracted from the peat and one bulk sample were dated. The measurements were made at the underground laboratory of Climate and Environmental Physics at the University of Bern using the conventional  $\beta$ -counting method. The samples yielded ages between  $3789 \pm 25$  and  $3622 \pm 31$  conventional radiocarbon years before present. According to the small range of <sup>14</sup>C-dated samples, we conclude that they belong to the same fen peat unit.

### Material and Methods

Thirteen peat samples of 1 kg to 5 kg were processed. They were sawed into two halves. One half was used for insect analysis, the second half put aside for <sup>14</sup>C and other investigations. The compressed and hard peat was composed of sedges, brown moss, flattened twigs and silty layers. Twelve kilograms of sediment yielded more than 5000 insect fragments.

To extract the insects, the standard process described by Coope (1986) and Elias (1994) was applied. In a first step, the sediment was put in a plastic bowl and soaked in water, then gently split by hand along the layers. Due to its solid consistency, the material had to be soaked in a weak sodium carbonate ( $\text{Na}_2\text{CO}_3$ ) solution for several days and heated at low temperature for a few hours. The sediment was dispersed again by hand. To remove mineral particles such as silt and fine sand, the slurry was washed several times through a sieve of 300 micron mesh diameter. Once the detritus had been screened, a handful of the residue was placed in a bowl and thoroughly mixed with paraffin. This treatment makes the smooth impermeable insect sclerites adhere to the oil. When cold water is added to the mixture, the oil and insects float on the surface. The floating fraction was carefully decanted. This process had to be repeated until no insect remains were floating on the surface anymore. To remove the paraffin, the concentrate was first washed with detergent and then with ethanol. Since some plant debris was inevitable decanted together with the insect remains, the decanted fraction had to be sorted under a binocular microscope. Heavily sclerotised fragments of Coleopterans were mounted on cards used in micropalaeontology with a water-soluble glue. Frail and small elements were put in vials and stored in ethanol.

### Insect analysis



Identification of fossil insects requires a modern reference collection. Keys of present-day Coleoptera are based on exoskeletal criteria and therefore useful to distinguish fossil species. However, as they are always based on whole specimens, they are helpful only to certain extent. An exception is Goecke (1943) who provides a key to the elytra of European Donaciinae (leaf beetles). In general, we worked with the series of Freude et al. (1964–83) and the supplementary volumes of Lohse & Lucht (1992). The illustrations in Reitter (1908–16) were useful in cases where the assignment to a family was difficult. Special literature was needed for *Helophorus* sp. (Angus 1992), the Omaliines (Zanetti 1987), *Olophrum* sp. (Campbell 1983) and *Aphodius* sp. (Dellacasa 1983). In many cases, direct comparison with modern material combined with taxonomic instruction led to successful identification.

However there are groups of beetles that cannot be identified to species level if only preserved in fragments. One example is the subfamily Aleocharinae. Individuals that could not be unambiguously assigned to a species (indicated by "cf.") were not included in ecological and climatic interpretation of the fauna. Specimens identified by the author were checked by different specialists.

In some cases, male genitalia found in the sediment could be assigned to a genus. The aedeagus (male genitalia) is diagnostic for many coleopteran species and the most certain criterion for identification. Several abdomens of the staphilinid beetle *Stenus bifoveolatus* were preserved together with the

---

<sup>2</sup> World Glacier Monitoring Service, Swiss Federal Institute of Technology, Laboratory of Hydraulics, Hydrology, and Glaciology, Gloriastrasse 37/39, CH-8092 Zurich, Switzerland



**Plate 1.**  
Fragments of fossil Coleoptera photographed with different techniques

The scale bar is 1 mm in a, and 0.25 mm in b-i

Fig. a photographed with digital camera  
left elytron of *Nebria castanea* Bon.

Fig. b and c conventional photographs on microscope  
b modern (left) and fossil (right) aedeagus of *Agabus congener* (Thunb.)  
c pronotum of *Agabus congener* (Thunb.)

Figs. d-i scanning electron micrographs  
d head of male *Anthophagus alpinus* (F.)  
e head of *Eucnecosum brachypterum* (Grav.) / *tenuis* (LeConte)  
f pronotum of *Olophrum consimile* (Gyll.)  
g left elytron of *Helophorus glacialis* Villa  
h right elytron of *Gynmusa brevicollis* (Payk.)  
i head and pronotum of *Stenus bifoveolatus* Gyll.

aedeagus. After the provisional identification of a pronotum as *Agabus congener* (Dytiscidae) (Plate 1, Fig. c), we recognised an isolated male genitalia of this species in the same sample (Plate 1, Fig. b), which confirmed the identification.

### Faunal assemblage

The most abundant sclerites found in the peat are derived from beetles. Remains from spiders, ants, caddisflies, mites, Chalcidoidea, Tipulidae and Diptera were picked out and stored but not yet further identified. In the discussion of the beetle assemblage, the fossil content of the thirteen samples will be considered as a single unit since they are very similar to one another and the radiocarbon dates suggest that they all belong to the same fen peat unit.

For each species in the faunal list (Table 1), the minimum number of individuals is calculated from the maximum number of any identifiable skeletal element. An example: Four left elytra, two right elytra, three pronota and one head of a certain species gives a minimum number of four individuals.

The abundance of staphilinids (rove beetles) in species and number of individuals is remarkable. *Eucnecosum brachyptenum/tenue*, *Olophrum consimile* and *Stenus bifoveolatus* (Plate 1, Fig. e, f, i) are represented by hundreds of fragments. All three species are hygrophilous and require a similar habitat. It has to be mentioned that *Stenus bifoveolatus* and *Gymnusa brevicollis* (Plate 1, Fig. h) are not occurring in the alpine zone (Koch 1989).

<i>Olophrum consimile</i> (Gyll.) formerly <i>Olophrum alpinum</i> (Heer)	associated with Cyperaceae, Graminae and moss along running water in alpine habitats (pers. comm. Zanetti 1999)
<i>Eucnecosum tenue</i> (LeConte)	associated with Cyperaceae in alpine lakes and marshes (pers. comm. Zanetti 1999)
<i>Stenus bifoveolatus</i> Gyll.	typical marsh inhabitant, occurring from the plain up to 1800 m a.s.l. (pers. comm. Puthz 1998)
<i>Gymnusa brevicollis</i> (Payk.)	mires, marshy surroundings, typhophilous, often in <i>Sphagnum</i> , not alpine (Koch 1989)

Aquatic species such as the dytiscid beetles *Agabus congener*, *Hydroporus memnonius* and *Hydroporus nigrita* are found in marshy waters. Focarile (1981) reports *H. memnonius* as typical for small lakes with vegetation at altitudes of 1900–2000 m in Kanton Tessin.

<i>Hydroporus nigrita</i> (F.)	cold-stenothermal, acidophilous, shady springs, fen pools and mountain streams, forest streams rich with leaves (Koch 1989)
<i>Hydroporus memnonius</i> Nic.	stenotopic, acidophilous, marshy waters, pools in wood rich with leaves (Koch 1989)
<i>Agabus congener</i> (Thunb.)	stenotopic, marshy waters (Koch, 1989). Recorded as high as 2200 m a.s.l. (Freude, Harde & Lohse 1971)

There are several cold-stenothermic species belonging to different families. Among the Carabidae (ground beetles) there is *Nebria castanea* (Plate 1, Fig. a), a wingless species, which is often found near melting snow. The dung beetle *Aphodius abdominalis* is mainly reported from the dwarf-shrub heath and along snowfields. Focarile (1985) mentions the staphilinid *Quedius punctatellus* as a species living in alpine meadows and accidentally occurring in the *Alnetum*. According to Koch (1989) it is also found in snow depressions. Horion (1965) reports that it is frequently occurring in subalpine forests. Among the cold tolerant taxa the water beetle *Helophorus glacialis* (Plate 1, Fig. g) needs the most severe climatic conditions. It lives on the edges of snow patches and in meltwater ponds. The presence of this hydrophilid "suggests the persistence of snow-patches in summer" (Ponel 1997).

<i>Nebria castanea</i> Bon.	everywhere in mountain areas, damp and wet sites, mostly close to melting snow (Marggi 1992)
<i>Helophorus glacialis</i> Villa	"a northern and mountain species ... It is typically found in areas just uncovered by melting snow" (Angus 1992)
<i>Quedius punctatellus</i> Heer	numerous in moss, at high altitudes (Comellini 1997). Forests, zone of larch and green alder, alpine meadows, snow depressions (Koch 1989)
<i>Aphodius abdominalis</i> (Bon.)	mainly in the dwarf-shrub heath, also in <i>Loiseleurietum</i> and on the edges of snow patches (Koch 1989)

Beside the already mentioned species living in moss and reeds, the surrounding vegetation is indicated by some Omaliines, a subfamily of staphilinids. They indicate flowering meadows and brushwood (Zanetti 1987). The aquatic leaf beetles *Donacia* and *Plateumaris* feed on different sedges. As its name implies, the small weevil (Curculinidae) called *Rhynchaenus salicis* depends on the willow (*Salix* sp.).

<i>Eusphalerum alpinum</i> (Heer)	flowering meadows in subalpine and alpine regions (Zanetti 1987)
<i>Amphichroum hirtellum</i> (Heer)	flowering shrubs, especially <i>Alnus</i> and <i>Rhododendron</i> (Zanetti 1987)
<i>Anthophagus bicornis</i> (Block)	on shrubs, mountainous and subalpine regions (Zanetti 1987)
<i>Anthophagus alpinus</i> (F.)	on shrubs, especially <i>Rhododendron</i> , subalpine regions (Zanetti 1987)
<i>Anthophagus omalinus</i> (Zett.)	on shrubs, mountainous and subalpine regions (Zanetti 1987)
<i>Plateumaris sericea</i> (L.)	oligophagous on <i>Carex</i> sp. (Koch 1992)
<i>Rhynchaenus salicis</i> (L.)	oligophagous on <i>Salix</i> sp. (Koch 1992)

Before drawing any conclusions about paleoecology, we have to ask whether the fauna in this present case is autochthonous (Ponel & Richoux 1997). Many beetles are good fliers or can be transported by wind. Further, streams could have brought in some of the insects. Ponel (1997) examined a high altitude site in the Northern Apennines and reported an influx of wind-blown elements. Kenward (1975, 1976) investigated death assemblages from a drain and from roof gullies. He emphasises

that the background fauna (i. e. the transported component) may be predominant where the local fauna is sparse. For Kenward, reliable evidence that fossils lived close to the point of deposition is the abundance of a group of species with similar requirements, as long as the total fauna is large. If a certain species is represented by numerous individuals, it could be an indication that it was attracted to the site of the deposit or its surroundings or maybe even that it did breed there. These conditions recommended for the reconstruction of past environments at the Unteraar site are fulfilled: The fauna is large, some species are represented by a large number of individuals. The fossil assemblage contains many species of different families with very similar ecological requirements. Even very frail sclerites are in a good state of preservation, which indicates that they have not been transported over long distances after death. Fossils of insects and their host plants when found together in the same samples confirm that the assemblage is largely in situ.

The ecology depicted by the beetle fauna is very consistent in itself. An overwhelming abundance of hygrophilous Coleoptera suggests a marshy habitat with a pond but also running water nearby. Species exclusively dependent on trees are absent. There are some species in the assemblage though that are not known to occur in alpine areas. The information furnished by different specimens about the vegetation indicates a position in the dwarf-shrub heath. We locate the ancient fen peat in the upper subalpine zone, above the forest limit but below treeline.

### Climatic reconstruction

Quantitative estimates of paleotemperatures based on coleopteran assemblages can be made by using the Mutual Climatic Range method (MCR). This method, developed by Atkinson et al. (1986) uses the correlation between the modern distribution of a particular beetle species and thermal climatic factors, which are measured by meteorological stations. Each species can be plotted on a climatic envelope defined by  $T_{\max}$  (the mean temperature of the warmest month) and  $T_{\text{range}}$  (the difference between the mean temperature of the warmest month and the mean temperature of the coldest month).  $T_{\min}$  (the mean temperature of the coldest month) can be calculated from these two co-ordinates. The area of overlap of the climatic envelopes of the species comprising the fossil community is called the mutual climatic range and defines the paleoclimatic conditions. Tests of this method on modern beetle faunas showed slight systematic deviations which can be avoided using correction equations (Atkinson et al. 1987). In the following case we applied the slightly revised version of the regression equations as published in Coope et al. (1998).

To reconstruct  $T_{\max}$  and  $T_{\min}$  from the Unteraar fen peat, twelve species with established climatic envelopes were used:

*Nebria rufescens*  
*Pterostichus diligens/strenuus*

*Hydroporus memnonius*  
*Agabus congener*  
*Helophorus glacialis*  
*Omalium excavatum*  
*Eucnecosum brachypterum/tenue*  
*Acidota crenata*  
*Anthophagus bicornis*  
*Tachinus elongatus*  
*Gymnusa brevicollis*  
*Aegialia sabuleti*

Calculation of temperature for this assemblage provided the following results:

Unteraar, 1920 m a.s.l., 3789 ± 25 to 3622 ± 31 yr BP (<sup>14</sup>C age)  
 $T_{\max}$  median (calibrated) = 9.0°C  
 $T_{\min}$  median (calibrated) = -7.7°C

These temperatures may be compared with the actual ones, measured at the Grimsel Hospiz, 5.5 km east of the Unteraar site (Fig. 1):

Grimsel Hospiz, 1950 m a.s.l., 1964–1998  
 average temperature of July = 9.5°C  
 average temperature of January = -5.3°C

It is conspicuous how similar  $T_{\max}$ , inferred from the fossil beetle assemblage, is to the average of the measured July temperatures of the last 35 years at Grimsel Hospiz. According to these values it was slightly cooler during the warmest month (0.5°C) between 3800 and 3600 yr BP than today. But this small difference is probably within the experimental limits of the MCR method. The value calculated by the MCR method for the coldest month ( $T_{\min}$ ) suggests a temperature in January that was about 2.5°C lower.

Comparing temperatures from Unteraar and the Hospiz however we have to bear in mind that in spite of the short distance microclimate can be different. Frey (1921, p.107) reports that temperatures measured in summer 1918 at the bottom of the valley near the glacier were 2° C higher than further along the valley. Further on it is likely that the water sheet of the dammed Unteraar Lake influences (buffers) the temperatures at Grimsel Hospiz today.

### Discussion and Conclusions

During the growth of the fen peat, the Unteraar Glacier was smaller than today. The fossil beetle assemblage contains no tree-dependent species suggesting that trees may not have been growing in the immediate vicinity of the peat at that time. Among the dated wood fragments however (Hormes et al. 1998), there were three samples within the range of peat growth. One sample was *Pinus cembra*, another *Picea* sp. and a third either *Picea* or *Larix*. In the peat we found a male in-

Table 1. List of Coleoptera arranged in the systematic order proposed by Freude, Harde and Lohse (1965–83). The numbers indicate the minimum number of individuals per sample.

Species	Sample number													Total number of individuals
	1	2	3	4	5	6	7	8	9	10	11	12	13	
<b>CARABIDAE</b>														
<i>Cicindela hybrida transversalis</i> Dej.	1	-	-	-	-	-	-	-	-	-	-	-	-	1
<i>Nebria rufescens</i> Ström	1	-	-	-	-	-	-	-	-	-	-	-	-	1
<i>Nebria castanea</i> Bon.	-	-	1	-	-	-	-	-	-	1	-	-	1	3
<i>Pterostichus strenuus</i> (Panz.) / <i>diligens</i> (Sturm)	-	-	-	-	-	-	-	-	-	-	1	-	-	1
<b>DYTISCIDAE</b>														
<i>Hydroporus nigrita</i> (F.)	-	2	2	-	-	-	-	1	-	8	1	1	-	15
<i>Hydroporus memnonius</i> Nic.	1	-	9	-	-	-	-	-	-	-	2	-	-	12
<i>Hydroporus</i> spp.	1	-	5	1	-	4	-	1	-	7	3	-	-	22
<i>Agabus biguttatus</i> (Ol.)	1	-	-	-	-	-	-	-	-	-	-	-	-	1
<i>Agabus congener</i> (Thunb.)	-	-	1	-	-	-	-	-	-	1	-	1	-	3
<i>Agabus</i> cf. <i>congener</i>	-	-	-	-	-	-	-	-	-	-	-	-	1	1
<i>Agabus</i> sp.	-	1	-	-	-	-	-	-	-	-	-	-	-	1
<b>HYDROPHILIDAE</b>														
<i>Helophorus glacialis</i> Villa	-	1	-	1	-	4	-	1	1	19	3	-	1	31
<i>Enochrus ochropterus</i> (Marsh.)	-	-	-	-	-	-	-	-	-	1	-	-	-	1
<i>Enochrus ochropterus</i> (Marsh.) / <i>quadripunctatus</i> (Hrbst)	-	-	-	-	-	-	-	-	-	1	-	-	-	1
<b>STAPHILINIDAE</b>														
cf. <i>Proteinus</i>	-	-	-	-	-	-	-	1	-	-	-	-	-	1
<i>Eusphalerum alpinum</i> (Heer)	2	-	1	-	-	-	-	-	1	3	2	-	-	9
<i>Eusphalerum</i> cf. <i>alpinum</i>	-	-	-	-	-	-	-	-	-	2	-	-	-	2
<i>Eusphalerum anale</i> (Er.)	-	-	-	-	-	1	-	-	-	-	-	-	-	1
<i>Eusphalerum</i> cf. <i>limbatum</i> (Er.)	-	-	-	-	-	-	-	-	-	1	-	-	-	1
<i>Eusphalerum</i> spp.	2	2	-	-	1	-	2	-	2	6	2	1	-	18
<i>Omalium caesum</i> Grav.	-	1	-	-	-	-	-	-	-	-	-	-	-	1
<i>Omalium excavatum</i> Steph.	-	-	-	-	-	-	-	-	-	-	-	1	-	1
<i>Omalium</i> sp.	-	-	1	-	1	1	-	1	-	-	-	-	2	6
<i>Olophrum consimile</i> (Gyll.), jun. syn. <i>O. alpinum</i> (Heer)	16	12	9	10	7	5	7	6	3	51	24	5	20	175
<i>Olophrum</i> sp.	31	2	2	11	2	2	-	6	1	1	12	1	23	94
<i>Eucnecosum brachypterum</i> (Grav.) / <i>tenuis</i> (LeConte)	68	122	13	53	23	10	8	26	8	117	95	122	59	724
<i>Acidota crenata</i> (F.)	-	-	-	-	-	-	-	-	1	-	-	-	-	1
<i>Amphichroum canaliculatum</i> (Er.) / <i>hirtellum</i> (Heer)	1	6	-	1	6	-	1	1	3	1	-	-	-	20
<i>Amphichroum hirtellum</i> (Heer)	6	2	-	4	1	11	-	-	-	30	11	1	5	71
<i>Lesteva</i> cf. <i>monticola</i> Kiesw.	-	-	-	-	-	-	-	-	-	1	-	-	-	1
<i>Anthophagus bicornis</i> (Block)	-	-	-	-	-	1	-	-	-	-	1	-	-	2
<i>Anthophagus</i> cf. <i>bicornis</i>	1	-	2	-	-	-	-	-	-	1	-	-	-	4
<i>Anthophagus alpinus</i> (F.)	5	1	-	-	2	5	-	-	-	9	4	-	-	26

<i>Anthophagus alpinus</i> (F.) / <i>fallax</i> Kiesw.	6	3	1	1	-	-	-	2	-	11	2	-	3	29
<i>Anthophagus omalinus</i> Zett.	4	2	1	2	-	3	-	2	-	7	4	-	1	26
<i>Anthophagus</i> cf. <i>omalinus</i>	-	-	-	-	-	-	-	-	-	1	-	-	-	1
<i>Anthophagus</i> cf. <i>alpestris</i> Heer	-	-	-	-	-	-	-	-	-	-	-	-	1	1
<i>Anthophagus</i> spp.	-	3	1	1	1	6	5	3	2	13	3	2	1	41
<i>Stenus</i> cf. <i>gallicus</i> Fauv.	-	-	1	-	-	-	-	-	1	-	-	1	1	4
<i>Stenus</i> cf. <i>pusillus</i> Steph.	-	-	-	1	-	-	-	-	-	-	-	-	-	1
<i>Stenus</i> cf. <i>oscillator</i> Rye	-	-	-	-	-	-	-	-	1	-	-	-	-	1
<i>Stenus bifoveolatus</i> Gyll.	45	19	13	14	27	3	7	2	4	8	21	65	10	238
<i>Stenus</i> sp.	-	2	-	-	-	-	-	2	-	-	1	2	-	7
<i>Philonthus</i> spp.	-	2	-	-	-	-	-	-	-	1	1	-	1	5
<i>Gabrius</i> spp.	-	2	-	-	-	-	-	-	-	-	1	-	-	3
Quediini indet	-	-	-	-	-	1	-	-	-	-	-	-	1	2
<i>Quedius punctatellus</i> Heer	1	1	1	1	1	-	-	1	1	1	-	-	1	9
<i>Quedius</i> spp.	-	-	1	1	-	1	-	-	1	-	-	-	-	4
<i>Mycetoporus</i> s.l.	-	-	1	-	-	-	-	-	-	-	-	-	-	1
<i>Tachinus elongatus</i> Gyll.	-	-	-	-	-	-	-	-	-	1	-	-	-	1
Aleocharinae indet	9	28	10	8	4	2	-	2	2	19	14	18	5	121
<i>Gymnusa brevicollis</i> (Payk.)	1	1	5	1	1	-	-	-	-	1	4	2	-	16

#### SCARABAEIDAE

<i>Aegialia sabuleti</i> (Panz.)	-	1	-	-	-	-	-	-	-	-	-	-	-	1
cf. <i>Aegialia sabuleti</i>	1	-	-	-	-	-	-	-	-	-	-	-	-	1
<i>Aphodius abdominalis</i> (Bon.)	1	1	-	1	1	-	-	1	-	1	1	-	-	7

#### SCIRTIDAE

<i>Cyphon</i> sp.	-	-	-	-	-	-	-	-	-	4	-	-	-	4
-------------------	---	---	---	---	---	---	---	---	---	---	---	---	---	---

#### BYRRHIDAE

<i>Cytilus sericeus</i> (Forst.)	1	-	-	-	-	-	-	-	-	-	-	-	-	1
<i>Cytilus sericeus</i> (Forst.) / <i>auricomus</i> (Duft.)	-	-	2	-	-	-	-	-	-	-	-	-	-	2
<i>Byrrhus</i> sp.	-	-	-	-	-	1	-	-	-	1	-	-	-	2

#### CHRYSOMELIDAE

<i>Donacia impressa</i> Payk.	-	-	-	-	-	-	-	-	-	1	-	-	-	1
<i>Donacia impressa</i> Payk. / <i>aquatica</i> (L.)	-	-	-	-	-	-	-	-	-	1	-	-	-	1
<i>Donacia</i> cf. <i>brevitarsis</i> Thoms.	-	-	-	-	-	-	-	-	-	1	-	-	-	1
<i>Donacia obscura</i> Gyll.	-	-	1	-	-	-	-	-	-	-	-	-	-	1
<i>Donacia</i> cf. <i>obscura</i>	1	-	-	1	-	-	-	-	-	-	-	-	-	2
<i>Donacia</i> cf. <i>thalassina</i> Germ.	-	-	-	-	-	-	-	-	-	1	-	-	-	1
<i>Donacia vulgaris</i> Zsch.	-	-	-	-	-	-	-	-	-	1	-	-	-	1
<i>Donacia</i> spp.	-	1	-	1	1	1	-	-	-	1	-	-	-	5
<i>Plateumaris sericea</i> (L.)	1	-	1	-	1	-	-	-	-	1	-	-	-	4
<i>Phratora</i> sp.	-	-	-	-	-	-	1	-	-	-	-	1	1	3
<i>Altica</i> cf. <i>lythri</i> Aubé	-	-	-	-	-	-	-	-	-	-	2	-	-	2
<i>Altica</i> sp.	-	-	-	-	-	-	-	-	-	-	-	2	-	2

#### CURCULIONIDAE

Curculinidae indet	1	1	-	-	1	-	-	1	-	1	-	-	-	5
<i>Rhynchaenus salicis</i> (L.)	-	-	-	1	-	-	-	-	-	1	-	-	-	2
<i>Rhynchaenus</i> sp.	-	-	-	1	-	-	-	-	-	-	1	-	-	2






Fig. 1. Map of Switzerland and the Grimsel area with the location of the site

florescence of *Pinus* (not *P. cembra*, pers. comm. M. Winter 1999). We do not consider these facts to be contradictory. It has to be taken into account that the fen peat was located in the Vorfeld of the former glacier where the conditions for tree growth are not favourable due to the unstable geological regime (braided river system). On the vegetation map by Frey (1921), where the situation before inundation is depicted, we find no trees in the Aarboden (Vorfeld / bottom of the valley). He reports that *Pinus cembra*, *Larix decidua*, *Betula alba* prefer to grow on the slopes where they are spread as far as the terminus of the glacier. *Pinus cembra* occurs up to 2100 m a.s.l. on the southerly exposed slope, called "Sonnige Aar" (Fig. 1), and up to an elevation of 2080 m a.s.l. on the northerly exposed slope. A similar situation for the period of 3800–3600 yr BP is conceivable but further upvalley. The fact that the Gletschervorfeld was not forested, indicated by the absence of tree-dependent Coleoptera and the abundance of species living on flowering shrubs and herbs, rules out a tim-

berline much higher than today. This conclusion is supported by other authors who have tried to locate the elevation of timberline in the Swiss Central Alps for this period. Welten (1982), who has made extensive investigations about vegetation history in the Western Swiss Alps, has shown that the era of good alpine forests and high elevation of the timberline lasted until 5000–4000 yr BP. Use by Neolithic man and cattle brought about an end to this period. The event coincides with the mass expansion of *Alnus viridis*. Palaeobotanical data of Tinner et al. (1996) show that Alpe Essertse in the Valais was covered with a forest of *Pinus cembra* and *Larix decidua* up to at least 2343 m a.s.l. from 9500 to 3600 yr BP. After 4700 yr BP the forest became more open and timberline moved down either due to climatic changes or human impact. Wick (in: Wick & Tinner 1997) reports mass expansion of *Alnus viridis* and a continuous downward movement of the timberline at Lago Basso (2250 m a.s.l.) in the Italian Central Alps between 4500 and 3000 yr BP.

Although fluctuation of timberline can indicate climatic changes, estimations of paleotemperatures are difficult to make (Markgraf 1974). A rough approximation is the coincidence of the timberline with the 10°C July isotherm (Eggenberg 1995). Investigations of fossil coleopteran assemblages render a helpful contribution to the question of paleotemperatures (Coope et al. 1998; Ponel et al. 1999).

The period between 4400 and 3600 yr BP is known as the Postglacial Climatic Optimum in the Swiss Alps. Dendroclimatology indicates the longest time span with temperatures above the average between 3760 and 3600 yr BP (Gamper & Suter 1982). For the Eastern Alps, Patzelt suggested mean summer temperatures had been the same as today between 4100 and 3500 yr BP (in: Slupetzky et al. 1998). He postulated (Patzelt 1977) that summer temperatures during the last 9500 years did not fluctuate more than 1.5°C over a longer time span. The study of the fossil beetle assemblage suggests a similar July temperature (9.0°C) at Unteraar between 3800 and 3600 yr BP as today (9.5°C). Calculated mean temperature for the coldest month is -7.7°C compared to the average of the last 35 years at Grimsel Hospiz with -5.3°C in January.

#### Acknowledgements

In particular I would like to thank Professor Ch. Schlüchter (Bern) who initiated the project and provided the financing. In the Quaternary Entomology Laboratory of Fargo, North Dakota State University, I was introduced to the technique of extracting fossil beetles and collecting living ones in the field. The weeks I spent with Professor A. Ashworth and his team were very stimulating! I am indebted to Professor R. Coope who has always had time to teach me identification during his visits to Switzerland. His valuable comments greatly improved the content of this paper. Many questions were answered by Dr. P. Ponel (Marseille) and Dr. S. Elias (Boulder) and I thank them for their support. I am grateful for the help of several specialists who made or checked the identifications of my beetles: R. Angus, M. Brancucci, M. Döberl, J. Gebert, E. Geiser, C. Huber, B. Klausnitzer, W. Marggi, M. Miler, V. Puthz. Special thanks are due to Dr. A. Zanetti in Verona whose comments on the subfamily of Omaliinae improved the accuracy of the faunal list. At the Museum of Natural History in Bern I always felt welcome. I had access to the collection and was given assistance by the members of the Invertebrates Department. With Anne Hormes I had inspiring discussions about the Unteraar peat. It was nice to have someone working on the same site. Many thanks go to Emma Jane Sayer and Dr. W. Shotyk for proof-reading.

#### REFERENCES

ANGUS, R. 1992: Insecta: Coleoptera: Hydrophilidae: Helophorinae. In: Schwoerbel, J. & Zwick, P., editors, Süßwasserfauna von Mitteleuropa 20/10-2. Stuttgart: Gustav Fischer Verlag, 1-144.

ATKINSON, T. C., BRIFFA, K. R., COOPE, G. R., JOACHIM, M. J. & PERRY, D. W. 1986: Climatic calibration of coleopteran data. In: Berglund, B. E., editor, Handbook of Holocene Palaeoecology and Palaeohydrology, London: John Wiley, 851-858.

ATKINSON, T. C., BRIFFA, K. R. & COOPE, G. R. 1987: Seasonal temperatures in Britain during the past 22,000 years, reconstructed using beetle remains. *Nature* 325, 587-592.

CAMPELL, J. M. 1983: A revision of the North American Omaliinae (Coleoptera Staphilinidae): The genus *Olophrum* Erichson. *Canadian Entomologist* 115, 577-622.

COMELLINI, A. 1974: Notes sur les Coléoptères Staphylinides de haute altitude. *Revue suisse de zoologie* 81/2, 511-539.

COOPE, G. R. 1986: Coleoptera analysis. In: Berglund, B. E., editor, Handbook of Holocene Palaeoecology and Palaeohydrology, London: John Wiley, 703-713.

COOPE, G. R., LEMDAHL, G., LOWE, J. J. & WALKING, A. 1998: Temperature gradients in northern Europe during the last glacial-Holocene transition (14-9 14C kyr BP) interpreted from coleopteran assemblages. *Journal of Quaternary Science* 13/5, 419-433.

DELLACASA, G. 1983: Monografie, 1 - Sistematica e nomenclatura degli Aphodiini italiani (Coleoptera Scarabaeida: Aphodiinae). Torino: Museo regionale di Scienze Naturali, 463 p.

EGGENBERG, S. 1995: Ein biogeographischer Vergleich von Waldgrenzen der nördlichen, inneren und südlichen Schweizeralpen. *Mitteilungen der Naturforschenden Gesellschaft in Bern, N.F.* 52, 97-120.

ELIAS, S. A. 1994: Quaternary Insects and Their Environments. Washington: Smithsonian Institution Press, 284 p.

FOCARILE, A. 1981: Connaissance actuelles sur les Coléoptères de haute altitude du Tessin. *Bollettino della Società Ticinese di Scienze Naturali* 69/2, 21-51.

- 1985: Le cenosi fitosaprobie di Coleotteri in lettiera di Ontano Verde (*Alnus viridis* Chaix) nelle Alpi occidentali. *Bollettino del Museo regionale di scienze naturali*, Torino 3/1, 79-126.

FRENZEL, B. (editor), 1993: Oscillations of the Alpine and Polar Tree Limits in the Holocene. *Palaeoclimate Research* 9. Stuttgart: Gustav Fischer Verlag, 234 p.

FREUDE, H., HARDE, K. W. & LOHSE, G. A. (editors) 1964-83: Die Käfer Mitteleuropas 1-11. Krefeld: Goecke & Evers Verlag.

FREY, E. 1921: Die Vegetationsverhältnisse der Grimselgegend im Gebiet der zukünftigen Stauseen. *Mitteilungen der Naturforschenden Gesellschaft in Bern*, Heft VI, 85-281.

- 1922: Die Arven- und Lärchenbestände im Unteraartal. *Schweizerische Zeitschrift für Forstwesen* 73/1 + 2, 10.

FURRER, G. 1990: 25 000 Jahre Gletschergeschichte dargestellt an einigen Beispielen aus den Schweizer Alpen. Veröffentlichung der Naturforschenden Gesellschaft in Zürich im Anschluss an den Jahrgang 135 der Vierteljahrsschrift, Heft 5.

GAMPER, M. & SUTER, J. 1982: Postglaziale Klimageschichte der Schweizer Alpen. *Geographica Helvetica* 37/2, 105-114.

GOECKE, H. 1943: Monographie der Schilfkäfer. II. Die fossilen Funde und ihre Bestimmung. In: Aberhalden, E., editor, *Nova Acta Leopoldina*, Neue Folge, Band 12, Nummer 86, 339-380.

GUDMUNSSON, G. H. 1994: Convergent glacier flow - a case study: The Unteraarglacier. *Mitteilungen der Versuchsanstalt für Wasserbau, Hydrologie und Glaziologie der Eidgenössischen Technischen Hochschule Zürich* 131, 120 p.

HAEFFELI, R. 1970: Changes in the behaviour of the Unteraargletscher in the last 125 years. *Journal of Glaciology* 9/56, 195-212.

HORION, A. 1965: Faunistik der mitteleuropäischen Käfer, Band 10. Überlingen-Bodensee: Kommissionsverlag Buchdruckerei Aug. Feyel.

HORMES, A., SCHLÜCHTER, C. & STOCKER, T. F. 1998: Minimal extension phases of Unteraarglacier (Swiss Alps) during the Holocene based on <sup>14</sup>C analysis of wood. *Radiocarbon* 40/2, 809-817.

KENWARD, H. K. 1975: Pitfalls in the Environmental Interpretation of Insect Death Assemblages. *Journal of Archaeological Science* 2, 85-94.

- 1976: Reconstructing ancient ecological conditions from insect remains: some problems and an experimental approach. *Ecological Entomology* 1, 7-17.

KOCH, K. 1989-92: Die Käfer Mitteleuropas, Ökologie 1-3. Krefeld: Goecke & Evers Verlag.

LOHSE, G. A. & LUCHT, W. H. 1992: Die Käfer Mitteleuropas, 2. Supplementband mit Katalogteil, Krefeld: Goecke & Evers Verlag.

- 1994: Die Käfer Mitteleuropas, 3. Supplementband mit Katalogteil, Krefeld: Goecke & Evers Verlag.

MARGGI, W. A. 1992: Faunistik der Sandlaufkäfer und Laufkäfer der Schweiz (Cicindelidae & Carabidae) Coleoptera. *Documenta Faunistica Helvetica* 13, Teil 1/Text, Teil 2/Verbreitungskarten, 477 p. and 243 p.

MARKGRAF, V. 1974: Palaeoclimatic evidence derived from timberline fluctuations. *Colloques Internationaux du C.N.R.S N° 219*, 67-77.

- PATZELT, G. 1977: Der zeitliche Ablauf und das Ausmass postglazialer Klimaschwankungen in den Alpen. In: Frenzel, B., editor, Dendrochronologie und postglaziale Klimaschwankungen in Europa. Erdwissenschaftliche Forschung, 13. Wiesbaden: Steiner, 248–259.
- PONEL, P., DE BEAULIEU, J.-L. & TOBOLSKI, K. 1992: Holocene palaeoenvironments at the timberline in the Taillefer Massif, French Alps: a study of pollen, plant macrofossils and fossil insects. *The Holocene* 2/2, 117–130.
- PONEL, P. 1997: Late Pleistocene Coleopteran Fossil Assemblages in High Altitude Sites: a Case Study from Prato Spilla (Northern Italy). *Quaternary Proceedings* 5. Chichester: John Wiley, 207–218.
- PONEL, P. & RICHOUX, P. 1997: Difficultés d'interprétation des assemblages de Coléoptères fossiles quaternaires en milieu d'altitude. *GEOBIOS, Mémoire spécial n° 21*, 213–219.
- PONEL, P., COOPE, G. R., ANDRIEU-PONEL, V. & REILLE, M. 1999: Coleopteran evidence for a mosaic of environments at high altitude in the eastern Pyrénées, France, during the climatic transition between the Allerød and Younger Dryas. *Journal of Quaternary Science* 14/2, 169–174.
- PUTHZ, V. 1981: Beitrag zur Faunistik der Staphiliniden der Schweiz (Coleoptera, Staphilinidae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 54, 165–170.
- REITTER, E. 1908–16: Die Käfer des Deutschen Reiches I–V, Fauna Germanica. Stuttgart: K.G. Lutz Verlag.
- SLUPETSKY, H., KRISAI, R. & LIEB, K. 1998: Hinweise auf kleinere Gletscherstände der Pasterze (Nationalpark Hohe Tauern, Kärnten) im Postglazial – Ergebnisse von  $^{14}\text{C}$ -Datierungen und Pollenanalysen. *Wissenschaftliche Mitteilungen aus dem Nationalpark Hohe Tauern Band 4*, 225–240.
- TESSIER, L., DE BEAULIEU, J.-L., COUTEAUX, M., EDOUARD, J.-L., PONEL, P., ROLANDO, C., THINON, M., THOMAS, A. & TOBOLSKI, K. 1993: Holocene palaeoenvironments at the timberline in the French Alps – a multidisciplinary approach. *Boreas* 22, 244–254.
- TINNER, W., AMMANN, B. & GERMANN, P. 1996: Treeline Fluctuations Recorded for 12,500 Years by Soil Profiles, Pollen, and Plant Macrofossils in the Central Swiss Alps. *Arctic and Alpine Research* 28/2, 131–147.
- WELTEN, M. 1982: Vegetationsgeschichtliche Untersuchungen in den westlichen Schweizer Alpen. Bern – Wallis. *Denkschriften der Schweizerischen Naturforschenden Gesellschaft* 95, 104p.
- WICK, L. & TINNER, W. 1997: Vegetation Changes and Timberline Fluctuations in the Central Alps as Indicators of Holocene Climatic Oscillations. *Arctic and Alpine Research* 29/4, 445–458.
- ZANETTI, A. 1987: Coleoptera Staphilinidae Omaliinae. *Fauna d'Italia* 25. Bologna: Calderini, 472 p.
- ZUMBÜHL, H. J. & HOLZHAUSER, H. 1988: Alpengletscher in der Kleinen Eiszeit. *Die Alpen* 64/3, Sonderheft. Bern: Schweizer Alpen-Club, 322 p.

Manuscript received October 28, 1999

Revision accepted May 25, 2000