**Zeitschrift:** Eclogae Geologicae Helvetiae

Herausgeber: Schweizerische Geologische Gesellschaft

**Band:** 96 (2003)

Heft: 3

**Artikel:** The oldest trigonalid wasp in the Late Albian amber of Charente-

Maritime (SW France) (Hymenoptera, Trigonalidae)

Autor: Nel, André / Perrichot, Vincent / Néraudeau, Didier

**DOI:** https://doi.org/10.5169/seals-169035

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 30.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

0012-9402/03/030503-6 DOI 10.1007/s00015-003-1104-3 Birkhäuser Verlag, Basel, 2003

# The oldest trigonalid wasp in the Late Albian amber of Charente-Maritime (SW France) (Hymenoptera: Trigonalidae)

André Nel<sup>1</sup>, Vincent Perrichot<sup>2</sup> & Didier Néraudeau<sup>2</sup>

Key words: Insecta, Hymenoptera, Trigonalidae, Albiogonalys elongatus gen. n., sp. n., amber, Late Albian, Charente-Maritime, France

#### ABSTRACT

Albiogonalys elongatus gen. n., sp. n., oldest known representative of the family Trigonalidae, is described from the Late Albian amber of France. It could be placed in a very basal position, as sister group of the modern representatives of the family. The positions of the fossil taxa currently attributed to this family are discussed. Except for *Cretogonalys taimyricus* RASNITSYN 1977, almost all of these taxa are too poorly preserved or described for accurate attributions to this family.

#### RESUME

Albiotrigonalis elongatus gen. n., sp. n., plus ancien représentant connu de la famille Trigonalidae, est décrit de l'ambre de l'Albien supérieur de France. Il est dans une position basale, comme groupe frère des représentants modernes de la famille. Les positions des taxons fossiles attribués à cette famille sont discutées. Mis à part Cretogonalys taimyricus Rasnitsyn, 1977, presque tous ces taxons sont trop mal conservés ou décrits pour une attribution correcte à cette famille.

#### 1. Introduction

The Trigonalidae is a small hymenopteran family that comprises 88 species in sixteen recent genera (Carmean & Kimsey 1998). Very few fossils have been attributed to this group, ranging from the Lower Cretaceous to the Oligocene: Turgonalus cooperi RASNITSYN & JARZEMBOWSKI 1998, described from the Lower Barremian of England (Rasnitsyn et al. 1998); Darbigonalus capitatus RASNITSYN 1986, from the Early Cretaceous of Gurvaneren Formation, West Mongolia (Rasnitsyn 1986); Cretogonalys taimyricus RASNITSYN 1977, from the Cenomanian amber of the Taimyr Peninsula, Russia (Rasnitsyn 1977); Turgonalus minor RASNITSYN 1990, from the Cretaceous of Transbaikalia, Russia (Rasnitsyn 1990); 'Trigonalys' bischofii STATZ 1938, from the Oligocene of Rott, Germany (Statz 1938). In addition, Martínez-Delclòs et al. (1999) listed undescribed specimens of the family 'Trigonalidae' in the Aptian/Albian Spanish amber of Alava. Furthermore, Cockerell (1917) described Trigonalys pervetus and (1920) listed the family 'Trigonalidae' in the Albian Burmese amber (recent Albian datation by Cruickshank & Ko 2003), although Antropov

(2000) transferred it in the Sphecidae. However, nearly all these fossils are too poorly preserved for really accurate systematic attributions. Thus, the present discovery of a fossil wasp belonging to this family in the Upper Albian amber of Archingeay (Charente-Maritime, France) is of great interest to estimate the age and origin of this family.

#### 2. Material and methods

The specimen studied here was discovered in a piece of amber together with four other insects (see the list of taxa in the systematic palaeontology section below). This lump, as the most of the amber lumps from the outcrop of Archingeay-Les Nouillers, is honey-coloured to clear brown and rather dusty. It was thus impossible to study this inclusion without preparing the specimen. First, each inclusion was separated from others, and the surrounding amber was removed in order to get as close as possible to the inclusion and to view its entire habitus. These operations were realized under a stereomicroscope using shaving blades transformed into micro-saws. Then, the separated specimens were embedded in Canada balsam be-

<sup>&</sup>lt;sup>1</sup> Laboratoire d'Entomologie and CNRS UMR 5143, Muséum National d'Histoire Naturelle, 45, Rue Buffon, F-75005, Paris, France. E-mail: anel@mnhn.fr

<sup>&</sup>lt;sup>2</sup> Géosciences Rennes, CNRS UMR 6118, Université Rennes1, Campus de Beaulieu, bât. 15, 263, Avenue du Général Leclerc, 35042 Rennes Cedex, France. E-mail: Vincent.Perrichot@univ-rennes1.fr, Didier.Neraudeau@univ-rennes1.fr




Fig. 1. – *Albiogonalys elongatus* gen. n., sp. n., holotype ARC 205.1; photography of general habitus. Scale bar represents 2 mm.

tween cover glasses, allowing an easier view of all sides of the insect during its study. We follow the terminology of Huber & Sharkey (1993) for the systematic description.

#### 3. Systematic palaeontology

Superfamily Trigonaloidea Cresson 1887 Family Trigonalidae Cresson 1887

## Genus Albiogonalys gen. n.

*Type species.* – *Albiogonalys elongatus* sp. n. *Gender.* – Masculine.

Etymology. – After the Albian age of this fossil, and 'gonalys' derived from the genus *Trigonalys*.

*Diagnosis.* – Easily distinguishable from modern taxa by the following characters: antennae with 25 cylindrical flagellomeres; in hind wing, part of M distal of r-m present and part of A




Fig. 2. – *Albiogonalys elongatus* gen. n., sp. n., holotype ARC 205.1; reconstruction of general habitus. Scale bar represents 2 mm.

distal of cu-a present as a nebulous vein, metasoma very elongate and narrow, about ten times as long as broad, tergite VII with two rather long lobes.

#### Albiogonalys elongatus sp. n. (Figs. 1-6)

Holotype. – Female specimen ARC 205.1, in the same piece of amber with two Coleoptera, one Hymenoptera: Proctotrupoidea, and one larva of Neuroptera: Myrmeleontoidea: Ascalaphidae? Deposited in the Laboratoire de Paléontologie, Muséum National d'Histoire Naturelle, Paris, France.

*Type locality.* – Archingeay, Charente-Maritime, SW France. *Stratigraphic horizon.* – Lower Cretaceous, Uppermost Albian (Néraudeau et al. 2002).

Etymology. – After its very elongate and narrow metasoma. *Diagnosis*. – That of the genus.

Description. – Body 5.4 mm long.




Fig. 3. – *Albiogonalys elongatus* gen. n., sp. n., holotype ARC 205.1; photography of left wings. Scale bar represents 1 mm.

Head 0.6 mm long, 1.0 mm wide, ratio length/width 0.6; labrum narrow elongate; right mandible strong, with four teeth, left mandible not visible; other mouthparts not or very poorly visible; three ocelli; a slight supra-antennal elevation above the torulus; antennae inserted on frons, 3.8 mm long, with a short scape, 0.2 mm long, ratio length of scape/length of antenna 0.05, pedicel 0.08 mm long, ratio length of pedicel/length of scape 0.4, 25 cylindrical flagellomeres, the apical one being conical; numerous sparce white scales on middle flagellomeres; malar space without a deep depression for reception of scape; eyes oval, large, well separated, 0.4 mm apart.

Mesosoma 1.88 mm long, 0.88 mm wide, ratio length/width 2.14; no triangular axilla and median groove on mesoscutum; propodeum broader than long, 0.4 mm long, 0.52 mm wide, ratio length/width 0.77, without any visible sculpturing ornamentation.

Legs. Only middle and hind legs are preserved, with five tarsomeres; tarsomeres 1–4 each with plantar lobe, 0.02 mm long; mid femur about 1 mm long, 0.2 mm wide, ratio length/width 5.0; tibia 1.12 mm long, 0.12 mm wide, ratio length/width 9.3, tarsus 1.48 mm long; hind femur about 1.2 mm long, 0.24 mm wide, ratio length/width 5; tibia 1.72 mm long, 0.16 mm wide, ratio length/width 10.75, tarsus 1.84 mm long; coxae, trochanters and trochantelli not preserved; tarsal claws with a preapical small tooth.

Fore wing hyaline, 3.28 mm long, 1.12 mm wide, ratio length/width 2.93; veins C and Sc+R distinct; pterostigma black, 0.52 mm long, 0.12 mm wide, ratio length/width 4.33; Rs separating from Sc+R distinctly basal of pterostigma (0.16 mm); basal part of Rs straight, relatively (0.28 mm) long before its fusion with M; basal part of vein M moderately curved, 0.24 mm long; base of M 0.96 mm from wing base, aligned with cross-vein 1cu-a; Rs+M rather long (0.4 mm long); distal free part of M, before its contact with 1m-cu short (0.08 mm long); distal free part of Rs, before its contact with r-rs longer (0.32 mm long); cross-vein 2r-m aligned with r-rs; 2m-cu ending in cell [2Rs]; 3r-m present but with its median part nebulous; distal part of vein Cu strongly bent posteriorly; 2cu-a very short (0.04 mm long), distinctly shorter than 1cu-a; 1m-cu ending in M distal of fork of Rs+M; 2m-cu well defined but with a short nebulous part near its anterior end; distal ends of

M and Cu well defined and reaching wing margin; no vein 2A+3A and cross-vein a; all cells [1M], [2M], [1R1], [1Rs], [2Rs], and [2R1] closed; [2R1] longer than other distal cells, 1.24 mm long, 0.32 mm wide, ratio length/width 3.87; [1R1] 0.72 mm long, 0.24 mm wide, ratio length/width 3.0; [1Rs] 0.32 mm long, 0.12 mm wide, ratio length/width 2.66; [2Rs] 0.36 mm long, 0.16 mm wide, ratio length/width 2.25; [1M] 0.6 mm long, 0.2 mm wide, ratio length/width 3.0; [2M] 0.68 mm long, 0.24 mm wide, ratio length/width 2.83.

Hind wing hyaline, 1.6 mm long, 0.56 mm wide, ratio length/width 2.85; distal part of Rs 0.36 mm long, not reaching wing margin; cross-vein r-m long and oblique, 0.12 mm long; distal parts of M and Cu reaching wing margin; a distinct angle between cu-a and vein A; distal part of vein A present but nebulous; vein cu-a 0.08 mm long; vein C present at wing base but only 0.2 mm long; no visible posterior lobe at wing base; five strong distal hamuli.

Metasoma pedunculate, very elongate, higher than broad, 3.44 mm long, 0.32 mm wide in broadest part, ratio length/width 10.75, sterna hard, convex; no metasomal armature on sternum II or III; spiracles not visible, probably absent, ovipositor very reduced, not surpassing apex of metasoma, sternite VII with two broad and long lobes, 0.4 mm long, 0.28 mm wide, ratio length/length of metasoma 0.11; cerci visible, 0.08 mm long.

#### 4. Discussion

If we follow the key to modern wasp families proposed in Mason (1993), *Albiogonalys* gen. n. falls in the Trigonalidae. There is still no real consensus on the phylogenetic relationships of the Trigonalidae. Among other papers, Dowton et al. (1997) considered the Trigonalidae, in their molecular phylogeny of the Apocrita, as the sister group of the Megalyridae, in the clade Evaniomorpha. Ronquist et al. (1999: figs 5–9) proposed several other phylogenetic hypotheses, depending on the method to obtain them. Lastly, Rasnitsyn (2002: fig. 331), following Rasnitsyn (1988), considered the Trigonalidae as a 'sister group' of a paraphyletic lineage Megalyridae (+ Stephanidae). He characterized the group Megalyridae+ Stephanidae+Trigonalidae by the hind wing vein A lost beyond cu-a. In *Albiogonalys* gen. n., the vein




Fig. 4. – *Albiogonalys elongatus* gen. n., sp. n., holotype ARC 205.1; reconstruction of wings: a. left fore wing, b. left hind wing. Scale bar represents 1

A, although nebulous, is not absent. Rasnitsyn (1988, 2002) also characterized the Trigonalidae by the following characters: (1) mandibular dentition asymmetrical (four teeth in right mandible and three in left, unlike three in Megalyridae). Carmean & Kimsey (1998) noted that this character is secondarily lost in three modern genera. The presence of four teeth in the right mandible of Albiogonalys gen. n. strongly suggests that it might be of the trigonalid type; (2) median mesoscutal sulcus absent, as in Albiogonalys gen. n. but also in the modern megalyrid genus Carminator SHAW 1988 (Shaw 1988: 107); (3) metasomal sterna hard, convex, as in Albiogonalys gen. n.; (4) ovipositor internalised, as in Albiogonalys gen. n. Furthermore, after the diagnosis of the Trigonalidae of Carmean & Kimsey (1998), Albiogonalys gen. n. has the following further derived characters: (5) presence of sparse white scales on the female flagellomeres ('feature unique to all members of this family'); (6) presence of a supra-antennal elevation. It also has the following other, plesiomorphic characters: (7) tarsi with plantar lobes; (8) fore wing with complete venation; (9) hind wing with two closed cells; (10) numerous antennomeres (25 in Albiogonalys gen. n.). Albiogonalys gen. n. has not the 'metasomal armature' proper to some modern Trigonalidae.

If we follow Shaw (1988) in his redefinition of the Megalyridae as a monophyletic group, *Albiogonalys* gen. n. does not fit in it because it has not the main apomorphic character of the family, i.e. hind wing venation greatly reduced. It can also be separated from this family after the following key characters (but some are either plesiomorphies or of more uncertain polarity): median mesoscutal sulcus present, long ovipositor, mandibles symmetrical, twelve antennal flagellomeres, fore wing venation reduced, broad metasoma, presence of a subantennal groove in the malar space. Thus there is no argument supporting a close relationship between *Albiogonalys* gen. n. and the modern Megalyridae.

Rasnitsyn (2002) considered the Megalyridae as a paraphyletic group. All the other Ceraphronoidea *sensu* Rasnitsyn and more especially the Stephanidae and Trigonalidae are supposed to be derived from it. We prefer to reject this confusing

point of view and to use monophyletic groups. In particular, Rasnitsyn (1975) created the Mesozoic subfamily Cleistogastrinae for several genera with complete fore wing venation very similar to those of the modern Trigonalidae. As Shaw (1988) already noted, the main body characters necessary for a family attribution are not preserved in all these fossils (bases of antennae, mandibles, thoracic ornamentation, structure of tarsi, structure of metasomal sterna). Thus, their attribution to the Megalyridae is not demonstrated. Nevertheless, many of these Cleistogastrinae have long ovipositors, unlike the modern Trigonalidae. Their trigonalid-like wing venation is not sufficient alone for their attribution to this family or its stem group. Their positions are uncertain. All these Cleistogastrinae can be easily distinguished from *Albiogonalys* gen. n. after their rounder metasoma.

Carmean & Kimsey (1998) separated the Trigonalidae in the two subfamilies Orthogonalinae CARMEAN & KIMSEY 1998 and Trigonalinae on the basis of the presence of tyloids on the male antennae as a synapomorphy of the former subfamily. These structures are not available in *Albiogonalys* gen. n. This fossil genus differs from *Orthogonalys* SCHULZ 1905, unique genus of the Orthogonalinae and most basal modern genus, in the absence of areolate-rugose propodeal sculpturing ornamentation. *Albiogonalys* gen. n. differs from all modern Trigonalidae in its very narrow elongate metasoma and the presence of a hind wing vein A (Schulz 1907; Yamane & Kojima 1982; Carmean & Kimsey 1998). This last character state is supposed to be plesiomorphic for the Trigonalidae after Rasnitsyn (2002), thus it would suggest a very basal position for *Albiogonalys* gen. n. in the trigonalid lineage.

# Position of the fossil taxa previously attributed to the Trigonalidae

The Mesozoic Cleistogastrinae, especially *Cleistogaster* RASNITSYN 1975 have a wing venation very similar to those of the modern Trigonalidae but they differ from this family in their very long ovipositor. As already indicated above, their position is uncertain. But their existence demonstrates that the wing venational characters are not sufficient for the attribution of a fossil taxon to the Trigonalidae *sensu stricto*.

The following taxa have been attributed to the Trigonalidae:

- Cockerell (1917) described *Trigonalys pervetus* and the same author (1920) listed the family 'Trigonalidae' in the Albian Burmese amber (recently dated as Albian by Cruickshank & Ko 2003). However Antropov (2000) transferred it in the Sphecidae.
- Statz (1938) described 'Trigonalys' bischofii from the Oligocene of Rott (Germany). The type of this species is stored in the Natural History Museum of Los Angeles County (Sphon 1973). Its wing venation is of the trigonalid-type but there is no body character supporting its attribution to the Trigonalidae. 'Trigonalys' bischofii differs from Albiogonalys gen. n. in the following characters of the fore




Fig. 5. – *Albiogonalys elongatus* gen. n., sp. n., holotype ARC 205.1; photography of ovipositor. Arrow shows the cerci. Scale bar represents 0,25 mm.

wing venation: cross-vein 2r-m reaching Rs well distal to the point of fusion between Rs and r-rs; point of separation between Rs and M is distal to anterior end of 1m-cu.

- Martínez-Delclòs et al. (1999) listed the family 'Trigonalidae' in the Aptian/Albian Spanish amber of Alava.
- Rasnitsyn (1977) described the fossil genus and species *Cretogonalys taimyricus* from the Cenomanian amber of the Taimyr Peninsula (Russia). He erected the fossil subfamily Cretogonalinae RASNITSYN 1977 for this taxon. *Cretogonalys* shows dissymmetric mandibles, as in modern Trigonalidae. It is the unique argument of Rasnitsyn to include it in this family, but its wing venation strongly differs from those of modern Trigonalidae. It also has areolate-rugose propodeal sculpturing ornamentation, as in 'primitive evaniomorph' and *Orthogonalys* (Carmean & Kimsey 1998: 53). *Cretogonalys taimyricus* strongly differs from *Albiogonalys* gen. n. in the following characters of the fore wing venation: cell [1M] very small, cell [1R1] very large; cell [2R1] small, not elongate; propodeum sculptured.
- Rasnitsyn (1986) described the fossil genus and species Darbigonalus capitatus from the Early Cretaceous of Gurvaneren Formation, West Mongolia. It is a very poorly preserved specimen with the body structures nearly useless and a bad preservation of the wing venation. It is an incertae sedis. Darbigonalus capitatus differs from Albiogonalys gen. n. in its rounded metasoma, but also in the shape of its fore wing cell [1R1].
- Rasnitsyn (1990: 189, pl. 15, fig. 1) described the fossil genus and species *Turgonalus minor* from the Cretaceous of Transbaikalia (Russia). It is based on a rather poorly preserved impression, with no detail in the body structure and a wing venation in bad condition.




Fig. 6. – *Albiogonalys elongatus* gen. n., sp. n., holotype ARC 205.1; photography of head and mesosoma. Scale bar represents 1 mm.

 Rasnitsyn et al. (1998) described an isolated fore wing from the Lower Barremian of England under the name *Turg-onalus cooperi* RASNITSYN & JARZEMBOWSKI, 1998. As indicated above, if this wing venation is of trigonalid-type, it is not sufficient for an attribution to this family *sensu stricto*.

The genus *Turgonalus* can only be considered as an *incertae* sedis that could be related to the Trigonalidae. It differs from *Albiogonalys* gen. n. in its rounded metasoma and fore wing venation

In conclusion, except maybe for *Cretogonalys taimyricus*, all the fossil taxa that are currently attributed to the Trigonalidae are of uncertain familial affinities. *Albiogonalys* gen. n. is thus the first and oldest accurate fossil representative of this family.

#### 5. Palaeoecological remarks

Many modern Trigonalidae have a very particular biology, laying their eggs on foliage. These eggs have to be consumed by the host to develop. The reduction of the ovipositor and the metasomal armature are related to this behaviour. However, none of the known hosts, except for sawflies, can be considered ancient. The most widely distributed trigonalid, Taeniogonalos, is a relatively derived genus known to directly parasite sawflies as well as indirectly tachinids and ichneumonids (Weinstein & Austin 1991). The least derived trigonalid, Orthogonalys, is widely distributed, absent only from Europe and Australia. Host information for the genus is limited to few rearings from Lepidoptera pupae that are believed to be intermediate hosts and a tachinid parasitoid of Lepidoptera. Considering its great antiquity and the lack of metasomal armature, Albiogonalys gen. n. was probably still a direct parasitoid, with sawflies as possible hosts.

Modern Trigonalidae have a nearly cosmopolitan distribution, with the exception of arctic and alpine regions. However, most taxa occur in the intertropical areas, under warm climates, and only one species is found in Europe. This intertropical distribution is congruent, for *Albiogonalys* gen. n., with the warm palaeoclimate generally admitted in Europe for the Mid-Cretaceous period (Frakes 1979; Jenkyns et al. 1994), as well as with the global temporary view shown by both the arthropod assemblage found in the amber and by the associated palynological remains identified in the outcrop of Archingeay-Les Nouillers thus far (Perrichot, in press; Perrichot et al., in press, 2002; Nel et al., in press; Azar et al. 2003; Néraudeau et al. 2002). The distribution of modern trigonalid genera linking widely separated regions appears to be relictual and supports evidence for the ancient origin of the group, at least Upper Jurassic or Lower Cretaceous, as shown by the present discovery of *Albiogonalys* gen. n.

#### Acknowledgements

The authors would like to acknowledge Thierry Guyot, collector of the specimen reported here, who generously gave us his collection of amber lumps containing fossil inclusions from Archingeay. We also thank François Locu for drawing the specimen and two anonymous referees for their comments. This paper is a contribution to the 'ECLIPSE' CNRS program: 'Interactions Climat/Ecosystèmes entre l'Aptien et lePaléocène'.

#### REFERENCES

- ANTROPOV, A.V. 2000: Digger wasps (Hymenoptera, Sphecidae) in Burmese amber. Bull. Nat. Hist. Mus., London, (Geol.) 56 (1), 59–77.
- AZAR, D., PERRICHOT, V., NÉRAUDEAU, D. & NEL, A. 2003: New psychodids from the Cretaceous ambers of Lebanon and France, with a discussion of *Eophlebotomus connectens* Cockerell, 1920 (Diptera, Psychodidae). Annals of the Entomological Society of America 96 (2), 117–126.
- CARMEAN, D. & KIMSEY, L. 1998: Phylogenetic revision of the parasitoid wasp family Trigonalidae (Hymenoptera). Syst. Entomol. 23 (1), 35–76.
- COCKERELL, T.D.A. 1917: Descriptions of fossil insects. Proc. Biol. Soc. Wash. 30, 79–82.
- 1920. Fossil arthropods in the British Museum. 4. Ann. Mag. Nat. Hist. (9) 6, 211–214.
- CRUICKSHANK, R.D. & Ko, K. 2003: Geology of an amber locality in the Hukawng Valley, Northern Myanmar. J. Asian Earth Sci. 21, 441–455.
- DOWTON, M., AUSTIN, A.D., DILLON, N. & BARTOWSKY, E. 1997: Molecular phylogeny of the apocritan wasps: the Proctotrupomorpha and Evaniomorpha. Syst. Entomol. 22 (3), 245–255.
- FRAKES, L.A. 1979: The Cretaceous period. In: Climates throughout geological time (Ed. by L.A. FRAKES). Amsterdam: Elsevier Scientific Publishing Company, 168–178.
- HUBER, J. T. & SHARKEY, M.J. 1993: Chapter 3. Structure. In: Hymenoptera of the World: an identification guide to families (Ed. by GOULET, H. & HUBER, J.T.). Research Branch Agriculture Canada Publication, Ottawa, 13–59.
- JENKYNS, H.C., GALE, A.S. & CORFIELD, R.M. 1994: Carbon- and oxygen-isotope stratigraphy of the English Chalk and Italian Scaglia and its palaeoclimatic significance. Geol. Mag. 131 (1), 1–34.
- MARTÍNEZ-DELCLÒS, X., PEÑALVER-MOLLA, E. & RASNITSYN, A. 1999: Hymenopteran insects from the Lower Cretaceous amber of Alava (Spain). Abstracts of the 7th International Symposium on Mesozoic Terrestrial Ecosystems, Buenos Aires, 42.
- MASON, W.R.M. 1993: Chapter 5. Key to superfamilies of Hymenoptera. In: Hymenoptera of the World: an identification guide to families (Ed. by GOULET, H. & HUBER, J.T.). Research Branch Agriculture Canada Publication, Ottawa, 65–100.

- NEL, A., PERRICHOT, V., DAUGERON, C. & NÉRAUDEAU, D. In press: A new Microphorites in the Lower Cretaceous amber of the Southwest of France (Insecta: Diptera: Dolichopodidae: 'Microphorinae'). Zoological Journal of the Linnean Society.
- NÉRAUDEAU D., PERRICHOT V., DEJAX J., MASURE E., NEL A., PHILIPPE M., MOREAU P., GUILLOCHEAU F. & GUYOT T., 2002: Un nouveau gisement à ambre insectifère et à végétaux (Albien terminal probable): Archingeay (Charente-Maritime, France). Geobios 35, 233–240.
- PERRICHOT, V. In press: The litter fauna: a distinctive feature of the Cretaceous amber of southwest France. Geologica Acta.
- Perrichot, V., Azar, D., Néraudeau, D. & Nel, A. In press: New Psocoptera in the Lower Cretaceous ambers of Lebanon and southwestern France (Insecta: Psocoptera: Trogiomorpha). Geological Magazine.
- Perrichot, V., Néraudeau, D., Azar, D., Menier, J.J. & Nel., A. 2002: A new genus and species of fossil mole cricket in the Lower Cretaceous amber of Charente-Maritime, SW France (Insecta: Orthoptera: Gryllotalpidae). Cretaceous Res. 23, 307–314.
- RASNITSYN, A.P. 1975: Vysshie pereponchatokrylye mezozoya [Hymenoptera Apocrita of Mesozoic]. Trudy Paleontologicheskogo Instituta Akademii nauk SSSR, Moscow, 147, 1–134.
- 1977: New Hymenoptera from the Jurassic and Cretaceous of Asia. Paleont. Zh. 3, 98–108. [in Russian, English translation in 1978, Paleont. J. 11 (3), 349–357].
- 1986: [Insecta Vespida (= Hymenoptera)],. In: Nasekomye v rannemelovykh ekosistemakh Zapadnoy Mongolii [Insects in the Early Cretaceous ecosystems of the West Mongolia] [in Russian] (Ed. by TATARINOV, L.P., AFANASEYVA, G.A., BARSBOLD, R., MOROSOWA, I.P., NOVITSKAYA, L.I., RESHETOV, V.Y., ROSANOV, A.Y., SYSOEV, V.A., TROFIMOV, B.A. & RASNITSYN, A.P.) Trudy Sovmestnaya Sovetsko-Mongol'skaya Paleontologicheskaya Ekspeditsiya, Moscow, 28, 154–164...
- 1988: An outline of evolution of the hymenopterous insects (order Vespida). Oriental Insects 22, 115–145.
- 1990: [Hymenoptera]. In: Pozdnemezozoyskie nasekomye Vostochnogo Zabaykal'ya [Late Mesozoic insects of Eastern Transbaikalia.] [in Russian]. (Ed. by RASNITSYN, A.P.). Trudy Paleontologicheskogo Instituta Akademii nauk SSSR, Moscow, 239, 177–205.
- 2002: Superorder Vespidea Laicharting, 1781. Order Hymenoptera Linné, 1758 (= Vespida Laicharting, 1781). In: History of insects (Ed. by RASNIT-SYN, A.P. & QUICKE, D.L.J.). Kluwer Academic Publishers, Dordrecht, Boston, London, 242–254.
- RASNITSYN, A.P., JARZEMBOWSKI, E.A. & ROSS, A.J. 1998: Wasps (Insecta: Vespida = Hymenoptera) from the Purbeck and Wealden (Lower Cretaceous) of southern England and their biostratigraphical and palaeoenvironmental significance. Cretaceous Res. 19 (3-4), 329–391.
- RONQUIST, F., RASNITSYN, A.P., ROY, A., ERIKSSON, K. & LINDGREN, M. 1999: Phylogeny of the Hymenoptera: a cladistic reanalysis of Rasnitsyn's (1988) data. Zool. Scr. 28 (1–2), 13–50.
- SCHULZ, W.A. 1907: Hymenoptera fam. Trigonalidae. In: Genera Insectorum (Ed. by Wystman, P) 61, 1–24.
- SHAW, S.R. 1988: Carminator, a new genus of Megalyridae (Hymenoptera) from the Oriental and Australian regions, with a commentary on the definition of the family. Syst. Entomol. 13 (1), 101–113.
- SPHON, G.G. 1973: Additional type specimens of fossil Invertebrata in the collections of the Natural History Museum of Los Angeles County. Contribution in Science, Los Angeles County Museum 250, 1–68.
- STATZ, G. 1938: Neue Funde parasitischer Hymenopteren aus dem Tertiär von Rott am Siebengebirge. Decheniana (A) 98 (1), 71–154.
- Weinstein, P. & Austin, A.D. 1991: The host-relationships of trigonalyid wasps (Hymenoptera: Trigonalyidae), with a review of their biology and catalogue to world species. Journal of Natural History 18, 209–214.
- YAMANE, S. & KOJIMA, J. 1982: *Pseudonomadina* new genus with a description of a new species from the Philippines (Hymenoptera, Trigonalidae). Kontyû 50 (2), 183–188.

Manuscript received May 4, 2003 Revision accepted October 10, 2003