

Zeitschrift:	Eclogae Geologicae Helvetiae
Herausgeber:	Schweizerische Geologische Gesellschaft
Band:	99 (2006)
Heft:	2
Artikel:	Echinoids from the Belpberg Beds (Obere Meeresmolasse, Middle Burdigalian) in the area of Bern (Switzerland)
Autor:	Kroh, Andreas / Menkveld-Gfeller, Ursula
DOI:	https://doi.org/10.5169/seals-169234

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 30.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Echinoids from the Belpberg Beds (Obere Meeresmolasse, Middle Burdigalian) in the area of Bern (Switzerland)

ANDREAS KROH¹ & URSULA MENKVELD-GFELLER²

Key words: Echinoidea, Belpberg Beds, Upper Marine Molasse (OMM), Early Miocene, Switzerland, palaeobiogeography
Schlüsselwörter: Echinoidea, Belpberg Schichten, Obere Meeresmolasse (OMM), Unter-Miozän, Schweiz, Paläobiogeographie

ABSTRACT

The present paper deals with the echinoid fauna of the Belpberg Beds, a lithostratigraphic unit in the Upper Marine Molasse cropping out southeast of Bern (Switzerland). In this unit, echinoids, in contrast to mollusks, are rare and of low diversity. The scarcity of echinoids is related to the environmental setting dominated by monotonous siliciclastic deposits and strong diagenetic alteration.

The echinoid fauna consists of three species: *Psammechinus dubius dubius*, *Echinocardium deikei* and *Hemipatagus ocellatus?* *Psammechinus dubius dubius* and *H. ocellatus* typically occur in Burdigalian sediments in the Paratethys and help to confirm the dating obtained from the microfossils and molluscs of the Belpberg Beds. Furthermore, the Belpberg echinoid fauna is especially relevant for the palaeobiogeography with species common to both the Rhône Basin and the Central Paratethyan faunas.

ZUSAMMENFASSUNG

Die vorliegende Arbeit beschäftigt sich mit der Echinidenfauna der Belpberg Schichten (Obere Meeresmolasse) bei Bern. Im Gegensatz zu den Mollusken sind die Echiniden selten und wenig divers. Dies kann einerseits auf das sedimentäre Regime, und andererseits auf diagenetische Ursachen zurückgeführt werden.

Insgesamt konnten drei Arten nachgewiesen werden (*Psammechinus dubius dubius*, *Echinocardium deikei* und *Hemipatagus ocellatus?*). Zwei davon treten in der Paratethys charakteristischerweise im Burdigalum auf und lassen ein ebensolches Alter für die Belpberg Schichten als wahrscheinlich erscheinen. Aus paläobiogeographischer Sicht ist die Belpberg-Echinidenfauna von besonderem Interesse, das sie ein Bindeglied zwischen Rhône Becken und Zentraler Paratethys darstellt.

1. Introduction

Apart from the early papers by Agassiz (1839, 1840) and de Loriol (1875–1876), the Neogene echinoid fauna of Switzerland and the Western Paratethys is poorly known. Additionally, a few general papers on the fauna of the Swiss Molasse mention echinoderm remains (Studer 1825; Heer 1865; Bachmann 1867, 1868; Kaufmann 1872; Kissling 1890; Thalmann 1923; Rutsch 1928). The only modern work deals with Pliocene echinoids from southern Switzerland (Ragaini 1996). In contrast, the echinoid faunas of the adjacent regions, the Rhône Basin and the Central Paratethys, have been recently the focus of renewed research (Philippe 1998; Kroh 2005). The echinoid fauna of the Belpberg Beds has never been studied in detail before. The echinoid collections of the Naturhistorisches Museum Bern give us this opportunity.

2. Geological setting

The Upper Marine Molasse (OMM, from the German term “Obere Meeresmolasse”) of the study area in the vicinity of Bern (Switzerland) was deposited during the Early Miocene in the perialpine Molasse Basin (Pfister & Wegmüller 1994). The special position of this area, represents a transitional zone between the Central Paratethys in the East and the Rhône Basin and the Mediterranean in the West, reflected by its mollusk (e.g. Pfister & Wegmüller 1994, 1998, 1999, 2000, 2001) and echinoid fauna (this study). Elements of both regions are represented.

Since Ritter (1742), the Belpberg Beds have been the focus of several palaeontological and geological publications. Studer (1825), Bachmann (1867) and Kissling (1890) described the fossil fauna around Bern. The work of Rutsch (1928) was for a

¹ Naturhistorisches Museum Wien, Geologisch-Paläontologische Abteilung, Burgring 7, A-1010 Wien, Österreich. E-mail: andreas.kroh@nhm-wien.ac.at

² Naturhistorisches Museum Bern, Abteilung Erdwissenschaften, Bernastrasse 15, CH-3005 Bern, Schweiz. E-mail: menkveld@nmbe.ch

	Niedermatt Beds (della Valle 1965)	
	Sädel Conglomerate (Rutsch 1928)	
Belpberg Beds sensu lato (Rutsch 1926)	Belpberg Beds sensu stricto (Rutsch & Schlüchter 1973)	Utzigen Beds (Gerber 1926) with Muschelsandstein (Rutsch 1926) and "Petrefactenlager" (Studer 1825) Upper Muschelsandstein (Gerber 1926) and/or Conglomerate with Oysters
	Bütschelbach Conglomerate (Rutsch 1967) and/or Lower Muschelsandstein (Gerber 1926)	
Sense Beds (Rutsch 1967)		

Fig. 1. Lithostratigraphic units and subunits of the Upper Marine Molasse in the study area (modified from Gruner 2001).




Fig. 2. Location and simplified geological map of the study area, indicating the different localities of the reported echinoids (modified from Pfister & Wegmüller 1994; Kellerhals et al. 1999).




Fig. 3. *Psammechinus dubius dubius* (AGASSIZ 1840). A) NMBE 5011426 in aboral view; B), C) NMBE 5011439 in oral and lateral view; D) NMBE 5011435, detail of the ambital tuberculation. All these specimens are from Hohburggraben (Middle Burdigalian, Belpberg, Switzerland). Scale bars: A-C (10 mm); D (5 mm).

long time the most comprehensive palaeontological and geological study of the Belpberg sediments. Keller (1989) and Schoepfer (1989) have published new results on sedimentological and stratigraphical aspects. More recently, Pfister & Wegmüller (1994, 1998, 1999, 2000, 2001) have published a series of papers on the mollusc fauna.

The sediments in the area have become known as the Belpberg Beds (Fig. 1). Initially, including a much wider range of lithostratigraphic units (Rutsch 1926), the beds have subsequently been restricted to the Upper Muschelsandstein and the Utzigen Beds (Rutsch & Schlüchter 1973). All the echinoid material originates from this restricted part of the Belpberg Beds, which crop out in the southeast of Bern (Fig. 2).

Gruner (2001) described the lithology of the Belpberg Beds s.s. as a fine- to medium grained sandstone, rich in mica, with intermediary marly layers and/or conglomerates. Rapid changes in facies are typical. The thickness of the Belpberg Beds s.s. ranges from 250 to 300 m and shows a tendency to decrease towards the northeast. Sedimentary interpretations by Schoepfer (1989) and Pfister & Wegmüller (1994) indicate a depositional environment within a storm influenced delta front, ranging into protected shallow-water prodelta bay environments.

Biostratigraphical correlations of the Belpberg Beds were given by Martini (1971), Rutsch (1971), Rutsch & Salaj (1974, 1980), Müller (1982), Schoepfer (1986), Schoepfer & Berger

(1989), Keller (1989), Demarq (1990) and Engesser (1990). Index forms are rare and age interpretations are controversial (Schoepfer 1989; Pfister & Wegmüller 1994). Strontium isotope measurements on pectinid shells from the Belpberg Beds (section Aarwald, Belpberg), in contrast, yielded an absolute age of 15.7 million years ago (Fischer 1985), correlating with the Early Langhian. However, these strontium data seem questionable in light of the strong diagenetic alteration of the material: aragonitic shells are preserved as moulds or casts only, pectinid and echinoid calcite shows a strong diagenetic overprint. Recently, Gruner (2001) assigned the Belpberg Beds to the Middle Burdigalian, based on conclusions of Schoepfer (1989) and Pfister & Wegmüller (1994). This is in agreement with earlier results obtained from calcareous nannoplankton samples, indicating an assignment from upper part of zone NN2 (equaling Early to Middle Burdigalian) to the border between zone NN3 and NN4 (correlating with the Middle Burdigalian; Schoepfer 1989; Demarq & Schoepfer 1990). In the Central Paratethys this interval corresponds to the Upper Eggenburgian to the Lower Ottangian (Schoepfer 1989; Pfister & Wegmüller 1994). See Harzhauser et al. (2003) for additional discussion on the age of the Belpberg Beds (and the "Helvetician", an outdated term used for the studied time interval) and Berger et al. (2005a, b) for an overview on the lithostratigraphy and palaeogeography of the Swiss Molasse.




Fig. 4. *Psammechinus dubius dubius* (AGASSIZ 1840). Drawing (using a camera lucida) of the apical disc of NMBE 5011427 from Hohburggraben (Middle Burdigalian, Belpberg, Switzerland). Poorly visible sutures are stippled, gonopore in genital plate 3 is not visible. Ocular plates are indicated by "oc" and their Lovenian number in Roman font. Genital plates are marked by "g" and their Lovenian number in Arabian font. Scale bar: 1 mm.

3. Material and methods

The material used in the present study is housed at the Naturhistorisches Museum Bern (NMBE). It originates from the collections of R. Hadorn, A. Hoerning, B. Hostettler, B. Schott, E. Wyttensbach and the excavations by the NMBE.

The described material originates from the upper part of the Belpberg Beds s.s. (Fig. 1). Most of the specimens come from the Hohburggraben section (Fig. 2E) and other localities at Belpberg. Additional material is derived from the localities Enggistein, Imihobel, Längenberg Seitenberg and Weinhalde (Fig. 2A-D, F, G).

The upper part of the Belpberg Beds s.s. (Fig. 1) is composed of a sequence of sandstones and marls. The sequence includes two conspicuous fossiliferous beds: the "Petrefactenlager" (fossiliferous horizons in the upper part of the sandstone and marl succession) in the middle part and the "Muschelsandstein" (coarse sandstone with bivalves, pebbles and shark teeth) at the top. Echinoids have only been found in the "Petrefactenlager". Only a single specimen (NMBE 5011432) can be assigned to a particular horizon (layer 13, Hohburggraben section), in the lower part of the succession of sandstones and marls. All the other sixteen specimens were collected loose from the uppermost layers of the sequence, or from fallen blocks belonging to these horizons.

During fieldwork, the investigated sections were repeatedly sampled to obtain washable material, but many horizons turned out to be too strongly lithified for washing and sieve analysis. The few samples that could be processed did not yield any echinoderm remain (J. Wegmüller, pers. comm.).

4. Systematic palaeontology

Phylum Echinodermata DE BRUGUIERE 1791 (ex KLEIN 1734)
Class Echinoidea LESKE 1778
Order Echinoida CLAUS 1876
Family Echinidae GRAY 1825
Genus *Psammechinus* AGASSIZ & DESOR 1846

Psammechinus dubius dubius (AGASSIZ 1840)

(Figs. 3A-D; Fig. 4)

- *1840 *Echinus dubius* AG. – AGASSIZ: 84–85; pl. 22, figs. 4–6
- 1875 *Psammechinus dubius*, AGASSIZ. – DE LORIOL: 29–30; pl. 2, figs. 6, 6a-c; 7, 7a, b
- 1883 *Psammechinus mirabilis*, NICOL. (*Echinometra*). – GUTZWILLER: 46
- 1998 *Psammechinus dubius* (AGASSIZ 1840) – PHILIPPE: 65–67; pl. 8, figs. 5–8 [cum syn.]
- 2005 *Psammechinus dubius dubius* (AGASSIZ 1840) – KROH: 25–26; pl. 14, figs. 1–4 [cum syn.]

Type material. – Figured in Agassiz (1840: pl. 22, figs. 4–6); current location unknown.

Type horizon and type locality. – ? Upper Marine Molasse, Middle Burdigalian, Chaux-de-Fonds, Switzerland.

Material. – 2 specimens from Cheergraben, Belpberg (NMBE 5014152, 5014155); 16 specimens from Hohburggraben, Belpberg, (NMBE 5011426–5011440, 5011442); 1 specimen from Seitenberg, near Niedermuhlern (NMBE 5011443); 1 specimen from between Seitenberg and Studweidholz, N Bütschelegg (NMBE 5011444).

Dimensions (in mm). –

Specimen no.	Diameter	Height	Remarks
NMBE 5011426	33.0	13.1	
NMBE 5011427	16.7	9.5	
NMBE 5011431	27.4	13.8	
NMBE 5011432	25.0	12.1	
NMBE 5011436	18.2	9.6	
NMBE 5011439	~28.0	12.9	deformed
NMBE 5011440	21.2	8.1	depressed
NMBE 5011442	19.5	8.9	
NMBE 5014152	28.0	19.0	
NMBE 5014155	27.0	6.0	

Description. – Test of small to medium size, with diameter ranging from 16.7 to 33 mm. The outline of the corona is more or less circular. The aboral surface is highly arched, the oral surface flattened and depressed around the peristome. The ambitus is tumid. The apical disc is preserved in specimen NMBE 5011427. It is dicyclic with clearly exsert ocular plates (Fig. 4). Ornamentation and/or tuberculation are obscured. The ambulacra consist of trigeminate plates belonging to the echinoid compound type. The pores are partitioned isopores and are arranged in arcs of three. Each plate bears one large noncrenulate, imperforate marginal tubercle and few smaller inner tubercles. The most perradially situated inner tubercle is distinctly enlarged. The interambulacra are about 1.5 times as




Fig. 5. *Echinocardium deikei* DESOR 1858. A) NMBe 5011449 in oral view, from Belpberg or Weinhalde (Tägertschi); B) NMBe 5011448 in aboral view, from Hohburggraben (Belpberg); C) NMBe 5011447 in aboral view, from Enggistein (Worbberg); D) NMBe 5011445 in oral view, collected between Vorder- and Hinterfultigen (Längenberg); E) NMBe 5011448, detail of ambulacrum III, from Hohburggraben (Belpberg). All the mentioned localities are from the Middle Burdigalian of Switzerland. Scale bars: A-D (10 mm); E (5 mm).

wide as the ambulacra at the ambitus. Each plate bears one noncrenulate, imperforate primary tubercle, lying slightly adorally of the centre of each plate. The secondary tuberculation is moderately dense. At the ambitus, two to four secondary tubercles are distinctly enlarged on each side of the primary tubercle. These secondary tubercles and the primary tubercles are arranged in more or less horizontal rows. The peristome is moderately large, being about 37 % of the test diameter. It has a circular outline and only faint buccal notches (formerly termed gill slits). The periproct is enclosed by the genital plates and has a circular outline. It is 2.1 mm in diameter in specimen NMBe 5011427.

Remarks. – Although poorly preserved, the specimens can be assigned to *Psammechinus dubius dubius* without doubt. All features correspond well to the original description. A thorough revision of closely related taxa was recently published by Philippe (1998). Reports of this subspecies from the Badenian (Middle Miocene) of the Central Paratethys (e.g. Mączyńska 1979, 1987, 1988) are doubtful and need to be re-examined (Kroh 2005).

Due to the strong diagenetic overprint, the skeletal material is strongly altered. This is expressed in the present specimens by a nearly complete loss of the stereom microstructure. Interestingly, the individual stereom types reacted differently to the alteration resulting in darker and lighter areas tracing the original structures. This accounts also for the dark horizon-

tal bands seen in many specimens (e.g. Fig. 3B, C), which might erroneously be interpreted as sutural gaps (in Fig. 3D, by contrast, sediment compaction led to partial crushing of the test and breaking of sutures).

Occurrence. – Paratethys: Eggenburgian (Early Burdigalian), Karpatian (Late Burdigalian), ? Badenian (Langhian-Serravallian); various localities in Austria and Hungary (Kroh 2005); Mediterranean: Aquitanian to Serravallian: circum-Mediterranean.

Order Spatangoidea CLAUS 1876
Family Loveniidae LAMBERT 1905
Genus *Echinocardium* GRAY 1825

Echinocardium deikei DESOR 1858

(Fig. 5A-D)

- *1858 [*Echinocardium*] *Deikei* DESOR nov. sp. – DESOR: 408
- 1875 *Echinocardium Deikei*, DESOR. – DE LORIOL: 122; pl. 22, fig. 6, 6a, b
- 1883 *Echinocardium Deikei*, DES. – GUTZWILLER: 46
- 1890 *Echinocardium Deikei* DESOR – KISSSLING: 68; pl. 1, fig. 1
- non1979 *Echinocardium deikei* DESOR 1857 – MĄCZYŃSKA: 33; pl. 9, figs. 2, 4 [= *E. peroni* fide PHILIPPE 1998]
- non1979 *Echinocardium deikei* DESOR 1857 – MĄCZYŃSKA: 33; pl. 9, fig. 3 [= deformed schizasterid]
- ?1985 *Echinocardium* cf. *deikei* DESOR 1858 – MIHÁLY: 245; pl. 5, fig. 6

Type material. – Reported to from the collection Deike according to Desor (1855–1858: 408), current location unknown.

Type horizon and type locality. – “Molasse de Krobell près St.-Gall” (Desor 1858: 408).

Material. – 2 specimens from between Vorder- and Hinterfultigen, Längenberg (NMBE 5011445, 5011446); 1 specimen from Enggistein, Worbberg (NMBE 5011447); 1 specimen from Hohburggraben, Belpberg (NMBE 5011448); 1 specimen from Belpberg or Weinalde, Tägertschi (NMBE 5011449).

Dimensions (in mm). –

Specimen no.	Length	Width	Height	Remarks
NMBE 5011445	31.1	22.5	12.9	slightly deformed
NMBE 5011446	23.3	23.8	26.0	deformed
NMBE 5011447	32.7	25.0	13.5	deformed
NMBE 5011448	41.6	34.0	21.9	posterior and left lateral parts missing
NMBE 5011449	34.9	37.6	15.2	deformed

Description. – Test of medium size with heart-shaped, antero-posteriorly elongated outline. A shallow frontal notch is developed. Aborally ambulacrum III and the adjacent interambulacral columns (2b and 3a) form a broad groove between the apical disc and the ambitus. Posterior end obliquely truncate with distinctly pointed subanal heel. In profile the test is slightly wedge shaped. The highest point lies posteriorly in interambulacrum 5. Apical disc is not preserved in any of the investigated specimens, but lies posteriorly, approximately 60 to 65 % of test length from the anterior margin. The ambulacra are petaloid and are moderately depressed on the aboral side between the apical disc and the ambitus. The paired ambulacra are greatly widened adapically. The frontal ambulacrum, in contrast, is narrower and more strongly depressed. Within the paired petals, the pores are large elongate isopores. In the frontal ambulacrum the pores seem to be minute partitioned isopores (poorly visible). The interambulacra are slightly inflated between the petals forming narrow, rounded keels. Tuberculation details are largely obscured due to poor preservation, but tubercles on the oral side and especially on the plastron are considerably larger than on the aboral surface. Fascioles are not preserved. The periproct lies high on the obliquely truncated posterior face of the test. It is subcircular with a diameter of 3.7 mm in NMBE 5011447. Peristome is not preserved.

Remarks. – Although the fascioles are not preserved, the material can clearly be assigned to the genus *Echinocardium* based on the overall shape and structure of the ambulacra. The studied specimens seem to be conspecific with *Echinocardium deikei* DESOR 1858 as re-described by de Loriol (1875: 122; pl. 22, fig. 6, 6a-b). Comparison with other species of *Echinocardium* from the Early Miocene of the Mediterranean and Paratethys is difficult due to the poor preservation of all known *E. deikei* specimens. *Echinocardium depressum* (AGASSIZ 1847) from the Burdigalian of the Rhône Basin differs from the present material by its subcentral apical disc, narrower ambulacrum III (respectively narrower groove) and more round-

ed outline (based on the re-description in Philippe 1998 and on material from the Early Burdigalian of the Central Paratethys). The relation between *E. deikei* and *E. depressum*, however, is a very close one and it may turn out that the two are conspecific upon a re-examination of the type material of *E. deikei*. *Echinocardium peroni* COTTEAU 1877 from the Burdigalian of the Rhône Basin and the Middle Miocene of Corsica differs by lacking a frontal notch, its rounded outline and its anteriorly eccentric apical disc (Cotteau 1877; Philippe 1998).

Echinocardium deikei DESOR 1857 from the Middle Miocene of Poland reported by Mączyńska (1979, 1991) was reassigned to *E. peroni* by Philippe (1998: 231).

Occurrence. – Paratethys: Obere Meeresmolasse (Middle Burdigalian) of Switzerland.

Echinocardium? sp.

(Fig. 6)

Material. – 1 fragment from the Belpberg Beds, Hohburg, Belpberg (NMBE 5011450).

Description. – Small (~ 22 x 20 mm) antero-lateral fragment of the aboral surface. The fragment preserves the aboral part of ambulacrum IV and interambulacral column 4a. The petal widens strongly adapically and seems to have been confluent with the posterior petal V. It bears widely spaced elongate isopores. Interambulacral column 4a shows traces of a dense and homogenous tuberculation. There are no traces of enlarged or camellate primary tubercles.

Remarks. – The present fragment is tentatively assigned to the genus *Echinocardium* based on the structure of the single preserved petal and the lack of enlarged or camellate primary tubercles in interambulacral column 4a. This genus is well documented by better preserved specimens from the locality Hohburg and other, nearby localities of the Belpberg Beds (this study).

Genus *Hemipatagus* DESOR 1858

Hemipatagus ocellatus? (DEFRANCE 1827)

(Fig. 7)

*1827 *Spatangus ocellatus* – DEFRANCE in BLAINVILLE et al.: vol. 50, 96-97 [not seen, fide PHILIPPE 1998]

1839 *Spatangus Nicoleti* AG. – AGASSIZ: 33; pl. 4, figs. 7; 8

1847 [*Spatangus*] *ocellatus* DEFR. – AGASSIZ & DESOR: 7

1858 [*Spatangus*] *ocellatus* DEFR. – DESOR: 422

non1864 *Spatangus ocellatus*, DEFRENCE. – WRIGHT: 487-488; pl. 21, fig. 1a, b [= *Lovenia duncani* GREGORY 1891 fide CHALLIS 1980]

1876 *Spatangus ocellatus*, DEFRENCE. – DE LORIOL: 132-134; pl. 23, figs. 2, 2a; 3

1915 *Hemipatagus ocellatus* DEFRENCE (*Spatangus*), 1827 – LAMBERT: 189-192; pl. 14, figs. 17; 18

1998 *Mareta ocellata* (DEFRENCE 1827) – PHILIPPE: 220-223; pl. 22, figs. 6; 7; 8a, b; 9a, b; 10a-c

2005 *Hemipatagus ocellatus* (DEFRENCE 1827) – KROH: 191-194; fig. 94; pl. 81, figs. 1-4; pl. 82, figs. 1; 2 [cum syn.]



Fig. 6. *Echinocardium?* sp. NMBE 5011450 in aboral view, from Hohburg (Middle Burdigalian, Belpberg, Switzerland). Scale bar: 10 mm.




Fig. 7. *Hemipatagus ocellatus?* (DEFRANCE 1827). NMBE 5011451, internal view of the aboral side, from Imihubel (Middle Burdigalian, Längenberg, Switzerland). Scale bar: 10 mm.

Type material. – Holotype unknown; Neotype designated by Lambert (1915: pl. 14, fig. 17); coll. Lambert, Muséum national, Paris (Philippe 1998: 220).

Type horizon and type locality. – Burdigalian of Saint-Paul-Trois-Châteaux, France (Philippe 1998: 220); Neotype from the Burdigalian of Taulignan.

Material. – 1 fragment from the Belpberg Beds, at Imihubel, Längenberg (NMBE 5011451).

Description. – Small (~15 x 43 mm) antero-lateral fragment of the aboral surface, exposed from the inside of the corona. Numerous camellate primary tubercles in interambulacra 3b and 4a are visible. The tubercles take up nearly the whole plate height and are extremely closely spaced, their areoles nearly touching each other. These tubercles extend from the adradial to the interradial suture both in interambulacra 3b and 4a. In the preserved fragment, there are at least 8 such tubercles in interambulacrum 3b and 16 in 4a. Additionally, the tip of petal IV is preserved. The petal is closed distally and bears large, conjugate, elongate isopores.

Remarks. – Although the specimen is a very small and poorly preserved fragment, it can be tentatively assigned to *Hemipatagus ocellatus* due to the presence and arrangement of the characteristic camellate primary tubercles and the shape

of the preserved petal. Another possible species, *Lovenia dun-canii* (GREGORY 1891), can be ruled out because of its less numerous and less densely arranged camellate tubercles. Moreover, the latter species seems confined to the Central Mediterranean region (Challis 1980). There is only a single record from the Southern Rhône Basin (Philippe 1998) and none from the Western and Central Paratethys. In contrast *Hemipatagus ocellatus* is a common member of the Burdigalian fauna of the Rhône Basin (Philippe 1998), the Western Paratethys (Agassiz 1839, under the name *Spatangus nicoleti*; de Loriol 1876) and the Central Paratethys (Schaffer 1912, under the name *Spatangus (Mareta) perornatus*; Kroh 2005). A possible attribution to *Spatangus*, another common genus in the Neogene of the Paratethys and the Mediterranean, can clearly be ruled out. *Spatangus* also has enlarged aboral primary tubercles (e.g. Néraudeau et al. 1998; Kroh 2005), but their structure is very different (Fig. 8). In *Hemipatagus ocellatus* and *Lovenia*, the aboral primary tubercles have deeply sunken areoles, clearly showing as bulbs on the inside of the corona (Fig. 8A). In *Spatangus*, on the contrary, the areoles are only slightly sunken and are not visible on the inside of the corona (Fig. 8B). In the present specimen, which is visible from the inside the *Hemipatagus/Lovenia*-type of tubercles is




Fig. 8. Schematic comparison between the enlarged aboral primary tubercles of loveniid type (A) and the spatangid type (B), in cross section. The fragment illustrated in Fig. 7 corresponds to the loveniid type, with pillar-like boss exposed through abrasion of the internal bulb.

clearly visible (Fig. 7). The innermost parts of the bulbs are partially eroded, exposing the pillar-like central part of the tubercles.

Earlier this species was assigned to the genus *Mareta* (e.g. Mortensen 1951), a member of the Spatangidae. However, Kroh (2005, in press) has shown that the type species of *Hemipatagus* as well as *H. ocellatus* are loveniids and not closely related to *Mareta*. *Hemipatagus* is structurally more closely related to *Lovenia* and it may often be difficult to distinguish from this genus on the base of poorly preserved material, a fact noted already by Mortensen (1951). This is notably expressed by the conspicuous adapical field of tubercles found in *Hemipatagus* that is invariably associated with an internal fasciole in all other known genera that share this feature. The presence of this specialized structure could indicate that an internal fasciole is initially present in juvenile *Hemipatagus*, but lost during ontogeny. Fasciole loss during ontogeny is not so uncommon in the Spatangoidea (e.g. Mortensen 1950, 1951; Jeffery 1998), and has caused considerable confusion when too much emphasis was placed in fasciole absence/presence in taxonomy. Other features include the depth of the frontal notch, the shape of the subanal fasciole, the structure of the phyllodes and primordial plates of the interambulacra and the constricted nature of the oral antero-lateral ambulacra.

Occurrence. – Paratethys: Burdigalian of the Swiss Molasse Basin and Early to Late Eggenburgian (Early Burdigalian) of the Austrian Molasse Basin; records from the Badenian (Langhian-Early Serravallian) of Hungary and Ro-

mania are highly questionable (Kroh 2005); Mediterranean: Burdigalian to ?Langhian of the Rhône Basin, Vence Basin and Sardinia.


5. Discussion and conclusions

5.1. Palaeoecology

The echinoid fauna of the Belpberg Beds is of low abundance and diversity. This is in strong contrast to the mollusc fauna which is both abundant and diverse (Pfister & Wegmüller 1994, 1998, 1999, 2000, 2001). The scarcity of the echinoderm material may also account for the lack of respective publications, with the exception of occasional notes in geological mapping reports (Studer 1825; Bachmann 1867; Kissling 1890; Thalmann 1923; Rutsch 1928) and the commonly mentioned ophiuroid resting traces. The latter were interpreted as asteroid body fossils or casts (e.g. as "*Astrerias aranciacea*" by Studer 1825 and as "*Astropecten helveticus*" by Mayer 1872 and Kaufmann 1872) until critically re-studied by Seilacher (1953). The strong diagenetic alteration and lithification of the sediment may in part explain the scarcity of echinoderm remains in the Belpberg Beds. Additionally, the sedimentary facies of the succession is rather monotonous, being dominated by fine grained, sandy sediments from a setting influenced by fluvial input (cf. 2. Geological setting). Such sediments facilitate only a reduced spectrum of echinoderms, which are much more common and diverse in biogenic carbonates or fully marine clays (Smith 1984). Moreover, the lack of washable sediment reduces the chances to find rare echinoderm groups (e.g. Donovan 2001), that can in most instances only be documented by bulk sample analysis. Cidaroid remains, for example, reported as rare by earlier authors (Bachmann 1867; Kissling 1890), were not recovered.

Extant sedimentary settings similar to those documented from the Belpberg Beds are inhabited by the same echinoid genera. *Echinocardium* and *Psammechinus*, for example, are very common in extant near-shore settings influenced by siliciclastic fluvial input (e.g. the German North Sea coast, Schäfer 1962; Krönke et al. 2001). Unfortunately, the majority of the reported specimens comes from scree and could not be related precisely to the horizons. It could not be established whether the species under discussion belong to a single assemblage or originate from different beds. The sediment adhering to the investigated specimens suggests the former, but provides no conclusive evidence due to the rather monotonous sedimentary succession.

Echinocardium is relatively rare in Miocene strata (Néraudeau et al. 2001; Kroh 2005), with the exception of the Burdigalian of the Southern Rhône Basin (Philippe 1998). It is usually found in infralittoral sands, often in association with echinaceans (e.g. *Psammechinus*, *Schizechinus*) and sometimes with *Spatangus* (Videt & Néraudeau 2002). In contrast *Psammechinus* is a generally found in Miocene shallow water echinoderm assemblages, both in siliciclastic and carbonate settings.



Species	Rhône Basin	Swiss Molasse						Central Paratethys
		Ⓐ Les Verrières, NE	Ⓑ Ste-Croix, VD	Ⓒ La Chaux-de-Fonds, NE	Ⓓ Crêt du Locle, NE	Ⓔ Berne Region	Ⓕ Aargau Region	Ⓖ St. Gallen Region
<i>Prionocidaris avenionensis</i> (DES MOULINS 1837)	6		3, 5	1, 3, 5				
<i>Cidaroida</i> indet.					2, 4			
<i>Arbacia catenata</i> (DESOR 1858)	6			5				7
<i>Psammechinus dubius dubius</i> (AGASSIZ 1840)	6	3, 5	3, 5	1, 3, 5	5	x		7
<i>Parascutella paulensis</i> (AGASSIZ 1841)	6					3	3	7
<i>Echinolampas scutiformis</i> (LESKE 1778)	6	3, 5		3, 5				
<i>Schizaster</i> sp.	6				5			
<i>Brissopsis crescentica</i> WRIGHT 1855	6	3, 5						
<i>Spatangus delphinus</i> DEFRENCE 1827	6	3, 5						
<i>Echinocardium deikei</i> (DESOR 1858)					x		3	
<i>Echinocardium depressum</i> (AGASSIZ 1847)	6			5	5			7
<i>Hemipatagus ocellatus</i> (DEFRENCE 1827)	6	3, 5		1, 3, 5		x		7

1 by AGASSIZ 1839, 1840; 2 by BACHMANN 1867; 3 by DE LORIOL 1875, 1876; 4 by KISLING 1890; 5 by LAMBERT 1925; 6 by PHILIPPE 1998; 7 by KROH 2005; x in the present paper; NE = Neuenburg; RG = Rhine Graben; VD = Vaud

Fig. 9. Comparison of the Belpberg echinoid fauna with that of the whole Swiss Molasse and its adjacent basins (Rhône Basin and Central Paratethys). The inset shows the position of the localities mentioned in the table and the general distribution of the marine Molasse sediments (in grey).

Various authors have commented on the potential environmental preferences of fossil species of *Psammechinus*, noting a possible association with detrital facies, coralline algal debris, seagrass meadows, and possibly reefal environments (Roman 1984; Roman & Lachkhem 1993; Blondel & Philippe 1992). *Psammechinus dubius dubius* is morphologically similar to the extant *P. miliaris*, the ecology of which is well known thanks to ecological and aquacultural studies (Kelly & Cook 2001 for a synthesis). *Psammechinus miliaris* is an omnivore, found in a variety of habitats from low tide level to deeper, sublittoral settings (down to 100 m depth). It is obviously not very selective in respect to the settled sediment, as it has been reported from boulders and rocky bottoms, as well as sands and mud. In many cases, *P. miliaris* is associated with brown algal patches, oyster banks and *Zostera* meadows.

Hemipatagus ocellatus is usually encountered in coarse sands and biotdetrites in shallow-water palaeoenvironments (Roman 1984; Kroh & Harzhauser 1999). It probably had similar ecological requirements as the members of its most closely related extant genus, *Lovenia*, which is a shallow burrowing detritivore living in sand or gravel bottoms in sheltered settings

from the low tide zone of the sublittoral (Miskelly 2002; Schultz 2005).

5.2. Biostratigraphy

From a biostratigraphic point of view, the dating indicated from the echinoids of the Belpberg Beds is consistent with that defined earlier based on microfossils, which indicated Early to Middle Burdigalian age (Schoepfer 1989). Based on the ranges of the studied taxa in the Western and Central Paratethys, the presence of *Psammechinus dubius dubius* and *Hemipatagus ocellatus*? indicates a late Early Burdigalian, respectively Late Eggenburgian age (cf. Kroh 2005). Although they have also been mentioned from younger strata of the Central Paratethys, these records were interpreted as doubtful notably because of misidentifications (Kroh 2005).

5.3. Palaeobiogeography

The Swiss Molasse Basin is a transitional area between the French Rhône Basin (westwards) and the Central Paratethys

(eastwards) (Fig.9). As such it is a key region for the understanding of the biogeographic relation of these two areas. The Belpberg echinoid fauna, despite its low diversity, includes forms common to both the Rhône Basin and the Central Paratethyan faunas. This is consistent with the echinoid fauna of the whole Swiss Molasse (Fig. 9). All species, except one (*E. deikei*), reported from the Swiss Molasse are also found in the Rhône Basin, and about 50 percent of the species are shared with the Eggenburg region of the Central Paratethys. This can be interpreted as a further evidence to support the theory of an echinoid immigration of the Late Eggenburgian (Early Miocene) Central Paratethyan forms via the Molasse trough (Kroh & Harzhauser 1999).

Acknowledgements

This study was supported by the Austrian Science Fund (FWF) via project no. P-13466-Bio to Werner E. Piller (Univ. Graz). Our sincere thanks go to Matthias Harzhauser and Oleg Mandic (both at the Naturhistorisches Museum Wien) for valuable discussion, to Thérèse Pfister (Naturhistorisches Museum Bern) for her support and providing material and to Alice Schumacher (Naturhistorisches Museum Wien) for taking photographs. The critical reviews and helpful comments of David A.T. Harper (Geologisk Museum, Copenhagen), James H. Nebelsick (Univ. Tübingen), Didier Néraudeau (Univ. Rennes), Roger Portell (Florida Museum of Natural History) and Jürg Wegmüller (Gunter) are gratefully acknowledged.

REFERENCES

- AGASSIZ, L. 1839: Description des Échinodermes fossiles de la Suisse: première partie, Spatangoides et Clypéasteroides. *Mém. Soc. helvét. Sci. nat.* 3, i-viii, 1–101.
- AGASSIZ, L. 1840: Description des Échinodermes fossiles de la Suisse: seconde partie. *Cidarides. Mém. Soc. helvét. Sci. nat.* 4, i-iv, 1–97.
- BACHMANN, I. 1867: Ueber die in der Umgebung Bern vorkommenden versteinerten Thierresten. 41 p., Weingart, Bern.
- BACHMANN, I. 1868: Über den Muschelstein in der Gegend von Reiden. *Mitt. naturf. Ges. Bern.* 1867, 247–260.
- BERGER, J.-P., REICHENBACHER, B., BECKER, D., GRIMM, M., GRIMM, K., PICOT, L., STORNI, A., PIRKENSEER, C., DERER, C. & SCHAEFER, A. 2005a: Paleogeography of the Upper Rhine Graben (URG) and the Swiss Molasse Basin (SMB) from the Eocene to Pliocene. *Int. J. Earth Sci. (Geol. Rundsch.)* 94, 697–710.
- BERGER, J.-P., REICHENBACHER, B., BECKER, D., GRIMM, M., GRIMM, K., PICOT, L., STORNI, A., PIRKENSEER, C., & SCHAEFER, A. 2005b: Eocene–Pliocene time scale and stratigraphy of the Upper Rhine Graben (URG) and the Swiss Molasse Basin (SMB). *Int. J. Earth Sci. (Geol. Rundsch.)* 94, 711–731.
- BLONDEL, T. & PHILIPPE, M. 1992: Les Echinides du Burdigalien terminal–Langhien de Tunisie. Données paléobiologiques et paléogéographiques. *Revue de Paléobiologie* 11, 433–446.
- CHALLIS, G.R. 1980: Palaeoecology and Taxonomy of Mid-Tertiary Maltese echinoids. 401 p., PhD Thesis, Bedford College, University of London (UK).
- COTTEAU, G.H. 1877: Description des Echinides. In: Locard, A. (Ed.): Description des Faunes des terrains Tertiaires moyen de la Corse. *Ann. Soc. Agric., Hist. Nat. Arts utiles Lyon* 1877, 227–335.
- DEMARO, G. 1990: Pectinidés néogènes: proposition d'échelle biostratigraphique pour la Méditerranée. *Geobios* 23, 149–159.
- DEMARO, G. & SCHOEPFER, P. 1990: Les Pectinidés du Miocène de Suisse occidentale. *Eclogae geol. Helv.* 83, 751–791.
- DESOR, E. 1855–1858: Synopsis des échinides fossiles. Lxviii + 490 p., Reinwald, Paris.
- DONOVAN, S.K. 2001: Evolution of the Caribbean echinoderms during the Cenozoic: moving towards a complete picture using all of the fossils. *Palaeogeogr., Palaeoclimatol., Palaeoecol.* 166, 177–191.
- ENGESSER, B. 1990: Die Eomyidae (Rodentia, Mammalia) der Molasse der Schweiz und Savoyens. Systematik und Biostratigraphie. *Schweiz. Paläont. Abb.* 112, 1–144.
- FISCHER, H. 1985: Strontium-Isotopen-Stratigraphie der Oberen Meeressmolasse (OMM): erste Resultate. *Schweiz. mineral. petrogr. Mitt.* 65, 360–361.
- GERBER, E. 1926: Ueber die Molasse im Amtsbezirk Bern, oder: Lassen sich aus den Molasseprofilen diluviale Krustenbewegungen nachweisen? *Mitt. natr. Ges. Bern* 1925, 38–58.
- GRUNER, U. 2001: Blatt 1167 Worb (Atlasblatt 104). Erläuterungen (mit Beiträgen von R. Burkhalter). *Geol. Atlas Schweiz* 1:25000, Erläut. 104, 1–51.
- HARZHAUSER, M., MANDIC, O. & ZUSCHIN, M. 2003: Changes in Paratethyan marine molluscs at the Early/Middle Miocene transition: diversity, palaeogeography and palaeoclimate. *Acta Geol. Pol.* 53, 323–339.
- HEER, O. 1865: Die Urwelt der Schweiz. 622 p., Schultheiß, Zürich.
- JEFFERY, C.H. 1998: Carrying on regardless: the echinoid genus *Cyclaster* at the Cretaceous-Tertiary boundary. *Lethaia* 31, 149–157.
- KAUFMANN, F.J. 1872: Rigi und Molassegebiet der Mittelschweiz. *Beitr. Geol. Karte Schweiz* 11, 1–474.
- KELLER, B. 1989: Fazies und Stratigraphie der Oberen Meeressmolasse (unteres Miozän) zwischen Napf und Bodensee. *Inauguraldiss. University of Bern*. (Switzerland), 403 p.
- KELLERHALS, P., HAEFELI, C. & RUTSCH, R.F. 1999: Blatt 1167 Worb. Topographie: Landeskarte der Schweiz 1:25000 (Atlasblatt 104). *Geol. Atlas Schweiz* 1:25000, Karte 104.
- KELLY, M. S. & COOK, E. J. 2001: The ecology of *Psammechinus miliaris*. In: Lawrence, J. M. (Ed.): *Edible Sea Urchins: Biology and Ecology. Developments in Aquaculture and Fisheries Science*, 217–224. Elsevier. Amsterdam.
- KISSLING, E. 1890: Die versteinerten Thier- und Pflanzenreste der Umgebung von Bern. *Excursionsbüchlein für Studirende* (sic). 70 p., Wyss, Bern.
- KROH, A. 2005: Catalogus Fossilium Austriae. Echinoidea Neogenica. Ivi + 210 p., Österr. Akad. Wiss., Wien.
- KROH, A. in press: *Hemipatagus* – a misinterpreted Loveniid (Echinodermata: Echinoidea). *J. of Syst. Palaeont.*
- KROH, A. & HARZHAUSER, M. 1999: An Echinoderm Fauna from the Lower Miocene of Austria: Paleoecology and Implications for Central Paratethys Paleobiogeography. *Ann. Naturhist. Mus. Wien* 101A, 145–191.
- KRÖNCKE, I., ZEISS, B. & RENSING, C. 2001: Long-term Variability in Macrofauna Species Composition off the Island of Norderney (East Frisia, Germany) in Relation to Changes in Climatic and Environmental Conditions. *Senckenbergiana maritima* 31, 65–82.
- LAMBERT, J. 1915: Description des échinides des terrains néogènes du bassin Rhône. *Mém. Soc. Paléont. Suisse* 41, 155–240.
- LAMBERT, J. 1925: Échinides de la molasse marine du Jura neuchâtelois. *Eclogae geol. Helv.*, 19 (1925–26), 250–254.
- LORIOL, P. DE 1875–1876: Echinologie Helvétique. Description des Échinides tertiaires de la Suisse. Troisième partie. Échinides de la Période Tertiaire. *Mém. Soc. Paléont. Suisse* 2 (1875): 1–88, pls. 1–8; 3 (1876): 89–142, pls. 9–23.
- MĄCZYŃSKA, S. 1979: Echinoids from the Miocene deposits of the Roztocze Region, South-eastern Poland. *Prace Muz. Ziemi* 32, 29–32.
- MĄCZYŃSKA, S. 1987: A supplementary account on the echinoids from the Kotryńska Basin (Middle Miocene; Holy Cross Mountains, Central Poland). *Acta Geol. Polonica*, 37, 145–153.
- MĄCZYŃSKA, S. 1988: Echinoids from the Middle Miocene (Badenian) sands from southern Poland. *Prace Muz. Ziemi* 40, 59–64.
- MĄCZYŃSKA, S. 1991: Characteristics of the Miocene echinoid remains from the exposure at Niechobrz near Rzeszów (In Polish). *Paleontologia a bathymetria - Materiały XIV Konferencji Paleontologów w Karpatach Rzeszowskich*, 36–37.
- MARTINI, E. 1971: Standard Tertiary and Quaternary calcareous nannoplankton zonation. In: Farinacci, A. & Matteucci, R. (Eds.): *Proceedings of the*

- II planktonic conference Roma 1970, vol. 2, 739–785, Tecnoscienza, Roma.
- MAYER, K. 1872: Systematisches Verzeichnis der Versteinerungen des Helvetian der Schweiz und Schwabens. Beitr. Geol. Karte Schweiz 11, 475–511.
- MISKELLY, A. 2002: Sea Urchins of Australia and the Indopacific. 180 p., Capricornica Publications, Sydney.
- MORTENSEN, T. 1950: A Monograph of the Echinoidea. V, 1. Spatangoida I. Protosternata, Meridosternata, Amphisternata I. Palaeoneustidae, Palaeostomatidae, Aëropsidae, Toxasteridae, Micrasteridae, Hemiasteridae. 432 p., C. A. Reitzel, Copenhagen.
- MORTENSEN, T. 1951: A Monograph of the Echinoidea. V, 2. Spatangoida II. Amphisternata II. Spatangidae, Loveniidae, Pericosmidae, Schizasteridae, Brissidae. 593 p., C.A. Reitzel, Copenhagen.
- MÜLLER, C. 1982: Nannoplankton der Schweizer Molasse. In: Jung, P. (coord.): Nouveaux résultats biostratigraphiques dans le bassin molasique, depuis le Vorarlberg jusqu'en Haute-Savoie. Doc. Lab. Géol. Fac. Sci. Lyon, H.S. 7, 35–38.
- NÉRAudeau, D., BORGHI, E. & ROMAN, J. 1998: Le genre d'échinide *Spatangus* dans les localités du Pliocène et du Pléistocène d'Émilie (Italie du Nord). Ann. Paléont. 84, 243–264.
- NÉRAudeau, D., GOUBERT, E., LACOUR, D. & ROUCHY, J.M. 2001: Changing biodiversity of Mediterranean irregular echinoids from the Messinian to Present-Day. Palaeogeogr., Palaeoclimatol., Palaeoecol. 175, 43–60.
- PFISTER, T. & WEGMÜLLER, U. 1994: Beschreibung, Vergleich und Verbretitung der Bivalven-Arten aus den Belpbergschichten (Obere Meeresmolasse, mittleres Burdigalien) in der Umgebung von Bern, Schweiz. 1. Teil: Palaeotaxodonta und Pteriomorphia, exklusive Ostreacea. Eclogae geol. Helv. 87, 895–973.
- PFISTER, T. & WEGMÜLLER, U. 1998: Beschreibung, Vergleich und Verbretitung der Bivalven-Arten aus den Belpbergschichten (Obere Meeresmolasse, mittleres Burdigalien) in der Umgebung von Bern, Schweiz. 2. Teil: Ostreacea, Heterodontia pro parte (Lucinacea, Chamacea, Carditacea und Cardiacea). Eclogae geol. Helv. 91, 457–491.
- PFISTER, T. & WEGMÜLLER, U. 1999: Beschreibung, Vergleich und Verbretitung der Bivalven-Arten aus den Belpbergschichten (Obere Meeresmolasse, mittleres Burdigalien) in der Umgebung von Bern, Schweiz. 3. Teil: Heterodontia pro parte (Mactracea, Solenacea und Tellinacea). Eclogae geol. Helv. 92, 395–449.
- PFISTER, T. & WEGMÜLLER, U. 2000: Beschreibung, Vergleich und Verbretitung der Bivalven-Arten aus den Belpbergschichten Obere Meeresmolasse, mittleres Burdigalien) in der Umgebung von Bern, Schweiz. 4. Teil: Veneracea. Eclogae geol. Helv. 93, 445–470.
- PFISTER, T. & WEGMÜLLER, U. 2001: Beschreibung, Vergleich und Verbretitung der Bivalven-Arten aus den Belpbergschichten (Obere Meeresmolasse, mittleres Burdigalien) in der Umgebung von Bern, Schweiz. 5. Teil: Heterodontia pro parte (Myacea, Hiatellacea, Pholadacea), Anomalodesmata (Pholadomyacea, Pandoracea, Clavagellacea), Nachtrag zu Palaeotaxodonta (Nuculacea, Nuculanacea), Pteriomorphia (Pectinacea) und Heterodontia (Carditacea, Cardiacea, Solenacea). Eclogae geol. Helv. 94, 399–426.
- PHILIPPE, M. 1998: Les Échinides miocènes du Bassin du Rhône: révision systématique. Nouv. Arch. Mus. Hist. Nat. Lyon 36, 3–241, 249–441.
- RAGAINI, L. 1996: L'echinofauna pliocenica del Canton Ticino nelle collezioni del Museo Cantonale di Storia Naturale di Lugano. Geol. Insubrica 1, 47–51.
- RITTER, J.J. 1742: De Patellite minimo et cucullato brevissimo. Acta Physico-Medica Academiae Caesareae Leopoldino-Carolinae Naturae Cursorum Exhibentia Ephemerides 6, 48–50.
- ROMAN, J. 1984: Les échinides et les milieux de vie du Tertiaire. Quelques exemples. Géobios, M.S. 8, 115–124.
- ROMAN, J. & LACKHEM, H. 1993: Le bassin sédimentaire de Melilla (Maroc septentrional) et sa faune d'échinoides du Messinien (Miocène supérieur). In: 118^e Congr. Nat. Soc. Hist. Scient., 4^e Coll. Géol. Afric. 445–460, Pau.
- RUTSCH, R.F. 1926: Zur Stratigraphie und Tektonik der Molasse südlich von Bern. Eclogae geol. Helv. 19, 673–678.
- RUTSCH, R.F. 1928: Geologie des Belpbergs. Beiträge zur Kenntnis der Stratigraphie, Palaeontologie und Tektonik der Molasse südlich von Bern. Separatabdr. aus den Mitt. natf. Ges. Bern 1927, 1–194.
- RUTSCH, R.F. 1967: Blätter 332 Neuenegg 333 Oberbalm 334 Schwarzenburg 335 Rüeggisberg (Siegfried-Atlas) (Atlasblatt 26). Erläuterungen. Geol. Atlas Schweiz 1:25000, Erläut. 26, 1–69.
- RUTSCH, R.F. 1971: Helvetian. G. Geol. Ann. Mus. geol. Bologna (2a) 37/2, 93–105.
- RUTSCH, R.F. & SALAJ, J. 1974: Eine Foraminiferenfauna aus dem Stratotyp des Helvétien (Mayer-Eymar 1858). Eclogae geol. Helv. 67, 431–434.
- RUTSCH, R.F. & SALAJ, J. 1980: Validité du Stratotype de l'Helvétien d'Imihubel [sic]. Actes du VI^e Colloque Africain de Micropaléontologie, Tunis 1974. Ann. Mines Géol. Tunis 28, 397–407.
- RUTSCH, R.F. & SCHLÜCHTER, C. 1973: Stratigraphische Gliederung der Molasse im bernischen Mittelland. Mitt. natf. Ges. Bern, N.F. 30, 86–90.
- SCHAFFER, F.X. 1912: Die Echiniden der Miocänbildung von Eggenberg. Abh. Geol. R.-A., 22, 187–192.
- SCHÄFER, W. 1962: Aktuo-Paläontologie nach Studien in der Nordsee. VIII + 666 p., Waldemar Kramer, Frankfurt am Main.
- SCHOEPFER, P. 1986: Sur l'âge de l'«Helvétien» du Guggisberg (BE): premières données micropaléontologiques. Eclogae geol. Helv. 79, 913–916.
- SCHOEPFER, P. 1989: Sédimentologie et stratigraphie de la Molasse Marine Supérieure entre le Gibloux et l'Aar. 211 p., PhD Thesis, University of Fribourg (Switzerland).
- SCHOEPFER, P. & BERGER, J.-P. 1989: Burdigalian and Helvetian in Western Switzerland. Geologica Carpathica 40, 17–21.
- SCHULTZ, H. 2005: Sea Urchins. 484 p., Heinke & Peter Schultz Partner, Hemdingen.
- SEILACHER, A. 1953: Studien zur Palichnologie, II. Die fossilen Ruhespuren (Cubichnia). N. Jb. Geol. Paläont. Abh. 98, 87–124.
- SMITH, A. B. 1984: Echinoid Palaeobiology. Special topics in palaeontology 1, x + 190 p., Allen & Unwin, London.
- STUDER, B. 1825: Beyträge zu einer Monographie der Molasse, oder Geognostische Untersuchungen über die Steinarten und Petrefacten, die zwischen den Alpen und dem Jura gefunden werden; mit besonderer Rücksicht auf den Kanton Bern und die angrenzenden Theile von Freyburg, Luzern und Solothurn. 427 p., Jenni, Bern.
- THALMANN, H. 1923: Die Fauna des Vindobonien vom Imihubel. Eclogae geol. Helv. 18, 366–371.
- VALLE, G. DELLA 1965: Geologische Untersuchungen in der miozänen Molasse des Blasenfluhgebietes (Emmental, Kt. Bern). Mitt. Natf. Ges. Bern, N.F. 22, 87–181.
- VIDET, B. & NÉRAudeau, D. 2002: Distribution paléoenvironmentale des huîtres dans le Messinien du bassin de Sorbas (Andalousie, SE Espagne). Ann. Paléont. 88, 147–166.
- WRIGHT, T.W. 1864: On the fossil Echinidae of Malta. With additional notes on the Miocene beds of the Island, and the stratigraphical distribution of the species therein; by A. Leith Adams. Quart. Journ. Geol. Soc. London 20, 471–491.

Manuscript received April 22, 2005

Revision accepted July 7, 2006

Published Online First October 5, 2006

