

**Zeitschrift:** Entomologica Basiliensia  
**Herausgeber:** Naturhistorisches Museum Basel, Entomologische Sammlungen  
**Band:** 5 (1980)

**Artikel:** Coleoptera: Fam. Cryptophagidae, genus Himascelis  
**Autor:** Gupta, T.Sen / Pal, T.K.  
**DOI:** <https://doi.org/10.5169/seals-980733>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 04.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Ergebnisse der Bhutan-Expedition 1972 des Naturhistorischen Museums in Basel

### Coleoptera: Fam. Cryptophagidae, Genus *Himascelis*

by T. Sen Gupta and T.K. Pal

Abstract: Ten specimens of the genus *Himascelis* Sen Gupta (Atomariinae) collected from Darjeeling district of West Bengal, Kashmir, Nepal and Bhutan are examined. *H. nigratus* Sen Gupta is synonymised with *H. brunneus* Sen Gupta and one species, *H. kashmirensis*, is described as new to science.

One of us (SEN GUPTA, 1978) erected the genus *Himascelis* for the two species *brunneus* and *nigratus*, from Himachal Pradesh and Darjeeling district of West Bengal respectively. He separated the species *nigratus* from *brunneus* by its blackish brown colour, smaller puncturation, antennal segment 3 slightly longer than segments 4–6 and the elytra being broadest across anterior one-third. In the present study we examined some more examples of this genus from Darjeeling, Kashmir Nepal and Bhutan including the male genitalia (Fig. 1). It is observed that the differences mentioned above between the species *nigratus* and *brunneus* are not constant. Moreover, the aedeagus, which was not studied earlier, is identical. Therefore, these two species have been merged here into one and *nigratus* is synonymised with *brunneus*.

#### 1. *Himascelis brunneus* Sen Gupta

*Himascelis brunneus* SEN GUPTA, 1978, Oriental Ins. 12 (2): 277–279 (Himachal Pradesh: India)

*Himascelis nigratus* SEN GUPTA, 1978, Oriental Ins. 12 (2): 279 (Darjeeling dist., W.B.: India) **Syn. nov.**

This species can be recognised by its prothorax being broad (1.00:1.77), lateral margin not uniformly curved, more than three times wider than front margin: elytra slightly elongate (1.19:1.00). Aedeagus (Fig. 1) inverted cucujoid-type, median lobe markedly elongated and its apex broadly pointed; articulated parameres well developed and moderately long. Length: 1.35–1.45 mm.

Specimens examined: India: West Bengal, Darjeeling district, Chimkhona (Ghum), 2200 m, 1 ex., 4. VI. 1975, W. Wittmer; Lebong, 1600/1800 m, 1 ex., 8. V. 1975, W. Wittmer; Kashmir, Daksum,

2400–2700 m, 2 ex., 9–13.VII.1976, W. Wittmer. Nepal: Thodung via Those, 3100 m, 1 ex., 29–31.V.1976, W. Wittmer & C. Baroni. Bhutan: 3100 m, 1 ex., 1.IX.1976, Dorjee Khandu; Dorjula, 3100 m, 3 ex., 2.IX.1976, Dorjee Khandu.

Distribution: India: West Bengal, Himachal Pradesh, Kashmir; Nepal; Bhutan.

## 2. *Himascelis kashmirensis* n. sp.

This species differs from *H. brunneus* by the base of its prothorax being less than two times wider than apex, prothorax less transverse (1:1.37) and elytra more elongated (1.41:1). General appearance (Fig. 2) elongate-oval, convex, reddish brown, rather shiny with moderately dense, short and recumbent pubescence.

Head short, narrowed in front, partly exposed with its posterior part concealed under prothorax and with a transverse angular ridge, clypeus broadly transverse and its apical margin rounded, vertex with




Fig. 1: Aedeagus of *Himascelis brunneus* Sen Gupta. Fig. 2: Dorsal view of *Himascelis kashmirensis* n. sp.

minute and sparse punctures and pubescence projected posteriorly, eyes minute and hardly visible from dorsal side. Antenna moderately long, antennal insertions dorsal and in small depressions; scape moderately large, pedicel and segment 3 about as long as scape and narrower, segments 4–9 short, subequal and about as broad as long or slightly transverse, segment 10 transverse and segment 11 about as broad as long.

Prothorax transverse (1.00:1.37), gradually widening posteriorly, front angles slightly projected and acute, hind angles slightly obtuse, front margin emarginate, lateral margins outwardly curved and basal margin bisinuate, pronotum convex and finely bordered, moderately, densely and uniformly punctured, width of each puncture narrower than space between the punctures, pubescence moderately dense and projected posteriorly.

Scutellum transversely triangular and impunctate.

Elytra broadly elongate (1.41:1.00), widest beyond middle, lateral margins uniformly curved, puncturation coarser than that of pronotum, pubescence moderately dense and projected posteriorly.

Measurements: Total length: 1.36 mm, length of antenna: 0.38 mm, width of prothorax across front margin: 0.26 mm and across base: 0.51 mm, length of elytra: 0.86 mm and width: 0.60 mm.

Holotype ♀, India: Kashmir: Daksum, 2400–2700 m, 9.–13. VII. 1976, W. Wittmer. (Nat. Hist. Mus., Basle).

### Acknowledgements

We express our sincere thanks to Dr. W. Wittmer of Naturhistorisches Museum Basle, who kindly provided us this interesting material for study. We are also indebted to the Director, Zoological Survey of India for providing laboratory facilities.

### Literature

SEN GUPTA, T. (1978): *A new genus of Atomariinae (Coleoptera: Cryptophagidae) with descriptions of two new species from India*. *Oriental Ins.* 12 (2): 275–279.

Authors' address:

Tapan Sen Gupta & Tarun K. Pal, Zoological Survey of India, 34, Chittaranjan Avenue Calcutta – 700 012 (India)