Zeitschrift:	Entomologica Basiliensia
Herausgeber:	Naturhistorisches Museum Basel, Entomologische Sammlungen
Band:	6 (1981)
Artikel:	A new genus of Malachiidae from Asia (Coleoptera)
Autor:	Mayor, A. / Wittmer, W.
DOI:	https://doi.org/10.5169/seals-980724

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 11.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

A new genus of Malachiidae from Asia (Coleoptera)

by A. Mayor and W. Wittmer

Abstract: The genus Anhomodactylus is proposed to accommodate 4 species of *Malachius* and one species of Anthocomus. The genus is distinguished from Malachius and Anthocomus by the asymetrically lobed foretarsi.

Recent studies involving a classification of the species of *Malachius* into subgenera have disclosed the need for generic reassignment of several Japanese, Chinese and Siberian species of *Malachius* and *Anthocomus*. These species can not be assigned to any kown genus of Malachiidae, consequently a new name is being proposed. This action is being taken in anticipation of the results of our subgeneric classification of *Malachius*.

Anhomodactylus n.gen. Type species: Malachius eximius Lewis,1895

Elongate, moderately graciliform to moderately robust. Surface feebly shining to dull, microreticulate. Color variable, typically dark green above, legs often pale. Pubescence short, inconspicuous; elytra typically with pale reclining setae intermixed with black suberect to erect setae.

Head (Figs 1 & 2) transversely sulcate between eyes in male, unmodified in female. Interocular margins (including vertex posterior to interocular margins in *M. biimpressifrons*) longitudinally sulcate in both sexes, sulcus more pronounced in male. Clypeus short, sclerotized in basal 1/4-1/3, membranous apically; epistomal suture tangential to antennal sockets. Antennae 11-segmented, filiform to subserrate.

Pronotum variable; elongate, as wide as long to transverse; angles broadly rounded; apex slightly to strongly produced; sides slightly, evenly produced or converging anteriorly; posterior angles slightly reflexed or not. Elytra elongate, completely covering abdomen or leaving terminal 1 to 2 terga exposed; apex unmodified in both sexes. Wings present, fully developed in both sexes. Tarsi 5-segmented in both sexes, foretarsi vaguely asymetrical in female; male foretarsi (Fig. 3a) distinctly asymetrically lobed; anterior surface of segments I



Figs 1–3: 1–2. Head, frontal aspect: 1, Anhomodactylus eximius δ . 2, A. ohbayashii \Im . 3. Protarsus, A. eximius δ : a, anterior aspect. b, dorsal aspect, segment V.

to IV elongate, elongation most pronounced in segments III and IV; foretarsal claws (Fig. 3b) asymetrical in both sexes, anterior claw greatly reduced in male, slightly reduced in female.

Abdominal sterna with posterior margin narrowly membranous, sterna typically completely sclerotized; terminal abdominal sternum of male shallowly emarginate, broadly rounded in female; pygidium broadly rounded to truncate in both sexes. Aedeagus of male genitalia with median lobe elongate, becoming gradually acute at apex; internal sac covered with numerous small, inconspicuous spines.

The following species are included in Anhomodactylus:

Malachius biimpressifrons PIC, 1932, Ent. Anz. 12: 252 (China) Malachius eximius LEWIS, 1895, Ann. Mag. N. H. (6) 16: 117, Fig.2 (Japan) Malachius gotoi NAKANE, 1956, Sci. Rep. Saikyo Univ. Ser. A, 2 (3): 31 or A 161 (Japan) Malachius ohbayashii WITTMER, 1953, Ent. Rev. Japan 6: 37 (Japan) Anthocomus albilabrus PIC, 1914, L'Echange 30: 73 (Amur)

M. biimpressifrons, M. ohbayashii and *A. albilabrus* were described from females. Their inclusion in *Anhomodactylus* is based primarily on the presence in these species of a shallow longitudinal sulcus at the interocular margins. *M. biimpressifrons* differs from the others in that the sulcus begins near the middle of the interocular margins and extends to the vertex posterior to the eye. It is tentatively included in *Anhomodactylus* pending discovery of the male. Males of *Anhomodactylus gotoi* and *A. eximius* have antennal segments III to V broader than the following segments and antero-posteriorly compressed. In the female of *A. ohbayashii* and presumably *A. gotoi* and *A. eximius* antennal segments III and IV are broader than the following segments, while in *A. albilabrus* and *A. biimpressifrons* segments III and IV are not broader than the following segments.

Adults of Anhomodactylus are not likely to be confused with those of other malachiid genera. They are distinctive in having the head transversely sulcate in males, and the interocular margins longitudinally sulcate and the fortarsi asymetrically lobed in both sexes. The relationship of Anhomodactylus to other malachiid genera is currently problematic. It does not appear to be closely related to Malachius or Anthocomus. Anhomodactylus shares no derived features with Anthocomus, a genus whose species typically have the male elytral apicies excavated and appendiculate. Although the head of some species of Malachius is sulcate, the sulcus typically occurs in the epistomal region rather than more posteriorly as it does in Anhomodacty*lus.* This difference in the position of the cephalic sulcus suggests an independent derivation of this character in the two genera. The modification of the foretarsi in *Anhomodactylus* appears to be unique and other derived characters are not helpful in indicating relationship.

Acknowledgments

Illustration of this study was made possible through a grant from the Swiss National Foundation for Scientific Investigations. We thank Mr. Armin Coray, biological illustrator, Basel, Switzerland for the drawings, and Dr. John D. Pinto, University of California, Riverside for valuable suggestions concerning the manuscript.

Authors' address: Dr. A. Mayor and Dr. W. Wittmer Naturhistorisches Museum Augustinergasse 2 CH-4001 Basel