Zeitschrift:	Entomologica Basiliensia
Herausgeber:	Naturhistorisches Museum Basel, Entomologische Sammlungen
Band:	11 (1986)
Artikel:	The influence of acidity and chlorinity on the distribution of Hydroporus species (Coleoptera, Dytiscidae) in the Netherlands
Autor:	Cuppen, J. G. M.
DOI:	https://doi.org/10.5169/seals-980555

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 10.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

11

The Influence of Acidity and Chlorinity on the Distribution of Hydroporus Species (Coleoptera, Dytiscidae) in the Netherlands.

by J.G.M. Cuppen

Abstract: 1. Adult water beetles were collected from 732 localities in the Netherlands. The distribution of eighteen species of the genus *Hydroporus* has been related to acidity and chlorinity by use of the Index of Representation (I. R.).

2. H. tristis (Payk.), H. obscurus Strm., H. pubescens (Gyll.), H. gyllenhalii Schiödte (= H. piceus Steph.), H. melanarius Strm. and H. neglectus Schaum are acidobiont species in the Netherlands and acidity forms a main environmental variable in their distribution. H. scalesianus Steph., H. memnonius Nicol. and H. nigrita (F.) are acidophilous species. H. erythrocephalus (L.), H. umbrosus (Gyll.) and H. incognitus Sharp are significantly overrepresented in acid waters but have a wide tolerance. H. striola Gyll. and H. dorsalis (F.) are clearly over-represented at pH's between 6.1 and 7.5 (not significant). The very common H. palustris (L.) and H. angustatus Strm. are over-represented at pH's between 6.6 and 7.5 (not significant for H. angustatus) and they avoid acid waters. H. tessellatus Drap. is the only alkaliphilous species. H. planus (F.) is indifferent with respect to acidity.

3. *H. tessellatus* is halophilous and chlorinity is an important environmental variable for this species. The remaining species with the exception of *H. memnonius*, *H. nigrita*, *H. palustris* and *H. planus* are haloxenous.

4. Acidity explains more variation in the distribution patterns of *Hydroporus* spp. in the Netherlands than chlorinity.

Key words: Coleoptera Dytiscidae – *Hydroporus* spp. – acidity – chlorinity – distribution.

Introduction

The holarctic genus *Hydroporus* Clairville (Coleoptera; Dytiscidae) is comprised of 213 species (FRANCISCOLO, 1979); it is the most diverse dytiscid genus in the Netherlands, with 24 species represented (VAN NIEUKERKEN, 1982). The species inhabit a wide variety of habitats (e. g. peat-cuttings, brackish waters and brooklets).

This investigation is part of a greater study concerning the ecology and geographical distribution of aquatic beetles in the Netherlands. In this study the distribution of eighteen species of *Hydroporus* was related to pH and chlorinity; the other six species – *H. discretus* Fairm., *H. elongatulus* Strm., *H. glabriusculus* Aubé, *H. longulus* Muls., *H. notatus* Strm. and *H. rufifrons* (Duft.) – were omitted due to lack of data. Except for *H. discretus* these species are extremely rare in the Netherlands and known from only single or a few localities (EVERTS, 1903; VAN NIEUKERKEN, 1979). Relationships between species, pH and chlorinity, based on field data, only can be of a correlative nature. Although causal relationships can not be determined by analysis of correlations between species and environmental factors, useful predictions concerning the distribution of species in a certain area can be obtained and information about the coexistence or non-co-existence of species.

Acidity and chlorinity were chosen as environmental variables because they can be determined easily at low costs. Furthermore, many of the ecological data concerning relationships between species and habitat refer to acidity (acidophilous) or chlorinity (haloxenous, halophilous, halobiont) or this relationship (tyrphophilous, tyrphobiont, sphagnobiont) is indirectly involved (HEBAUER, 1974).

Correlations between species and environmental variables only refer to the investigated area (in this study: the Netherlands), because a species can occur in different habitats or niches in different geographic areas due to interactions between environmental variables. This phenomenon is particularly noticeable at the boundary of the distribution of a species. For example, the distribution of *H. tessellatus* Drap. is centered in southern Europe (ZIMMERMANN, 1931; FRANCISCOLO, 1979) and it just reaches the south-western part of the Netherlands. In the Netherlands *H. tessellatus* is restricted to stagnant, brackish waters (this study). However, in southern Europe this species inhabits lotic and lentic waters and fresh as well as brackish waters (GUIGNOT, 1947; FRANCISCOLO, 1979).

Material and methods

Collecting methods

Adult water beetles were collected during the years 1978–1983 from 732 localities throughout the Netherlands, but mainly in the south-eastern part. Almost every available water type was sampled, although the number of samples collected from some water types (lakes and springs) was low.

The beetles usually were collected using a standard macrofauna-net (aparture 30×20 cm; mesh-size 0.5 mm), but in very shallow waters or in waters overgrown by emergent vegetation a kitchensieve (same mesh-size) was used. Species living at the water's edge were collected by trampling the borders in order to dislodge the beetles; floating speci-

mens then were caught by hand. Mosses were rinsed in a white tray and large stones were inspected visually. Collections were either sorted in the field using a white tray or were transported in a plastic bag to the laboratory, where samples were sieved and sorted in a white tray, as soon as possible.

Specimens were collected for about $1-1\frac{1}{2}$ h at most localities. Sometimes, however, a longer or shorter time was adopted, depending on abundance and diversity. Sampling was usually only qualitative, therefore, only presence/absence data for *Hydroporus* spp. were used.

Collections were made throughout the year but the number of observations from November to February was low.

Chemical measurements

Acidity was measured with a Metrohm Herisau (E588) pH meter, while chlorinity was measured titrimetrically according to Mohr. Analysis was conducted no later than 36 h after a collection was made. The values of pH and chlorinity were divided in classes (Table 1 and 2 respectively).

Analysis of distribution patterns

The distribution of *Hydroporus* spp. was related to pH and chlorinity using the Index of Representation (I.R.) (HILDREW & TOWNSEND, 1976).

 $I.R. = O - E/\sqrt{E}$

where O = number of observations of a certain species in a certain class of the factor considered, and E = expected number of observations.

The statistical significance was tested by the chi-squared test. Calculation of I.R. values is based on the null hypothesis (H_o) that a species has no preference or aversion towards certain classes of the factor considered and is represented in all classes equally. H_o was accepted when the differences between observed and expected number of observations was not sufficient to obtain chi-squared values above the 5% level. H_o was rejected when chi-squared values were higher than 5%.

When H_o is rejected, this indicates under- or over-representation in one or more classes of the considered factor. Positive I.R. values indicate over-representation (in this study: preference) and negative values indicate under-representation (in this study: aversion). Following Tolkamp (1980) differences in I.R. values are considered to be significant when the values deviate 2 or more from zero. The Index of Representation has been used in stead of frequency distributions, because the number of observations in different classes is not equal and can lead to incorrect interpretations (CUPPEN, 1983).

Results

Tables 1 and 2, respectively, give the number of observations for pH-classes and Cl-classes and the distribution of the number of observations of the *Hydroporus* spp. over these classes. These summaries of the field data are the data set for the calculations of the I.R. values. They show that *H. palustris* (L.) is the most commonly collected species of the genus over a wide range of pH and Cl values; it is present in more than 60% of the samples, while *H. tessellatus* is present in less than 2% of the collections. Most *Hydroporus* spp. show a wide range for both parameters, though the number of observations for pH higher than 7.5 and chlorinities higher than 200 mg l⁻¹ is low. *H. tessellatus* has only been found at pH higher than 6.5 and chlorinities higher than 100 mg l⁻¹.

	pН	3.1 - 4.0	4.1 - 5.0	5.1-6.0	6.1 - 6.5	6.6 - 7.0	7.1 - 7.5	7.6 - 8.0	>8.1
	pH-classes	1	2	3	4	5	6	7	8
	N	33	55	54	105	184	165	73	62
ł	TT	4	10	20	50	0.4	70	2.2	17
	H. angustatus	4	10	20	50	84	70	22	17
	H. dorsalis	1	2	5	14	21	14	2	1
	H. erythrocephalus	19	27	17	24	33	24	6	3
	H. gyllenhalii	15	21	10	5	2	1	-	1
	H. incognitus	11	23	13	25	19	10	-	2
	H. melanarius	12	7	13	7	2	-	-	-
	H. memnonius	5	12	18	27	22	10	4	8
	H. neglectus	9	12	16	8	5	2		-
	H. nigrita	2	3	9	19	20	13	5	3
	H. obscurus	8	7	5	-	1		-	—
	H. palustris	7	13	23	62	141	130	52	41
	H. planus	8	16	15	24	37	40	20	10
	H. pubescens	12	20	3	6	6	2	-	1
	H. scalesianus	3	7	6	13	10	5	1	1
	H. striola	3	1	8	17	27	21	5	3
	H. tessellatus	-	—	-	-	2	3	2	7
	H. tristis	20	30	12	8	11	13	1	3
	H. umbrosus	10	24	18	24	33	26	4	4

Tab. 1: Number of observations for pH-classes (N) and number of observations of Hydroporus spp.

Cl ⁻ (mg l ⁻¹) Cl ⁻ -classes N	<15 1 56	15-30 2 124	$30-45\\3\\139$	45-60 4 125	$60-75\ 5\ 92$	75–100 6 71	100–200 7 64	$>200 \\ 8 \\ 61$
H. angustatus	6	51	62	43	44	34	26	17
H. dorsalis	3	7	17	11	8	8	5.	_
H. erythrocephalus	38	38	23	10	23	12	7	2
H. gyllenhalii	19	12	12	6	3	3	_	_ *
H. incognitus	17	25	30	8	12	7	3	2
H. melanarius	5	14	11	6	3	1	1	_
H. memnonius	3 ′	24	28	19	9	5	8	10
H. neglectus	4	16	19	8	4	1	-	_
H. nigrita	2	16	20	10	11	6	3	6
H. obscurus	13	6	1		1	_	-	_
H. palustris	14	63	82	79	74	55	54	46
H. planus	13	27	26	27	24	12	17	24
H. pubescens	18	14	9	2	4	3		-
H. scalesianus	8	12	8	7	4	6	1	_
H. striola	2	9	21	17	17	11	5	3
H. tessellatus	-	-	_		_	-	2	12
H. tristis	31	25	14	8	12	7	-	
H. umbrosus	31	20	33	19	20	17	2	1

Tab. 2: Number of observations for Cl⁻-classes (N) and numbers of observations of Hydroporus spp.

In table 3, the I.R. values of the *Hydroporus* spp. with respect to pH are given. This table shows that most species have significant preferences and/or aversions to certain pH-classes. Two species have no significant I.R. values: *H. dorsalis* (F.) and *H. planus* (F.). For the last mentioned species the I.R. values do not deviate much from zero and one may call this species indifferent with respect to pH. The other species, including *H. dorsalis* show a logical arrangement of the I.R. values, though these values are not always significant. It means that acidity is one of the factors that influence the species composition in a certain water body. The importance of acidity as environmental variable for a species can be deduced from the deviations of the I.R. values from zero and the number of pH-classes between significantly positive and significantly negative values, that is the smaller this number the more important is acidity as environmental variable for that species.

On the basis of the I.R. values the species can be arranged in a way that they form a list from species mainly living in acid waters to species mainly living in alkaline waters. Acidity is an important environmental variable for species marked with a *.

J. G. M. Cuppen

pH	3.1 - 4.0	4.1-5.0	5.1-6.0	6.1 - 6.5	6.6 - 7.0	7.1 - 7.5	7.6-8.0	>8.1
H. angustatus H. dorsalis H. erythrocephalus	$\frac{-2.45}{-1.04}$ 4.60	$\frac{-2.45}{-1.18}$ 4.56	$-0.20 \\ 0.27 \\ 1.69$	$1.47 \\ 1.83 \\ 0.23$	$1.51 \\ 1.52 \\ -0.89$	$1.52 \\ 0.12 \\ -1.79$	$-1.18 \\ -1.63 \\ -2.37$	$-1.43 \\ -1.81 \\ -2.77$
H. gyllenhalii H. incognitus H. melanarius	$\frac{7.94}{2.94}$	$\frac{8.29}{5.48}$	$\frac{2.94}{1.95}$ 5.73	-1.03 $\frac{2.65}{0.46}$	$\frac{-3.18}{-1.36}$ -2.59	$\frac{-3.23}{-2.75}$ -3.04	$\frac{-2.34}{-3.21}$	-1.70 -2.28 -1.86
H. memnonius H. neglectus H. nigrite	$\frac{1.10}{0.10}$ $\frac{4.34}{0.74}$	$\frac{1.10}{1.42}$ $\frac{4.09}{1.09}$	$\frac{3.63}{6.20}$	$\frac{3.02}{0.19}$	-0.91 -2.24 0.22	-2.85 -2.84 -2.84	-2.02 -2.28 0.88	-0.33 -2.10 1.21
H. nigrita H. obscurus H. palustris	-0.74 $\frac{7.24}{-3.08}$	-1.09 -4.31 -3.75	$\frac{2.77}{-1.98}$	$\frac{2.57}{-1.74}$ -0.65	-1.86 2.11	$\frac{-0.91}{2.34}$	-0.88 -1.45 0.75	-1.31 -1.33 0.19
H. planus H. pubescens H. scalesianus	$ \begin{array}{c c} 0.12 \\ \underline{6.48} \\ 0.65 \end{array} $	$ \begin{array}{r} 0.90 \\ \underline{8.37} \\ \overline{1.90} \end{array} $	$0.69 \\ -0.36 \\ 1.41$	$-0.08 \\ -0.44 \\ 2.49$	$-0.88 \\ -1.85 \\ -0.46$	$ \begin{array}{r} 0.26 \\ -2.76 \\ -1.67 \end{array} $	$ \begin{array}{r} 0.73 \\ -2.23 \\ -1.68 \end{array} $	-1.16 -1.57 -1.47
H. striola H. tessellatus H. tristis	$ -0.43 \\ -0.79 \\ 7.40$	$\frac{-2.13}{-1.03}$ 8.33	$0.69 \\ -1.02 \\ 1.77$	$1.37 \\ -1.42 \\ -1.72$	$1.21 \\ -0.81 \\ -2.75$	$0.41 \\ -0.09 \\ -1.94$	$-1.20 \\ 0.51 \\ -2.81$	-1.57 5.33 -1.84
H. umbrosus	1.39	4.04	2.29	0.76	-0.50	-1.10	-2.72	-2.33

Tab. 3: The I.R. values for the Hydroporus spp. with respect to pH (significant values are underlined).

		"significant"	"significant"
		over-representation	under-representation
	* H. tristis	≤5.0	> 6.0
	* H. pubescens	≤ 5.0	> 6.5
Acidobiont	* H. obscurus	≤ 6.0	> 6.0
	* H. gyllenhalii	≤ 6.0	> 6.5
	* H. melanarius	≤ 6.0	> 6.5
	* H. neglectus	≤ 6.0	> 6.5
	H. scalesianus	4.1 - 6.5	> 7.0
Acidophilous	H. memnonius	5.1 - 6.5	7.1 - 8.0
	H. nigrita	5.1 - 6.5	not significant
	H. erythrocephalus	≤ 5.0	> 7.0
	H. umbrosus	≤ 6.0	>7.5
	H. incognitus	≤ 6.5	>7.0
	H. dorsalis	6.1 - 7.0	> 7.5
	H. striola	6.1 - 7.0	$\leq 5.0 > 7.5$
	H. angustatus	6.1 - 7.5	$\leq 5.0 > 8.1$
	H. palustris	6.6 - 7.5	≤6.0
Alkaliphilous	H. tessellatus	>7.5	≤6.5
Indifferent	H. planus		

H. pubescens (Gyll.), H. tristis (Payk.), H. obscurus Strm., H. gyllenhalii Schiödte (=H. piceus Steph.), H. melanarius Strm. and H. neglectus Schaum are considered acidobiont species, while H. scalesianus Steph., H. memnonius Nicol. and H. nigrita (F.) are acidophilous species. H. erythrocephalus (L.), H. umbrosus (Gyll.) and H. incognitus Sharp have a preference for acid waters, but a wide tolerance. H. angustatus Strm. and H. palustris (L.) are over-represented at pH's between 6.6 and 7.5 and they avoid waters with a pH below 5.0. H. dorsalis (F.) and H. striola Gyll. are over-represented at pH's between 6.1 and 7.0 and have a wide tolerance. H. tessellatus is an alkaliphilous species.

The I.R. values for the *Hydroporus* spp. with respect to chlorinity are given in table 4. No significant I.R. values have been obtained for H. memnonius, H. nigrita and H. striola, and only negative ones for H. angustatus and H. dorsalis. A comparison with table 3 shows that negative or positive I.R. values do not deviate much from zero, significant values most often do not deviate much from 2 and the number of Clclasses between significantly positive and significantly negative values is large. Four acidobiont species (H. obscurus, H. pubescens, H. gyllenhalii and H. tristis) and one acidophilous species (H. scalesianus) are significantly over-represented in waters very low in chlorinity $(<15 \text{ mg } l^{-1})$ and are significantly under-represented in waters with a chlorinity $> 100 \text{ mg} \text{ }^{-1}$. For *H. obscurus* the under-representation is not significant due to the low number of observations for this species. The other acidobiont species (H. melanarius and H. neglectus) have significant positive I.R. values for somewhat higher chlorinities (15-45 mg 1⁻¹), but the same significant aversion. *H. memnonius* and *H. nigrita*, both acidophilous species, have a non-significant over-representation for chlorinities between 15-45 mg l⁻¹. Their occurrence in waters with more than 200 mg Cl⁻/l⁻¹ is most interesting; this occurrence concerned a small number of ditches behind seadikes, where a permanent seepage of brackish water through the dikes exists. H. incognitus, H. erythrocephalus and H. umbrosus have significant preference for low chlorinities and significant aversion for chlorinities $>100 \text{ mg } l^{-1}$. H. palustris and H. angustatus only have significant aversions for very low chlorinities ($<15 \text{ mg l}^{-1}$). H. planus and H. tessellatus have a significant preference for chlorinities above 200 mg l⁻¹, but the tolerance for lower chlorinities is great in case of H. planus. H. tessellatus is the only halophilous species in the Netherlands. Summarizing, all species except H. memnonius, H. nigrita, H. palustris and H. planus are haloxenous and *H. tessellatus* is halophilous.

J. G. M. Cuppen

Cl ⁻ (mg l ⁻¹)	<15	15-30	30-45	45-60	60-75	75–100	100-200	>200
H. angustatus	-3.36	0.44	1.13	-0.77	1.41	1.25	0.25	-1.36
H. dorsalis	-0.71	-0.95	1.73	0.29	0.21	0.95	-0.07	-2.22
H. erythrocephalus	7.69	2.37	-1.12	-3.15	0.86	-0.74	-1.74	-3.01
H. gyllenhalii	7.21	0.88	0.48	-1.11	-1.49	-1.01	-2.19	-2.14
H. incognitus	3.21	1.76	2.31	-2.32	-0.30	-0.97	-2.02	-2.26
H. melanarius	1.05	2.68	1.15	-0.38	-0.95	-1.49	-1.36	-1.85
H. memnonius	-1.79	1.43	1.75	0.21	-1.18	-1.65	-0.42	0.39
H. neglectus	0.01	2.42	2.90	-0.29	-0.99	-1.80	-2.13	-2.08
H. nigrita	-1.54	0.98	1.59	-0.74	0.56	-0.44	-1.37	-0.07
H. obscurus	8.99	1.29	-1.50	-1.89	-1.01	-1.43	-1.35	-1.32
H. palustris	-3.63	-1.81	-0.71	-0.08	2.00	1.44	2.06	1.13
H. planus	-0.00	-0.33	-1.10	-0.38	0.57	-1.10	0.55	2.61
H. pubescens	7.25	1.90	-0.16	-2.24	-0.91	-0.83	-2.09	-2.04
H. scalesianus	2.39	1.51	-0.25	-0.30	-0.74	0.73	-1.51	-1.96
H. striola	-1.77	-1.42	1.21	0.65	1.93	0.96	-0.89	-1.53
H. tessellatus	-1.03	-1.54	-1.63	-1.55	-1.33	-1.16	0.70	10.03
H. tristis	8.66	2.11	-1.03	-2.10	-0.05	-0.78	-2.91	-2.84
H. umbrosus	6.06	$-\overline{0.86}$	1.12	-1.10	0.48	0.84	-2.97	-3.16

Tab. 4: The I.R. values for the Hydroporus spp. with respect to Cl (significant values are underlined).

Discussion

Knowledge of larval habitat requirements is necessary for a detailed description of the habitat(s) of a certain species (GALEWSKI, 1971), but, so few Hydroporus larvae have been described (NILSSON, 1982), that habitat(s) of the larvae cannot be determined for the moment. According to JACKSON (1952) adults of many water beetles are very mobile (good developed wings and wing musculature) and they form an unstable part of macrofauna-coenoses. However, the restriction of many *Hydroporus* spp. to certain habitats is so evident, that, in general, their great migratory capacity (e. g. GALEWSKI, 1971; DROST & SCHREIJER, 1978) must be doubted. If mobility is assumed, a mechanism for detection of required water types must exist or, perhaps, species fall prey to fish, when colonizing unsuitable habitats, as shown by MACAN (1976) for Corixidae. This means that larvae will be found only in habitats where the adults form reproducing populations, at least during part of the year. Only species with a great migratory capacity (H. planus, H. tessellatus) can form populations in other habitats than their breeding grounds.

HEBAUER (1974) provides ecological nomenclature of water beetles according to the following definitions:

Tyrphobiont species: restricted to habitats with peaty soils

Tyrphophilous species: mainly in habitats with peaty soils, but occasionally elsewhere

Acidophilous species: mainly living in weakly acid waters.

Two characteristics refer to the preferred soil as the main characteristic of the distribution pattern of a species and all three characteristics – though indirectly – refer to acidity. Tyrphobiont – in terms of acidity – can be translated into: restricted to very acid waters, and tyrphophilous into: mainly in very acid waters, but occasionally elsewhere (e. g. *H. erythrocephalus* and *H. incognitus* in this study). When this assumption is made a comparison for some species in different parts of their geographical distribution area can be made.

	Netherlands this study	Germany Hebauer (1974) Schaeflein (1971)	Sweden Nilsson (1979)
H. obscurus	acidobiont	tyrphobiont	tyrphophilous
H. tristis	acidobiont	tyrphophilous	tyrphophilous
H. melanarius	acidobiont	acidophilous	tyrphophilous
H. umbrosus	"tyrphophilous"	tyrphophilous	tyrphophilous
H. erythrocephalus	"tyrphophilous"	acidophilous	tyrphophilous
H. scalesianus	acidophilous	?	tyrphobiont
H. angustatus	not acidophilous	acidophilous	tyrphophilous

This comparison clearly shows that the pH preference of a number of species of *Hydroporus* varies geographically. A better comparison only will be possible when more exact definitions for soil conditions (i.e. definitions not subject to a personal assessment based on the presence/ absence of *Sphagnum*) are designed or when actual measurements of the acidity in other countries are available.

Concerning chlorinity, the data for the *Hydroporus* spp. in the Netherlands are not in contradiction with data from literature as most species are haloxenous or – more or less – salt tolerant. In the Netherlands, acidity better describes the distribution patterns of *Hydroporus* spp. than chlorinity.

Acknowledgement

I would like to thank Prof. Dr C. den Hartog for critical remarks and the improvement of the text.

References

- CUPPEN, J.G.M., (1983): On the habitats of three species of the genus Hygrotus Stephens (Coleoptera: Dytiscidae). Freshw. Biol. 13: 579–588.
- DROST, B. & M. SCHREIJER, (1978): Waterkevertabel. Jeugdbondsuitgeverij, Zeist & 's Graveland. 222 pp.

EVERTS, E., (1903): Coleoptera Neerlandica. De schildvleugelige insekten van Nederland en het omringende gebied. I. Martinus Nijhoff, 's Gravenhage.

FRANCISCOLO, M.E., (1979): Coleoptera Haliplidae, Hygrobiidae, Gyrinidae, Dytiscidae. Fauna d'Italia 14: 1–804.

GALEWSKI, K., (1971): A study on morphobiotic adaptations of European species of the Dytiscidae (Coleoptera). Polskie Pismo Ent. 41: 487–702.

GUIGNOT, F., (1947): Coléoptères Hydrocanthares. Faune de France 48: 1-286.

HEBAUER, F., (1974): Über die ökologische Nomenklatur wasserbewohnender Käferarten (Coleoptera). Nachr. Bl. Bayer. Entomol. 23(5): 87–92.

HILDREW, A.G. & C.R. TOWNSEND, (1976): The distribution of two predators and their prey in an iron rich stream. J. Anim. Ecol. 45: 41–57.

JACKSON, D.J., (1952): Observations on the capacity for flight of water beetles. Proc. Ent. Soc. London, A, 27: 57–70.

MACAN, T.T., (1976): A twenty-one-year study of the water-bugs in a moorland fishpond. J. Anim. Ecol. 45: 93–922.

NIEUKERKEN, E.J. VAN, (1979): Faunistische notities over enkele soorten van het genus Hydroporus Clairville in Nederland (Coleoptera: Dytiscidae). Ent. Ber. 39: 116–120.

NIEUKERKEN, E.J. VAN, (1982): Handleiding voor het projekt waterkevers (Coleoptera). Instrukties voor medewerkers EIS-Nederland 6: 1–28.

NILSSON, A.N., (1979): The dytiscid (Coleoptera: Dytiscidae) fauna of the province Västerbotten, northern Sweden. Fauna Norr. 10: 1–32.

NILSSON, A.N., (1982): A key to the larvae of the Fennoscandian Dytiscidae (Coleoptera). Fauna Norr. 2: 1–45.

Schaeflein, H., (1971): 4. Familie: Dytiscidae, echte Schwimmkäfer. In: Die Käfer Mitteleuropas (Freude, H., Harde, K. W. & Lohse, G. A., eds). 3: 16–89.

TOLKAMP, H.H., (1980): Organism-substrate relationships in lowland streams. Thesis, Centre for Agricultural Publishing and Documentation, Wageningen.

ZIMMERMANN, A., (1931): Monographie der paläarktischen Dytiscidae. II. Hydroporinae (2. Teil: Die Gattung Hydroporus Clairv.). Kol. Rdsch. 17: 97–159.

Author's address: Dr Jan G.M. Cuppen Vakgroep Waterzuivering sektie Hydrobiologie De Dreijen 12, LH Wageningen 6703 BC Wageningen, The Netherlands