**Zeitschrift:** Entomologica Basiliensia

Herausgeber: Naturhistorisches Museum Basel, Entomologische Sammlungen

**Band:** 16 (1993)

Artikel: Inhibition of ovicidal activities and fertility in a water beetle, Cybister

tripunctatus asiaticus Sharp (Coleoptera, Dytiscidae) following topical

application of Diflubenzene, a Chitin inhibitor

Autor: Sen, N. S. / Nagvi, A. H. / Rani, Vineeta

**DOI:** https://doi.org/10.5169/seals-980483

# Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

# **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

# Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF: 28.11.2025** 

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Entomologica Basiliensia	16	73-78	1993	ISSN 0253-2484
--------------------------	----	-------	------	----------------

# Inhibition of ovicidal activities and fertility in a water beetle, Cybister tripunctatus asiaticus Sharp (Coleoptera, Dytiscidae) following topical application of Diflubenzene, a Chitin inhibitor

by N. S. Sen, A. H. Naqvi & Vineeta Rani

Abstract: Topical application of a chitin inhibitor, Diflubenzene (Dimilin) to the aquatic beetle, *Cybister tripunctatus asiaticus* affected the reproductive potential of both male and female. Control female mated with control male laid a maximum of 450 eggs and 95 % of them hatched, whereas high concentration treated female mated with similar male oviposited only 102 eggs and the hatching efficiency was reduced to 22 % only. Regarding production and hatching of eggs, males were more significantly affected (p < 0.01 & 0.001) by the treatment than the females (p < 0.05 & 0.01). However weight-loss in eggs and newly hatched larvae was equally affected (p < 0.01) on both the sexes.

Key words: Coleoptera, Dytiscidae – *Cybister* – topical application – Diflubenzene – reduction – ovicidal activities – fertility – weight loss – correlation coefficient.

# Introduction

Ovicidal activities and fertility of adult insects have been reported to be adversely affected by topical application of organophosphorous compounds (HANIFFA et al., 1986 and HANIFFA & JOSE, 1987). Secondary plant substances (MATHAVAN et al., 1984) and chitin inhibitors (SEXENA & KUMAR, 1982, CHOCKALINGAM & NOORJE-HAN, 1984). Such compounds are extensively used in India on a wide range of insects, especially on majority of the crops, vegetables and stored-grain pests not only because of the low cost involvement but also for their lesser residual effects. The basic toxicity of these compounds is targeted through the central nervous system via inhibition of acetyl cholinesterases (PANT & KATIYAR, 1983). Despite of such existing knowledge, little is known about the inhibitory role played by Diflubenzene [1-(4-Chloropheny1)-3-(2-6-diflurobenzyl) Ureal, a chitin syntheses inhibitor (HOLST, 1975 & BUTCHI, 1978) on the physiology of reproduction in insect. The present authors attempt to show the impact of topical application of Diflubenzene, a chitin inhibitor, on the ovicidal activities and fecundity rate in an aquatic beetle, Cybister tripunctatus asiaticus Sharp, a fish pest (DUTTAMUNSHI et al., 1979).

# **Material and Methods**

Healthy adults of C. tri. asiaticus were collected from a nearby pond and were acclimed to the laboratory conditions of  $26.5 \pm 2.3$  °C and  $43 \pm 10$  % relative humidity. Males and females were allowed to mate 4 times and the eggs laid were separated. Freshly hatched larvae were fed with mosquito larvae (LECLAIR et al., 1986) till they became ready for pupation. After a preliminary bioassay experiment (JOSE, 1986) different sublethal concentrations viz. 0.001 % (low) and 0.003 % (high) of Diflubenzene were prepared in distilled water and were topically applied to 2 of the 3 series of isolated, pre-weighed, healthy larvae (test individuals series II & III), while series I individuals were treated with distilled water alone. After the completion of metamorphosis the adult males and females were separated, weighed and allowed to mate in nine combinations: with control male (CM) and female (CF), low concentration-treated males (LM) with females (LF) and high concentration treated male (HM) with female (HF). The experimental setup was maintained in triplicate for statistical analysis (ZAR, 1974). For each combination, fecundity (egg-output) and egg weight were recorded. The eggs were then allowed to hatch in laboratory conditions (SEN, 1979) and the weights of newly hatched young ones were measured. The abdomen of postoviposited females was dissected out and the eggs in the ovary was counted.

### **Results and Discussion**

Total number of egg output was maximum for CF mated with CM which was significantly reduced in all the test individuals treated with either low or high concentration of Diflubenzene (Table 1). Topical application also resulted in a decrease in hatching rate of eggs (expressed in percentage); the range of decrease was also found to be of a significantly high one (95 %–22 %). Moreover, the eggs retained in the ovary was also reduced from an average value of 22 (CF × CM) to an average value of 7 (HF × HM) indicating an inhibitory effect in ovicidal activity and/or vitellogenesis (Table 1).

Similar reports on reduction in fecundity and percentage of hatching were reported in insects due to applications of moulting inhibitors (SAXENA & KUMAR, 1982; CHOCKALINGAM & NOORJE-

HAN, 1984), Pesticides (Kuribayashi, 1981) and antifeedants (ANSARI & KHAN, 1973, 1978). SINGH et al., 1986 were of the opinion that the insecticides interfer with endocrine process in insects and inhibit their development and fertility (LEWALLEN, 1964). The adverse effects in egg laying and hatching in *C. tri. asiaticus* could be attributed to the interference of toxicants on the synthesis and incorporation of macromolecules in the ovary (HALL et al., 1976; SRINIVASAN & KESHAVAN, 1979) and/or due to inhibition in the synthesis of required neurosecretory hormones (ADAMS, 1974; MC DANIEL & BERRY, 1974).

Increase in concentration of topical application to males and females also resulted in an inevitable weight-loss for the eggs. Table 1 also reveals that either one way – (either male or female) or both way – (both male and female) treatment affects the mating potential of males more significantly than of females, when fecundity and hatching of eggs are concerned. DAVIES, 1965; GILLOTT & FRIEDAL, 1977 and SUBRAMANIUM, 1984 stated that in addition to supplying sperms, mating in insects initiates oogenesis, accelerates egg maturation, stimulates ovulation, promotes oviposition and enhances the total egg output. The present findings conform their statement in the sense that even the low concentration treated individuals (LM or LF) are very conspicuous inhibitors of all those stages of ovicidal activities which culminate into an overall reduced hatching rate for the treated insects.

Correlation coefficient analysis confirmed that the inhibition in egg output and hatching in treated females was highly significant when mating was allowed with treated male (p < 0.01 and 0.001). It also showed that the treatment affected the reproductive potential of the females significantly (egg output: p < 0.05; hatching: p < 0.01). The inhibition of fecundity and fertility of the eggs by the toxicant showed a clear dosedependency and is very similar to the results obtained in  $Musca\ domestica\ (Srinivasan\ & Keshavan\ ,1979)$  and  $Bombyx\ mori\ (Haniffa\ et\ al., 1986)$ . The weight-loss of eggs as well as freshly emerged larvae is again dose-dependent and the efficacy of the toxicant (Diflubenzene) is equally exhibited in both male and female (p < 0.01; Table 1).

# Acknowledgements

The authors express their deep sense of gratitude to Prof. P. N. Mehrotra, Head, Department of Zoology, Ranchi University, Ranchi, India for facilities and encouragement. The work was financed by University Grants Commission, New Delhi, India in the form of a minor research project to one of us (NSS).

### References

- ADAMS, T. S. (1974): The role of juvenile hormone in house fly ovarian follicle morphogenesis. J. Insect. Physiol. 20: 263–276.
- Ansari, H. J. & Khan, M. A. (1973): Effect of triphenyltin acetate on the fecundity and fertility of Dysdercus cingulatus F. (Heteroptera: Pyrrhocoridae), Curr. Sci. 42: 280–281.
- Ansari, H. J. & Khan, M. A. (1979): Effect of triphenyltin acetate on the fertility of male and cotton bug, Dysdercus cingulatus F. (Heteroptera, Pyrrhocoridae). Comp. Physiol. Ecol. 4: 121–123.
- Büchi, R. (1978): Ovicidal and larvicidal action of Dimilin against the European corn-borer, Ostrinia nubailabis. Hl. Zeitsch. Angewandt. Entomol., 86: 67–71.
- CHOCKALINGAM, S. & NOORJEHAN, A. (1984): The ovicidal effect of Diflubenzene on hemipteran bugs, Dysdercus cingulatus and Chrysocoris purpureus. Curr. Sci., 53: 1112–1113.
- DANIEL, C. N. & BERRY, S. J. (1974): Effects of caffeine and aminophylline on adult development of the Cecropia, silkmoth. J. insect. Physiol. 20: 245–252.
- DATTAMUNSHI, J. S., BILGRAMI, K. S., SAHA, M. P. & JOHAR, M. A. O. (1979): Survey of fish-pawn collection grounds in the river Ganges. Environ. Conserv., 6 (2): 153–157.
- GILLOT, C. & FRIEDAL, T. (1977): Fecundity enhancing and receptivity inhibiting substances produced by male insects. In: Advances of invertebrate reproduction. Vol.1 (Eds. Adiyodi, K. S. & Adiyodi, R. G.) ISR, Peralamkenath, Kerala, India, p. 199–218.
- HANIFFA, M. A., DEVRAJ, M., MURUGESAN, A. G. & THOMSA, P. M. (1986): Inhibition of pupal development and egg production in the silkworm Bombyx mori following application of insecticide and secondary plant substances. Mitt. Zool. Mus. Berl., 62: 337–342.
- Haniffa, M. A. & Jose, S. S. (1987): Inhibition of egg production and fertility in the house cricket, Gryllodes sigillatus Walker (Orthoptera, Gryllidae) following topical application of cythion. Anz. Schädlingskde, 60: 136–138.
- HOLST, H. (1975): Effect of new insecticide PDD-6040 on the fertility of Epilachna varivestis Muls. (Coleoptera, Coccinellidae) & Leptinoterga decemlineata Say (Coleoptera, Chrysomelidae). Zeitsch. Pflanz. Pflanzenschutz., 82 (1): 8–12.
- Jose, S. S. (1986): Effects of DDT and cythion on preoviposition period, egg output, hatching, egg weight and first instar weight of the house cricket Gryllodes sigillatus (Walker). M.Sc. Dissertation, Madurai Kamraj Univ., Tamilnadu, India.

- KURIBAYASHI, S. (1981): Studies on the effect of pesticides on the reproduction of the silkworm Bombyx mori L. III. Ovicidal actions of organomercuric compounds administrated during the larval stages. J. Sericult. Sci. Japan, 51: 167–175.
- LECLAIR, R. JR., ALARIE, Y. & BOURASSA, J. P. (1986): Prey choice in larval Dytiscus harrisii Kirby and D. verticalis Say (Coleoptera, Dytiscidae). Entomologica Basiliensia, 11: 337–342.
- Lewallen, L. L. (1964): *Effect of farnesol and ziram on mosquito larvae*. Mosquito. News., 24: 43–45.
- MATHAVAN, S., PREMLATHA, Y. & CHRISTOPHER, M. S. M. (1985): Effects of caffeine and theophylline on the fecundity of four lepidopteran species. Exp. Biol. 44: 44–51.
- Pant, R. & Katiyar, S. K. (1983): Effects of malathion and acetylcholine on the developing larvae of Philosomia ricini (Lepidoptera, Saturniidae). J. Biosci. 5: 89–95.
- SAXENA, S. C. & KUMAR, V. (1982): Effect of two chitin synthesis inhibitors on reproduction of Trogoderma granarium Entomon. 7: 141–144.
- SEN, N. S. (1979): *Ecological studies of some aquatic insects of Ranchi*. Ph.D. Thesis, Ranchi University, Ranchi, India, p. 24–36.
- SINGH, S. P., BHATNAGAR, M. S., CHANDRA, R. & GANGULY, S. K. (1985): *Thiourea* as a growth inhibitor: Cockroach, Periplaneta americana L. J. Environ. Biol., 6: 169–178.
- Srinivasan, A. & Kesavan, P. S. (1979): Effect of caffeine on longivity and reproduction of housefly. Toxicol. Lett., 3: 101–105.
- Subramaniam, M. (1984): Studies on the male accessory reproductive glands and mateing physiology in the house cricket Gryllodes sigillatus W. (Orthoptera, Gryllidae). Ph.D. Thesis Univ. of Madras, Madras, India, p. 49–56.
- ZAR, J. H. (1974): Biostatistical analysis. Prentice-Hall Inc. N. J. 600-620 p.

### Author's Address:

Dr. Nisarga Sundar Sen, Associate Professor in Zoology Mr. Azfar Hussain Naqvi, Research Associate Mrs. Vineeta Rani, Research Associate Department of Zoology, Ranchi University, Ranchi 834 008, Bihar, India.

Table 1. Showing the effect of topical application of Diflubenzene on the fecundity and ovidical activities in C. tri. asiaticus

Mating Male	Pairing Female	Number of eggs laid	Percentage reduction in Hating (%)	Number of eggs retained	Egg weight (mg)	First instar weight (mg)		
C C C L L H H	C L H C L H C L	$\begin{array}{c} 458 \pm 9.1 \\ 352 \pm 9.3 \\ 265 \pm 8.0 \\ 380 \pm 21.6 \\ 290 \pm 32.7 \\ 242 \pm 28.7 \\ 321 \pm 16.6 \\ 166 \pm 5.6 \\ 102 \pm 6.0 \end{array}$	$4.8 \pm 0.66$ $10.7 \pm 2.64$ $21.3 \pm 1.99$ $28.2 \pm 2.81$ $43.4 \pm 3.21$ $52.1 \pm 5.11$ $44.3 \pm 1.51$ $57.9 \pm 2.65$ $87.9 \pm 4.57$	$22.7 \pm 1.73$ $18.9 \pm 3.51$ $14.6 \pm 3.19$ $19.0 \pm 2.06$ $13.7 \pm 1.63$ $8.7 \pm 1.36$ $14.7 \pm 1.15$ $10.0 \pm 1.73$ $7.0 \pm 2.0$	$\begin{array}{c} 0.66 \pm 0.03 \\ 0.60 \pm 0.01 \\ 0.58 \pm 0.02 \\ 0.62 \pm 0.02 \\ 0.60 \pm 0.01 \\ 0.57 \pm 0.03 \\ 0.60 \pm 0.03 \\ 0.53 \pm 0.02 \\ 0.51 \pm 0.05 \end{array}$	$\begin{array}{c} 0.74  \pm  0.021 \\ 0.70  \pm  0.016 \\ 0.71  \pm  0.026 \\ 0.73  \pm  0.028 \\ 0.73  \pm  0.018 \\ 0.71  \pm  0.116 \\ 0.74  \pm  0.012 \\ 0.72  \pm  0.015 \\ 0.70  \pm  0.0126 \\ \end{array}$		
CORRELATION COEFFICIENT								
Male Female	P P	< 0.01 < 0.05	< 0.001 < 0.01	_ _	< 0.01 < 0.01	< 0.01 < 0.05		

C = control, L = low concentration treated, H = high concentration treated,

P < 0.05 = significant, < 0.01 = highly significant.