

**Zeitschrift:** Entomologica Basiliensia  
**Herausgeber:** Naturhistorisches Museum Basel, Entomologische Sammlungen  
**Band:** 19 (1996)

**Artikel:** Taxonomic revision of the genus Hydrovatus Motschulsky (Coleoptera, Dytiscidae)

**Autor:** Biström, O.

### Inhaltsverzeichnis

**DOI:** <https://doi.org/10.5169/seals-980453>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 10.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Taxonomic Revision of the Genus *Hydrovatus* Motschulsky (Coleoptera, Dytiscidae)

(Contribution to the study of Dytiscidae 55)

by **O. Biström**

**Abstract:** This revision of the genus *Hydrovatus* Motschulsky, 1853 is based on examination of adult specimens. In all, 202 species are tentatively recognized in the genus (the status of some species remains unclear). A new classification based on a character analysis is introduced, dividing the genus into 15 species groups. The phylogeny of Hydrovatini is discussed; the tribe is regarded as monophyletic. Two monophyletic genera are associated with Hydrovatini: *Queda* Sharp and *Hydrovatus*. A combination of Methlini and *Hydrovatus* excluding *Queda* is rejected. Diagnoses and descriptions for all recognized taxa are given together with keys for their identification. Known distribution and biological data are given for all species. A brief zoogeographical survey indicates that the place of origin of Hydrovatini and its genera is Gondwanaland, 100 million years ago or more, before the opening of the South Atlantic rift.

New taxa are described as follows: *Hydrovatus hintoni* n.sp. (Mexico); *H. kavanaughi* n.sp. (Mexico); *H. youngi* n.sp. (Mexico); *H. longior* n.sp. (Mexico); *H. asemus* n.sp. (Uganda, Zaire); *H. visendus* n.sp. (Tanzania, Zimbabwe); *H. fernandoi* n.sp. (Angola); *H. eximius* n.sp. (Zimbabwe, Mozambique); *H. angusticornis* n.sp. (Angola); *H. imitator* n.sp. (Zaire); *H. nilssoni* n.sp. (Gambia, Senegal, Congo, Angola); *H. unguicularis* n.sp. (Uganda, Sudan); *H. tydaeus* n.sp. (Sudan, Ethiopia); *H. maai* n.sp. (Malaysia, Indonesia); *H. similis* n.sp. (Philippines); *H. balfourbrownii* n.sp. (Malaysia); *H. laosensis* n.sp. (Laos); *H. saundersi* n.sp. (Malaysia, Singapore); *H. pisiformis* n.sp. (Thailand, Singapore); *H. latipalpis* n.sp. (Zaire); *H. roccii* n.sp. (Guinea Bissau); *H. irianensis* n.sp. (Indonesia); *H. schawalleri* n.sp. (Tanzania); *H. mollis* n.sp. (Sudan); *H. weiri* n.sp. (Australia); *H. navigator* n.sp. (Philippines); *H. cessatus australis* n. ssp. (Angola); *H. vulpinus* n.sp. (Zimbabwe); *H. unguiculatus* n.sp. (Ghana), *H. witteti* n.sp. (Zaire); *H. vulneratus* n.sp. (Nigeria, Zaire); *H. soror* n.sp. (Tanzania, Zimbabwe); *H. stridulus* n.sp. (Thailand, Malaysia, Singapore); *H. samuelsoni* n.sp. (Indonesia); *H. enigmaticus* n.sp. (Indonesia); *H. jaechi* n.sp. (Indonesia); *H. agathodaemon* n.sp. (Indonesia).

New synonymies are established as follows (first valid name): *H. pictulus* Sharp, 1882 = *H. scymnoides* Régimbart, 1895; *H. subrotundatus* Motschulsky, 1859 = *H. carbonarius* (Clark, 1863), *H. carbonarius* v. *fuscobrunneus* (Clark, 1863), *H. ferrugatus* Régimbart, 1877, *H. elevatus* Sharp, 1882, *H. orientalis* Sharp, 1882 and *H. javanus* Csiki, 1938; *H. leconteii* (Clark, 1862) = *H. inornatus* Sharp, 1882; *H. pustulatus* (Melsheimer, 1846) = *H. indianensis* Blatchley, 1910; *H. parvulus* Régimbart, 1899 = *H. noctivagus* Guignot, 1953, *H. socors* Guignot, 1954 and *H. ocnerus* Guignot, 1958; *H. granosus* Guignot, 1958 = *H. similaris* Bilardo & Rocchi, 1987; *H. confossus* Guignot, 1958 = *H. platycerus* Bilardo & Pederzani, 1978; *H. flammulatus* Sharp, 1882 = *H. gravicornis* Omer-Cooper, 1957; *H. absonus* Guignot, 1948 = *H. lachnaeus* Guignot, 1958 and *H. absonus* *borkuanus* Bruneau de Mirè & Legros, 1963; *H. abraeoides* Régimbart, 1895 = *H. metrius* Guignot, 1961; *H. glaber* Guignot,

1953 = *H. katangensis* Guignot, 1958 and *H. medioximus* Omer-Cooper, 1963; *H. spissicornis* Régimbart, 1904 = *H. procus* Guignot, 1942; *H. madagascariensis* Régimbart, 1903 = *H. lophotus* Guignot, 1948; *H. cibratus* Sharp, 1882 = *H. laticornis* Régimbart, 1895 and *H. dyscheres* Guignot, 1955; *H. longicornis* Sharp, 1882 = *H. berdoa* Bruneau de Miré & Legros, 1963; *H. globulosus* Gschwendtner, 1943 = *H. pauxillus* Gschwendtner, 1943; *H. pudicus* (Clark, 1863) = *H. pusillus* Régimbart, 1881; *H. gabonicus* Régimbart, 1895 = *H. pravus* Guignot, 1954; *H. difficilis* Guignot, 1956 and *H. discifer* Guignot, 1958; *H. ovalis* Sharp, 1882 = *H. armstrongi* Watts, 1978; *H. collega* Guignot, 1955 = *H. rutilus* Omer-Cooper, 1963; *H. acuminatus* Motschulsky, 1859 = *H. obscurus* Motschulsky, 1859, *H. sordidus* Sharp, 1882, *H. humilis* Sharp, 1882, *H. affinis* Régimbart, 1895 and *H. obscurus* Régimbart, 1895; *H. nimbaensis* Guignot, 1954 = *H. ornatus* Guignot, 1958; *H. parallelipennis* Régimbart, 1895 = *H. histeroides* Régimbart, 1899; *H. seminarius* Motschulsky, 1859 = *H. fusculus* Sharp, 1882, *H. tinctus* Sharp, 1882 and *H. matsuii* Nakane, 1990; *H. cuspidatus* (Kunze, 1818) = *H. simplex* Sharp, 1882; *H. insolitus* Guignot, 1948 = *H. propinquus* Guignot, 1955, *H. cryptus* Guignot, 1956, *H. tritus* Guignot, 1959 and *H. nachtigali* Bruneau de Mirré & Legros, 1963; *H. castaneus* Motschulsky, 1855 = *H. reticulatus* Zimmermann, 1924 and *H. bironicus* Guignot, 1956; *H. obsoletus* Peschet, 1922 = *H. adelphus* Guignot, 1956; *H. simoni* Régimbart, 1894 = *H. glomeratus* Guignot, 1942 and *H. consimilis* Omer-Cooper, 1957; *H. regimbarti* Zimmermann, 1919 = *H. laeviusculus* Gschwendtner, 1943 and *H. subsimilis* Guignot, 1958; *H. facetus* Guignot, 1942 = *H. continentalis* Guignot, 1957 and *H. quezeli* Bruneau de Miré & Legros, 1963; *H. reticuliceps* Régimbart, 1895 = *H. liberatus* Guignot, 1945 and *H. compotor* Guignot, 1955; *H. impunctatus* Guignot, 1953 = *H. cunctator* Guignot, 1958, *H. vagetinctus* Guignot, 1958 and *H. pallidus* Bilardo & Pederzani, 1979; *H. contumax* Guignot, 1954 = *H. rohani* Guignot, 1959; *H. villiersi* Guignot, 1955 = *H. albertianus* Guignot, 1959, *H. uncus* Guignot, 1959 and *H. browneanus* Guignot, 1959; *H. pulcher* Gschwendtner, 1934 = *H. legroisi* Guignot, 1954; *H. badeni* Sharp, 1882 = *H. cicer* Guignot, 1942; *H. obtusus* Motschulsky, 1855 = *H. acutus* Sharp, 1882.

Lectotypes are designated for many taxa.

*Hydrovatus pasiricus* Csiki, 1938 and *H. p.* var. *simplicior* Csiki, 1938 and *H. p.* var. *unicolor* Csiki, 1938 should not be placed in *Hydrovatus* as here defined. Possibly they should be placed somewhere close to *Hyphydrini*. The taxonomic status of a few *Hydrovatus* species remains unclear because no material have been available for study.

**Key words:** Coleoptera Dytiscidae - *Hydrovatus* - revision - description - new species.

## Contents

<b>1. Introduction . . . . .</b>	59
<b>2. Material and methods . . . . .</b>	60
<b>3. Monophyly of Hydrovatini and its genera . . . . .</b>	65
3.1. Character distribution and discussion . . . . .	65
3.2. Conclusions . . . . .	75

<b>4. Subdivision of <i>Hydrovatus</i></b>	76
4.1. Historical review	76
4.2. An attempt of subdivision and reconstruction of the phylogeny	77
<b>5. Zoogeographical considerations</b>	83
<b>6. Classification and descriptions</b>	85
6.1. Genus <i>Hydrovatus</i> Motschulsky	85
6.2. Special taxonomic cases	91
6.3. Key to the species groups of <i>Hydrovatus</i>	94
6.4. Key to the species	96
6.5. Descriptions of the species	125
6.5.1. Species group 1 (sp. gr. <i>pictulus</i> )	125
6.5.2. Species group 2 (sp. gr. <i>turbinatus</i> )	133
6.5.3. Species group 3 (sp. gr. <i>pustulatus</i> )	134
6.5.4. Species group 4 (sp. gr. <i>confossus</i> )	172
6.5.5. Species group 5 (sp. gr. <i>balfourbrownei</i> )	310
6.5.6. Species group 6 (sp. gr. <i>pumilus</i> )	314
6.5.7. Species group 7 (sp. gr. <i>gabonicus</i> )	327
6.5.8. Species group 8 (sp. gr. <i>roccii</i> )	334
6.5.9. Species group 9 (sp. gr. <i>acuminatus</i> )	336
6.5.10. Species group 10 (sp. gr. <i>parallelipennis</i> )	359
6.5.11. Species group 11 (sp. gr. <i>oblongipennis</i> )	368
6.5.12. Species group 12 (sp. gr. <i>villiersi</i> )	480
6.5.13. Species group 13 (sp. gr. <i>pulcher</i> )	494
6.5.14. Species group 14 (sp. gr. <i>confertus</i> )	521
6.5.15. Species group 15 (sp. gr. <i>clypealis</i> )	551
<b>7. Checklist of the genus <i>Hydrovatus</i></b>	555
<b>8. References</b>	558
<b>9. Index to species group names</b>	577
<b>Appendix</b>	580

## 1. Introduction

With this revision of the diving-beetle genus *Hydrovatus* Motschulsky I conclude my survey of the taxonomy and systematics of the hydroporine tribe Hydrovatini.

The traditional composition of Hydrovatini, introduced by SHARP (1882a), includes two genera, viz. *Hydrovatus* and *Queda* Sharp. A separate revision of the latter genus was published by BISTRÖM (1990). A reorganisation of the Hydrovatini has recently been sug-