

Zeitschrift: Entomologica Basiliensia
Herausgeber: Naturhistorisches Museum Basel, Entomologische Sammlungen
Band: 19 (1996)

Artikel: Taxonomic revision of the genus Hydrovatus Motschulsky (Coleoptera, Dytiscidae)
Autor: Biström, O.
Anhang: Appendix
DOI: <https://doi.org/10.5169/seals-980453>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

- *savanicola* Guignot ... 382
schawalleri n.sp ... 348
scholaeus Guignot ... 194
**scymnoides* Régimbart ... 126
seminarius Motschulsky ... 368
semirufus Zimmermann ... 312
senegalensis Régimbart ... 198
seydeli Guignot ... 478
**sharpī* Régimbart ... 444
sharpī Van den Branden ... 145
**similaris* Bilardo & Rocchi ... 179
similis n.sp ... 306
simoni Régimbart ... 441
**simplex* Sharp ... 373
singularis Guignot ... 93
sinister Sharp ... 521
sitistus Omer-Cooper ... 496
sobrinus Omer-Cooper ... 435
**socors* Guignot ... 174
**sordidus* Sharp ... 352
soror n.sp ... 514
spadix Guignot ... 178
spissicornis Régimbart ... 252
sporas Guignot ... 454
stappersi Guignot ... 432
**straeleni* Guignot ... 402
stridulus n.sp ... 529
subparallelus Gschwendtner ... 263
**subpunctatus* Kolbe ... 507
subrotundatus Motschulsky ... 137
**subsimilis* Guignot ... 444
subtilis Sharp ... 526
sumatrensis Sharp ... 547
suturalis Bilardo & Pederzani ... 438
tardiosus Guignot ... 93
testudinarius Régimbart ... 467
**tinctus* Sharp ... 369
tristis Guignot ... 261
**tritus* Guignot ... 377
turbinatus Zimmermann ... 133
**turgidus* Guignot ... 482
tydaeus n.sp ... 280
**uncus* Guignot ... 490
unguicularis n.sp ... 273
unguiculatus n.sp ... 481
uniformis (Fairmaire) ... 516
**vagetinctus* Guignot ... 472
validicornis Régimbart ... 254
verisae Bilardo & Rocchi ... 259
vicus Guignot ... 172
villiersi Guignot ... 489
visendus n.sp ... 219
vividus Guignot ... 368
vulneratus n.sp ... 505
vulpinus n.sp ... 480
weiri n.sp ... 399
wittei n.sp ... 500
youngi n.sp ... 165

O. Biström
 Zoological Museum
 University of Helsinki
 P.O. Box 17
 FIN-00014 Helsinki, Finland.

Appendix

During the final preparation of the revision of *Hydrovatus* Motschulsky, published in this volume, an additional undescribed species was detected. Predominantly for practical reasons, the new species is described here.

***Hydrovatus perssoni* n.sp. Biström & Nilsson**

Type locality: Cacheu, 5 km W Bula, Guinéa-Bissau.

Type material (2 m, 6 f): Holotype, m: Guina-Bissau: 1, Cacheu: 5 km W Bula 19/7-92 S. Persson/*Hydrovatus* sp. *sordidus*-gr. Det. A N Nilsson-92 (Zool. mus.


Lund). – Paratypes: Same as holotype (1 ex. Zool. mus. Helsinki, 1 ex. coll. Nilsson); same as holotype but 19/7–92 (1 ex. Zool. mus. Helsinki, 2 exx. coll. Persson, 2 exx. coll. Nilsson); same as holotype but 31.VII.93/*Hydrovatus perssoni* sp.n. det. AN Nilsson –93 (1 ex. Zool. mus. Helsinki, 3 exx. coll. Nilsson & Persson); same as holotype but 4.VII.93/*H. perssoni* sp.n. det. Nilsson –93 (2 exx. Zool. mus. Helsinki; 5 exx. coll. Nilsson & Persson); same as holotype but 22.VII. 93/*H. perssoni* sp.n. det. Nilsson -93 (2 exx. coll. Nilsson & Persson); same as holotype but 12.VII.93/*H. perssoni* sp.n. det. Nilsson -93 (1 ex. coll. Nilsson & Persson); *Hydrovatus* (s.str.) nov. spec./Guinea Bissau Ilha de Bolama: Bolama; Vassado; Pte de Preco Leve; Rice-Field 7.XI.1992 leg. & det. J. Cuppen (1 ex. coll. Cuppen, the Netherlands).

Etymology: Named after Mr. Sven Persson, Sweden, who collected the type material.

Diagnosis and discussion: In the key to species groups (p. 94), the new species fits the species group 9 (sp.gr. *acuminatus*). In the key to species of species group 9 (p. 111), *H. perssoni* fits point 7, at which *H. schawalleri* and *H. collega* are distinguished. From both species, *H. perssoni* is easily separated by the penis, which is provided with a distinct medial constriction (*H. schawalleri* and *H. collega* lack similar medial constriction, see p. 349 and 346).

Description: only diagnostically important differences from redescription of *H. collega* are recognized.

Length of body: 2.32–2.56 mm, breadth: 1.44–1.64 mm. Habitus in Fig. 1.



Figs 1–6: *Hydrovatus perssoni*. – 1, habitus. – 2, head, frontal aspect. – 3, male antenna. – 4, penis, dorsal aspect. – 5, penis, lateral aspect. – 6, apical part of paramere. Horizontal scale 0.5 mm, head and antenna; left scale 1 mm, habitus; right scale 0.4 mm, genitalia.

Head: Ferruginous, frontally generally slightly paler; pale ferruginous. Head frontally narrowly margined; frontal outline medially straight or indistinctly concave (Fig. 2). Antenna slender (Fig. 3).

Pronotum: Ferruginous, laterally with vague paler areas and mediobasally with a narrow, vague, darkened area. Very finely microsculptured; laterally on disc meshes of microsculpture hardly discernible.

Elytra: Dark ferruginous. Laterally and posteriorly elytron becomes gradually paler; at epipleuron pale ferruginous. Without distinct colour pattern.

Male genitalia: Figs 4–6.

Female: Rather indistinctly dimorphous. Externally almost as male, or body-reticulation, although fine, slightly more distinct than in male.

Distribution: Guinea Bissau.

Biology: The new species was collected in two adjacent ponds with sparse marginal vegetation. The ponds were situated in more or less open land and were apparently permanent. Adults were found on 19th and 29th July, whereas subsequent visits in August and September produced no additional specimens.

Table 1. Distribution of relevant characters in taxa. Left vertical column with running numbers 1–19 indicating characters (cf. corresponding numbers in text). Horizontal top column indicating taxa (see below). 1, character represented only by plesiomorphous state; 2 character present both in plesiomorphous and apomorphous state; 3, character present only in apomorphous state. * special cases with further discussion in text. ?, status of character unclear. -, character not examined.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
1	2	1	2	2	1	1	1	?	1	1	1	1	1	1	1	1	1	1	1	1	1	2	1	?	1
2	1	1	3	3	1	2	2	1	1	3	3	3	1	1	1	1	1	1	1	1	1*	1	1	1	1
3	1	1	3	3	1	3	2	3	3	3	3	1	1	1	1	1	1	1	1	1	1	1	1	1	1
4	1	3	3	3	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	3*
5	1	1	3	3	1	1	1	1	1	1*	1*	1	1	1	1	1	1	1	1	1	1*	1*	1	1	1
6	1	1	1	3	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
7	1	1	1	1	1	1	3	3	3	3	3	1	1	1	1	1	1	1	1	1	1	1	1	1	1
8	1	3	2	3	3	3	3	3	3	3	3	3	1	1	1	1	1	1	1	1	1	1	1	?	1
9	1	3	3	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
10	1	3	1	1	3	?	?	?	?	3	3	3	1	1	1	1	1	1	1	1	1	1	3	3	?
11	3	3	1	1	3	3	2	1	?	1	1	1	1	1	1	1	1	?	1	1	1	1	?	?	?
12	3	3	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
13	1	3	1	1	1	1	1	—	3	3	3	3	1	1	3	3	3	3	3	3	1	1	1	1	1
14	1	3	3	1	3	1	2	1	1	1	1	1	1	1	1	?	?	?	?	1	1	1	1	1	3
15	2	1	2	2	1	1	1	?	?	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
16	1	3	1	3	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	3	1	2	1	3
17	1	1	1	1	1	!	3	1	1	1	1	1	1	1	1	?	1	1	1	1	1	?	1	1	1
18	2	3	2	1	1	1	2	1	1	1	1	2	1	1	1	1	1	1	1	3	1	1	1	1	1
19	2	1	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1

Taxa examined: 1, *Laccornis* – 2, *Methles*. – 3, *Hydrovatus*. – 4, *Queda*. – 5, *Derovatellus*. – 6, *Herophydrus*. – 7, *Bidessini*. – 8, “*Hydrovatus pasiricus*”. – 9, *Microdytes*. – 10, *Heterhydrus*. – 11, *Pachydrus*. – 12, *Hyphydrus*. – 13, *Platambus*. – 14, *Agabus*. – 15, *Ilybius*. – 16, *Colymbetes*. – 17, *Graphoderes*. – 18, *Acilius*. – 19, *Eretes*. – 20, *Dytiscus*. – 21, *Copelatus*. – 22, Noteridae. – 23, Haliplidiae. – 24, Gyrinidae. – 25, Hygrobiidae.

Table 2. Distribution of relevant characters within genus *Hydrovatus*. Left vertical column indicating characters (A-P cf. text). Horizontal columns indicating state of character in species groups distinguished (1–15) and *Queda* = Q. 1, character present only in plesiomorphous state; 2, character present both in plesiomorphous and apomorphous state; 3, character present only in apomorphous state. * special cases with further discussion in text.